

Ecological site R111XE002OH Limnic Muck

Last updated: 5/28/2020 Accessed: 05/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 111X–Indiana and Ohio Till Plain

111E – Indiana and Ohio Till Plain, Eastern Part. Most of this area is in the Till Plains Section of the Central Lowlands Province of the Interior Plains. The northeast tip of the area is in the Southern New York Section of the Appalachian Highlands. The entire area has been glaciated. It is dominated by ground moraines that are broken in places by kames, lake plains, outwash plains, terraces, and stream valleys. Narrow, shallow valleys commonly are along the few large streams in the area. Elevation ranges from 580 to 1,400 feet (175 to 425 meters), increasing gradually from west to east. Relief is mainly a few meters, but in some areas hills rise as much as 100 feet (30 meters) above the adjoining plain.

The extent of the major Hydrologic Unit Areas (identified by four-digit numbers) that make up this MLRA is as follows: Scioto (0506), 33 percent; Muskingum (0504), 31 percent; and Western Lake Erie (0410), 28 percent; Upper Ohio (0503), 5 percent; and Southern Lake Erie (0411), 3 percent. The headwaters of many rivers in central Ohio, including the Vermillion, Black Fork, Sandusky, Little Scioto, and Olentangy Rivers, are in this MLRA.

This MLRA is underlain by late Devonian shale and sandstone. Surficial materials include glacial deposits of till, glaciolacustrine sediments, and outwash from Wisconsin and older glacial periods.

Classification relationships

Major Land Resource Area (USDA-Natural Resources Conservation Service, 2006)

USFS Ecological Regions (USDA, 2007): Sections –Central Till Plains, Beech Maple (222H), Western Glaciated Allegheny Plateau (221F)

Subsections – Allegheny Plateau (221Fa), Bluffton Till Plains (222Ha), Miami-Scioto Plain – Tipton Till Plain (222Hb)

NatureServe Systems anticipated (NatureServe, 2011): Agriculture - Cultivated Crops and Irrigated Agriculture, Agriculture – Pasture/Hay, North-Central Interior Beech-Maple Forest,

LANDFIRE Biophysical Settings anticipated (USGS, 2010): Central Interior and Appalachian Floodplain Systems, North-Central Interior Beech-Maple Forest

Ecological site concept

This site is an upland site formed on herbaceous organic soil parent material overlaying coprogenous material in depressions on lake and till plains. Disturbance to the site by fire was driven by the seasonal presence of water on the site. In the spring and early summer the site is often ponded to a depth of 15 inches and for up to 30 days. The water table recedes as the summer progresses creating a drier situation. The drier conditions that can occur in very

early spring or late fall, coupled with the then dormant vegetation, increases the susceptibility of the site to fire. The seasonal flooding and relatively short fire return interval (~10 years or less) created a situation where the site is dominated by herbaceous vegetation.

The characteristic vegetation is herbaceous in nature, mostly dominated by tall prairie grass species, most notably big bluestem and eastern gamagrass. Sedge species are also common throughout the site and are the most abundant sub-dominant group of species. The influence of water via flooding and ponding along with fire maintain the herbaceous species dominance of the site. The absence of fire will lead this site to being converted to woodland and then potentially a closed canopy forest. The lack of fire will allow hydrophilic shrub species to get established, further altering the fire cycle. Continued absence of fire will have the site progress to a forest dominated by wetland tree species such as elm and ash. Invasive species, specifically, reed canarygrass and phragmites, can invade and come to dominate the site in the absence of management to curtail or prevent their establishment and proliferation. Currently, a large percentage of the site is in agricultural production, notably corn and soybean rotations, after the installation of drainage. These sites can be very productive, once drained, due in large part to the high amounts of organic matter.

Associated sites

R111XE001OH	Mineral Muck Underlain by mineral soil
R111XE003OH	Deep Muck Not underlain by mineral soil
F111XE501OH	Till Depression On till parent material

Similar sites

R111XE001OH	Mineral Muck Underlain by mineral soil
R111XE003OH	Deep Muck Underlying material is not limnic (coprogeneous earth).
R111XE401OH	Wet Outwash Mollisol Soil parent material is outwash; site generally higher on the landscape; soils are mollisols



Figure 1.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) Andropogon gerardii (2) Tripsacum dactyloides

Physiographic features

This site is located in the 111E – Indiana and Ohio Till Plain, Eastern Part MLRA. This site was formed on herbaceous organic material, occasionally with a top layer alluvium. It is located in depressions with slopes ranging from 0 to 2 percent.

Landforms	(1) Depression
Flooding frequency	None
Ponding duration	Long (7 to 30 days)
Ponding frequency	Frequent
Elevation	244–427 m
Slope	0–2%
Ponding depth	0–38 cm
Water table depth	15 cm
Aspect	Aspect is not a significant factor

Table 2. Representative physiographic features

Climatic features

The average annual precipitation in this area is 35 to 41 (890 to 1,040 millimeters). Most of the rainfall occurs as convective thunderstorms during the growing season. About half or more of the precipitation occurs during the freeze-free period. Snowfall is common in winter. The average annual temperature is 48 to 52 degrees F (9 to 11 degrees C). The freeze-free period averages about 185 days and ranges from 165 to 205 days.

Frost-free period (characteristic range)	131-150 days
Freeze-free period (characteristic range)	172-181 days
Precipitation total (characteristic range)	991-1,041 mm
Frost-free period (actual range)	127-153 days
Freeze-free period (actual range)	167-185 days
Precipitation total (actual range)	991-1,092 mm
Frost-free period (average)	142 days
Freeze-free period (average)	177 days
Precipitation total (average)	1,016 mm

Table 3. Representative climatic features



Figure 2. Monthly precipitation range



Figure 3. Monthly minimum temperature range



Figure 4. Monthly maximum temperature range



Figure 5. Monthly average minimum and maximum temperature



Figure 6. Annual precipitation pattern



Figure 7. Annual average temperature pattern

Climate stations used

- (1) MANSFIELD LAHM MUNI AP [USW00014891], Mansfield, OH
- (2) COLUMBUS PORT COLUMBUS INTL AP [USW00014821], Columbus, OH
- (3) LANCASTER [USC00334403], Lancaster, OH
- (4) ASHLAND 2 SW [USC00330256], Ashland, OH
- (5) BUCYRUS [USC00331072], Bucyrus, OH
- (6) GALION WTR WKS [USC00333021], Galion, OH

Influencing water features

These wetland systems are groundwater-dependent as well as being, generally, the lowest point in the landscape, making them a discharge depression. Water levels fluctuate seasonally, reaching their peak in spring and lows in late summer. Ponding can occur on the site, generally in the spring, to a depth of 15 inches and can have a very long duration (> 30 days). Water levels typically remain at or near the soils surface throughout the year. The hydrogeographic model classification for this site is DEPRESSION: Muck, Ground Water Influenced; coprogenous/herbaceous. This site has a Cowardin Classification of PEM1Eg; it is a Palustrine system with persistent, emergent vegetation that is seasonally flooded/saturated on organic soil.

Soil features

The soil series associated with this site are: Olentangy and Muskego. They are very deep, very poorly drained, and slow to moderately rapid permeable soils, with very acidic to moderately alkaline soil reaction, that formed in coprogenous material.

Parent Materials Kind: coprogenous material Surface Texture: silt loam Subsurface Texture group: loamy



Figure 8. map showing mapunit location in the MLRA

Table 4.	Representative	e soil features
10010 11	1100100011101111	oon routaroo

Parent material	(1) Coprogenic material
Surface texture	(1) Silt Ioam
Drainage class	Very poorly drained
Permeability class	Slow to moderately rapid
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (Depth not specified)	22.86–33.02 cm
Calcium carbonate equivalent (Depth not specified)	0–70%
Electrical conductivity (Depth not specified)	0 mmhos/cm
Sodium adsorption ratio (Depth not specified)	0
Soil reaction (1:1 water) (Depth not specified)	4.8–7.9
Subsurface fragment volume <=3" (Depth not specified)	0–2%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

The historic plant community of the Limnic Muck is a mucky prairie. This site is characterized by the soil parent material being composed of herbaceous organic matter with the subsoil being largely coprogenous, limnic material. The plant community is dominated by tall-grass prairie species along with an assortment of tall sedge species. The site is located in depressional areas often associated with glacial lake plains and till plains. Being the lowest portion of the local landscape, the site is dependent upon groundwater and its fluctuation, which reaches its peak in the spring and low in the late summer. This fluctuation and dominate herbaceous species create a fire dependent system. Replacement fires about every decade and seasonal flooding worked in concert to drive the species dominance and richness of the site. Both mechanisms work in concert to inhibit the establishment and perpetuation of woody species.

State and transition model



Figure 9. STM

Limnic Muck, R111EY002OH

Diagram Legend

T1A	No fire, no woody species management	
T1B	Drainage, site preparation, planting, management	
T1C	Invasion, no management	
T2A	Tree removal, planting, fire	
ТЗА	Drainage removal, planting, fire	
T4A	Chemical/mechanical treatment of invasive species, fire	
2.1A	No fire, no woody species management	
2.2A	Tree removal, no fire	
3.1A	Pasture/forage planting and maintenance	
3.2A	Tillage/no-till planting and management of row crops.	

Figure 10. Legend

State 1 Mucky Prairie

This is the reference or diagnostic plant community for this site. In reference conditions, this site was dominated by tall prairie grasses specifically big bluestem and eastern gammagrass. Sedge species were also a large component of this site, especially upright sedge and hairy sedge. This state was maintained by nearly seasonal flooding and a fire frequency of 10 years. Both mechanisms work in concert to inhibit the establishment and perpetuation of woody species.

Dominant plant species

- big bluestem (Andropogon gerardii), grass
- eastern gamagrass (Tripsacum dactyloides), grass

Community 1.1 big bluestem - Eastern gamagrass

This is the reference or diagnostic plant community for this site. In reference conditions, this site was dominated by tall prairie grasses specifically big bluestem and eastern gammagrass. Sedge species were also a large component of this site, especially upright sedge and hairy sedge.

Dominant plant species

- big bluestem (Andropogon gerardii), grass
- eastern gamagrass (Tripsacum dactyloides), grass

State 2 Fire Suppressed State

This state is characterized by a longer than normal fire return interval or the absence of fire as a disturbance agent. Absence of fire allows for a thick layer of leaf litter to accumulate which suppresses the production of the herbaceous species and allows the woody species to establish and flourish. Similarly, the woody species that colonize this site are generally intolerant of fire. The seasonal flooding and fire work in concert to help prevent woody species establishment. Continued lack of fire allows the site to transition from a shrub/herbaceous site, to a woodland, and eventually to a forest.

Dominant plant species

- dogwood (Cornus), shrub
- willow (Salix), shrub

Community 2.1 shrubs / small trees

This phase is characterized by the establishment and dominance shrubby species and small trees. Common species include dogwood, white sweetmeadow, and willow species.

Dominant plant species

- dogwood (Cornus), tree
- willow (*Salix*), tree

Community 2.2 American elm - blue ash

This phase is characterized by the establishment and dominance tree species that convert the site to a forest. Common species include American elm and blue ash as dominants.

Dominant plant species

- American elm (Ulmus americana), tree
- blue ash (Fraxinus quadrangulata), tree

Pathway P2.1A Community 2.1 to 2.2

Continued absence of fire and/or woody species management will move the site towards phase 2.2.

Pathway P2.2A Community 2.2 to 2.1

Removal of most or all the trees without the application of fire will move the site towards phase 2.1.

State 3 Agricultural State

This site has largely been converted to agricultural use. Most of the historic acres are now in row crop agricultural use. Most common is a corn and soybean rotation of various types. Roughly 8% of the site is used to grow hay or cool season forage and used for grazing. The conversion of the site to agricultural use begins with the installation of a drainage system which most often employs either or both ditching and field tile.

Community 3.1 Row crops (corn - soybeans)

This phase is characterized by row crop agriculture of small grains, primarily corn and soybeans.

Community 3.2 Cool season forage - pasture

This phase is characterized by forage or grazing agriculture. Different mixes of, generally, cool season grasses and forbs, largely clovers, are grown.

Pathway P3.1A Community 3.1 to 3.2

Planting of cool season pasture/forage species and management to maintain them.

Pathway P3.2A Community 3.2 to 3.1

Planting, either by conventional or no-till methods, of row crop. Management that keeps the site in row crop production

State 4 Invaded State

This site is characterized by the establishment of invasive species. Without management to control them, the invasive species come to be the dominant species to the exclusion of nearly all other species.

Community 4.1 phragmites - reed canarygrass

This phase is characterized by the establishment and dominance of invasive species. The two most common for this site are phragmites and reed canarygrass. Most often only one of the two species will come to dominate the site.

Dominant plant species

- reed canarygrass (Phalaris arundinacea), grass
- common reed (*Phragmites australis*), grass

Transition T1A State 1 to 2

No fire or woody species management will transition this site towards the fire suppressed state. This will result in a

loss of herbaceous species production and richness.

Transition T1B State 1 to 3

The installation of drainage either through ditches of field tile or both facilitates the start of the transition to the agriculture state. Planting and management of the selected crop complete the transition.

Transition T1C State 1 to 4

The establishment of invasive species without management, to include the use of fire, move the site towards the invaded state. This results in a loss in species richness of the site.

Restoration pathway R2A State 2 to 1

Removal of trees and other woody species then planting the site to the desired species. Following these actions with fire will help restore the site to state 1.

Restoration pathway R3A State 3 to 1

Removal of drainage system, site preparation, planting of the desired species, and regular application of fire move the site back to State 1.

Restoration pathway R4A State 4 to 1

Chemical and mechanical treatment of the invasive species is the first restoration step. At times biological treatment, to included grazing, can be helpful but is not common in this area. The reapplication of fire after seeding of the appropriate species helps maintain the species dominate in State 1.

Additional community tables

Inventory data references

Site concept developed through expert opinion, review of the literature, and field work.

Other references

Anderson, D. M. 1982. Plant communities of Ohio: A preliminary classification and description. Columbus, OH: Ohio Dept. of Natural Resources, Division of Natural Areas and Preserves.

Betz, R. (1973). The prairies of Indiana. Proceedings of the Fifth Midwest Prairie Conference (pp. 34-31). Ames: Iowa State University.

Federal Geographic Data Committee. 2013. Classification of wetlands and deepwater habitats of the United States. FGDC-STD-004-2013. Second Edition. Wetlands Subcommittee, Federal Geographic Data Committee and U.S. Fish and Wildlife Service, Washington, DC.

Gordon, R. B. 1969. The natural vegetation of Ohio in pioneer days. Columbus: Ohio State University. Homoya, M. A., Abrell, D. B., Aldrich, J. R., & Post, T. W. (1985). The Natural Regions of Indiana. Indiana Academy of Science, 94, 245-269.

Lafferty, M. B. 1979. Ohio's natural heritage. Columbus: Ohio Academy of Science. NatureServe. (2011). An online encyclopedia of life [web application]. NatureServe, Arlington, VA, USA [Online: www. natureserve. org/explorer]. Jackson, Marion T. 1997. The Natural heritage of Indiana. Bloomington: Indiana University Press, published in association with the Indiana Department of Natural Resources and the Indiana Academy of Science.

Transeau, E. (1935). The prairie peninsula. Ecology vol. 16 (3), 423-437.

USDA. (2007). Ecological Subregions: Sections and Subsections for the Conterminous United States. Washington, DC: USDA - Forest Service.

USDA-NRCS. 2008. Hydrogeomorphic Wetland Classification System: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service. Technical Note No. 190–8–76. Washington D.C.

USDA. (2006). Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U. S. Department of Agriculture, Natural Resources Conservation Service. U. S. Department of Agriculture Handbook 296.

USGS. (2010). LANDFIRE Biophysical Settings. Retrieved from http://www.landfire.gov

Whitaker, John O., Charles J. Amlaner, Marion T. Jackson, George R. Parker, and Peter Evans Scott. 2012. Habitats and ecological communities of Indiana presettlement to present. Bloomington: Indiana University Press.

Approval

Chris Tecklenburg, 5/28/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	TYLER STAGGS
Contact for lead author	
Date	05/19/2024
Approved by	Chris Tecklenburg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):

- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: