

## **Ecological site F115XA005IL Bedrock Backslope**

Last updated: 12/30/2024  
Accessed: 01/08/2025

---

### **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### **MLRA notes**

Major Land Resource Area (MLRA): 115X—Central Mississippi Valley Wooded Slopes

This MLRA is characterized by deeply dissected, loess-covered hills bordering well defined valleys of the Illinois, Mississippi, Missouri, Ohio, and Wabash Rivers and their tributaries. It is used to produce cash crops and livestock. About one-third of the area is forested, mostly on the steeper slopes. This area is in Illinois (50 percent), Missouri (36 percent), Indiana (13 percent), and Iowa (1 percent) in two separate areas. It makes up about 25,084 square miles (64,967 square kilometers).

Most of this area is in the Till Plains section and the Dissected Till Plains section of the Central Lowland province of the Interior Plains. The Springfield-Salem plateaus section of the Ozarks Plateaus province of the Interior Highlands occurs along the Missouri River and the Mississippi River south of the confluence with the Missouri River. The nearly level to very steep uplands are dissected by both large and small tributaries of the Illinois, Mississippi, Missouri, Ohio, and Wabash Rivers. The Ohio River flows along the southernmost boundary of this area in Indiana. Well defined valleys with broad flood plains and numerous stream terraces are along the major streams and rivers. The flood plains along the smaller streams are narrow. Broad summits are nearly level to undulating. Karst topography is common in some parts along the Missouri and Mississippi Rivers and their tributaries. Well-developed karst areas have hundreds of sinkholes, caves, springs, and losing streams. In the St. Louis area, many of the karst features have been obliterated by urban development.

Elevation ranges from 90 feet (20 meters) on the southernmost flood plains to 1,030 feet (320 meters) on the highest ridges. Local relief is mainly 10 to 50 feet (3 to 15 meters) but can be 50 to 150 feet (15 to 45 meters) in the steep, deeply dissected hills bordering rivers and streams. The bluffs along the major rivers are generally 200 to 350 feet (60 to 105 meters) above the valley floor.

The uplands in this MLRA are covered almost entirely with Peoria Loess. The loess can be more than 7 feet (2 meters) thick on stable summits. On the steeper slopes, it is thin or does not occur. In Illinois, the loess is underlain mostly by Illinoian-age till that commonly contains a paleosol. Pre-Illinoian-age till is in parts of this MLRA in Iowa and Missouri and to a minor extent in the western part of Illinois. Wisconsin-age outwash, alluvial deposits, and sandy eolian material are on some of the stream terraces and on dunes along the major tributaries. The loess and glacial deposits are underlain by several bedrock systems. Pennsylvanian and Mississippian bedrock are the most extensive. To a lesser extent are Silurian, Devonian, Cretaceous, and Ordovician bedrock. Karst areas have formed where limestone is near the surface, mostly in the southern part of the MLRA along the Mississippi River and some of its major tributaries. Bedrock outcrops are common on the bluffs along the Mississippi, Ohio, and Wabash Rivers and their major tributaries and at the base of some steep slopes along minor streams and drainageways.

The annual precipitation ranges from 35 to 49 inches (880 to 1,250 millimeters) with a mean of 41 inches (1,050 millimeters). The annual temperature ranges from 48 to 58 degrees F (8.6 to 14.3 degrees C) with a mean of 54 degrees F (12.3 degrees C). The freeze-free period ranges from 150 to 220 days with a mean of 195 days.

**Soils** The dominant soil orders are Alfisols and, to a lesser extent, Entisols and Mollisols. The soils in the area have

a mesic soil temperature regime, an aquic or udic soil moisture regime, and mixed or smectitic mineralogy. They are shallow to very deep, excessively drained to poorly drained, and loamy, silty, or clayey.

The soils on uplands in this area support natural hardwoods. Oak, hickory, and sugar maple are the dominant species. Big bluestem, little bluestem, and scattered oak and eastern redcedar grow on some sites. The soils on flood plains support mixed forest vegetation, mainly American elm, eastern cottonwood, river birch, green ash, silver maple, sweetgum, American sycamore, pin oak, pecan, and willow. Sedge and grass meadows and scattered trees are on some low-lying sites. (United States Department of Agriculture, Natural Resources Conservation Service, 2022)

## **LRU notes**

Most of this LRU (Land Resource Unit) is in the glaciated Till Plains Section of the Central Lowland Province of the Interior Plains. The southeast corner is in the Highland Rim Section (locally known as the Shawnee Hills Section) of the Interior Low Plateaus Province of the Interior Plains. The nearly level to very steep uplands in this LRU are dissected by both large and small tributaries of the Wabash and Ohio Rivers. Well defined valleys with broad flood plains and numerous stream terraces are along the major streams and rivers. The flood plains along the smaller streams are narrow. Broad summits are nearly level to gently sloping.

This area is covered almost entirely with Wisconsin loess. The loess can be more than 7 feet (2 meters) thick on stable summits. On the steeper slopes, it is thin or does not occur. The loess throughout the area is underlain dominantly by glacial till. Wisconsin outwash, alluvial deposits, and sandy eolian material are on some of the stream terraces and on dunes along the major tributaries in the area. The loess and glacial drift are underlain by Pennsylvanian-age bedrock. Bedrock outcrops are common in the walls of the valleys along the Wabash and Ohio Rivers and at the base of some steep slopes along minor streams and drainageways.

The dominant soil orders in this LRU are Alfisols, Entisols, Inceptisols, and Mollisols. The soils in the area have a mesic soil temperature regime, a udic or aquic soil moisture regime, and dominantly mixed or smectitic mineralogy. The soils are very deep, poorly drained to excessively drained, and loamy, silty, or clayey. Nearly level Endoaqualfs (Iva series) and Argiaquolls (Ragsdale series) formed in loess on broad upland summits and flats. Nearly level to steep Hapludalfs (Alford, Iona, Muren, Stoy, and Sylvan series) and Fragiudalfs (Hosmer series) formed in loess on uplands. Hapludalfs (Alvin, Bloomfield, and Princeton series) and Argiudolls (Ade series) formed in sandy eolian material in areas of dunes on uplands and stream terraces. Steep and very steep Hapludalfs (Hickory series) formed in Illinoian till along the major streams and dissected upland drainageways. Hapludalfs (Wellston series) formed in siltstone or sandstone residuum on strongly sloping to steep side slopes underlain by bedrock.

The soils in the major stream valleys include Hapludolls (Carmi series), Argiudolls (Elston series), and Hapludalfs (Skelton series), all of which formed in outwash on nearly level to moderately sloping stream terraces and outwash plains. Endoaquolls (Montgomery series), Endoaquepts (Zipp series), Epiaqualfs (McGary series), and Hapludalfs (Shircliff and Markland series) formed in clayey lacustrine sediments on nearly level to strongly sloping lacustrine terraces or lake plains. Endoaquepts (Evansville series), Endoaquolls (Patton series), and Hapludalfs (Henshaw and Uniontown series) formed in silty sediments on terraces and lake plains.

LRU notes (excerpts from Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. USDA Handbook 296, 2006)

## **Classification relationships**

Major Land Resource Area (MLRA) (USDA-NRCS, 2022):  
115X–Central Mississippi Valley Wooded Slopes

U.S. Forest Service Ecoregions (Cleland et al. 2007):

Domain: Humid Temperate Domain

Division: Hot Continental Division

Province: Eastern Broadleaf Forest (Continental)

Province Code: 222

Section Code: 222G, 222D

These PES sites are similar to other established ecological classifications. Field verification including community surveys are needed to confirm this hypothesis.

Class: 1. Forest & Woodland

Subclass: 1.B. Temperate & Boreal Forest & Woodland

Formation: 1.B.2. Cool Temperate Forest & Woodland

Division: 1.B.2.Na. Eastern North American Forest & Woodland

Macrogroup: M016. Southern & South-Central Oak - Pine Forest & Woodland

Group: G159. South-Central Interior Oak Forest & Woodland

Alliance: A3288. *Quercus alba* - *Quercus falcata* - *Quercus stellata* Ozark-Ouachita Forest Alliance

National Vegetation Classification System (NatureServe): the reference community of this ecological site is similar to: White Oak - Northern Red Oak - Mockernut Hickory / Flowering Dogwood Acidic Forest (CEGL002067)

## Ecological site concept

The historic pre-European settlement vegetation or reference plant community was a mature oak-hickory forest with a closed-canopy and a shade-tolerant herbaceous understory. Sites are on Wellston and Taftown soils which are formed from loess over residuum of shale, sandstone and/or siltstone. The Bedrock Backslope ecological site occurs on upland sites that range from relatively flat to steep slopes (2-35%).

The historic pre-European settlement vegetation or reference plant community was an oak-hickory forest with a shade-tolerant herbaceous understory. These communities are very common in adjacent MLRA 120B but less commonly in MLRA 115X. The reference ecological site has a closed tree canopy (>80%) dominated by northern red oak (*Quercus rubra*) and white oak (*Quercus alba*) along with multiple other species including mockernut hickory (*Carya tomentosa*), shagbark hickory (*Carya ovata*), white ash (*Fraxinus americana*), pignut hickory (*Carya ovalis*), blackgum (*Nyssa sylvatica*), southern red oak (*Quercus falcata*), and black oak (*Quercus velutina*). The subcanopy contained flowering dogwood (*Cornus florida*), blackhaw (*Viburnum prunifolium*) and black cherry (*Prunus serotina*). The shrub layer on these sites are quite variable depending on slope and aspect.

Historically, natural and anthropogenic disturbances such as fire, grazing and wind damage impacted the maintenance and regeneration of these sites. A lack of fire will increase the density of fire-intolerant species such as tulip poplar (*Liriodendron tulipifera*), red maple (*Acer rubrum*), sugar maple (*Acer saccharum*), and basswood (*Tilia americana*).

## Associated sites

F115XA001IL	<b>Silty Upland</b> Silty Upland. Soils are very deep, well drained and formed in loess. Slopes are less than 18%.
F115XA002IL	<b>Silty Backslope</b> Silty Backslope. Soils are very deep, well drained and formed in loess. Slopes are 18% or greater.

## Similar sites

F115XA001IL	<b>Silty Upland</b> Silty Upland. Soils are very deep, well drained and formed in loess.
F115XA002IL	<b>Silty Backslope</b> Silty Backslope. Soils are very deep, well drained and formed in loess. Slopes are 18% or greater.

Table 1. Dominant plant species

Tree	(1) <i>Quercus alba</i> (2) <i>Carya ovata</i>
Shrub	(1) <i>Cornus florida</i>

Herbaceous	(1) <i>Parthenocissus quinquefolia</i> (2) <i>Sanicula canadensis</i>
------------	--

## Physiographic features

These sites are located on various landforms including hill, ridge and structural bench. Elevation of these sites are between 341' to 1007'. Parent material kind is loess over residuum and origin is silty loess over sandstone, siltstone, and/or shale. Wet layer depth is greater than 6 feet. Permeability is impermeable and the runoff class is low to high. These ecological sites do not flood or pond. (NASIS 2020)

**Table 2. Representative physiographic features**


Landforms	(1) Upland > Hill (2) Upland > Hillside or mountainside (3) Upland > Hillslope (4) Upland > Ridge (5) Upland > Structural bench
Runoff class	Low to high
Flooding frequency	None
Ponding frequency	None
Elevation	341–1,007 ft
Slope	2–35%
Water table depth	0–40 in
Aspect	W, NW, N, NE, E, SE, S, SW

## Climatic features


About 60 percent of the precipitation falls during the freeze-free period. Most of the rainfall occurs as high-intensity, convective thunderstorms in summer. Snowfall is common in winter.

**Table 3. Representative climatic features**


Frost-free period (characteristic range)	171-179 days
Freeze-free period (characteristic range)	192-199 days
Precipitation total (characteristic range)	44-47 in
Frost-free period (actual range)	166-180 days
Freeze-free period (actual range)	190-204 days
Precipitation total (actual range)	40-48 in
Frost-free period (average)	175 days
Freeze-free period (average)	196 days
Precipitation total (average)	45 in




**Figure 1. Monthly precipitation range**




**Figure 2. Monthly minimum temperature range**




**Figure 3. Monthly maximum temperature range**



**Figure 4. Monthly average minimum and maximum temperature**



**Figure 5. Annual precipitation pattern**



**Figure 6. Annual average temperature pattern**

## Climate stations used

- (1) EVANSVILLE REGIONAL AP [USW00093817], Evansville, IN
- (2) PRINCETON 1 W [USC00127125], Princeton, IN
- (3) VINCENNES 5 NE [USC00129113], Vincennes, IN
- (4) TERRE HAUTE CAA AP [USW00093823], Terre Haute, IN
- (5) MT VERNON [USC00126001], Uniontown, IN

## Influencing water features

Bedrock Backslope sites are not influenced by wetland or riparian water features. Precipitation is the main source of water for vegetation. Sites are well drained, and permeability is very slow. Runoff ranges from low to high depending on slope. On sloping sites, surface runoff water occurs to downslope ecological sites.

## Soil features

Soil series associated with this site include Wellston and Taftown. Wellston soils are predominately in MRLA 120A and MLRA 120B with limited sites in MRLA 115X. Taftown, which was previously mapped as Wellston, is currently mapped in only one county in Indiana. Both soils are deep, well drained, and are formed in loess over loamy residuum over sandstone and/or shale. These soils have a sodium adsorption ratio of zero and a pH range of 3.5-7.3. Soils of this ecological site are classified as mesic Alfisols (NASIS, 2020)

OSD Taxonomic Classes:

Wellston: Fine-silty, mixed, active, mesic Ultic Hapludalfs

Taftown: Coarse-silty, mixed, active, mesic Typic Hapludalfs

**Table 4. Representative soil features**

Parent material	(1) Loess (2) Residuum—sandstone and shale (3) Residuum—sandstone
-----------------	---

Surface texture	(1) Silt loam (2) Silty clay loam
Drainage class	Well drained
Permeability class	Very slow
Depth to restrictive layer	33–64 in
Soil depth	40–60 in
Surface fragment cover ≤3"	0%
Surface fragment cover >3"	0%
Available water capacity (Depth not specified)	5–8 in
Calcium carbonate equivalent (Depth not specified)	0%
Electrical conductivity (Depth not specified)	0 mmhos/cm
Sodium adsorption ratio (Depth not specified)	0
Soil reaction (1:1 water) (Depth not specified)	3.5–7.3
Subsurface fragment volume ≤3" (Depth not specified)	0–13%
Subsurface fragment volume >3" (Depth not specified)	0–15%

## Ecological dynamics

Bedrock Backslope ecological sites are on well drained uplands with soils formed in loess and loamy residuum from shale and/or sandstone. The historic reference site was a mature oak-hickory forest. Dominant species include white oak (*Quercus alba*), northern red oak (*Quercus rubra*), and shagbark hickory (*Carya ovata*). Sites with a higher slopes and exposed aspect often include more southern red oak (*Quercus falcata*) and black oak (*Quercus velutina*). The sub-canopy includes flowering dogwood (*Cornus florida*). The shrub layer consists of Virginia creeper (*Parthenocissus quinquefolia*) and in some areas, blueberries (*Vaccinium* spp.). The herbaceous layer will include ticktrefoil (*Desmodium* spp.), bedstraws (*Galium* spp.), and Christmas fern (*Polystichum acrostichoides*). The shrub and herbaceous layers will be influenced by slope and aspect, so density and composition of the understory will vary on these sites.


Natural impacts such as drought, wind damage, ice storms and wildfires influenced these communities over centuries. Periodic fire was one of the disturbances that helped to maintain the oak dominance. Often these fires were of low-severity and helped to clear the litter from the forest floor and encourage oak regeneration. Lightning was the usual ignition source; although native Americans did use fire to reduce understory brush or increase forage. Long-term periods of drought did occur and would have influenced fire severity, species composition, shrub density, plant mortality, and tree growth rates.

Most lower slope sites have now been converted to agriculture – either cropland or hayland production. Cleared sites on higher slopes are often grazed; however, management is limited by slope. Landowner should be aware of potential overgrazing impacts such as soil erosion, soil compaction, water quality impacts, and noxious weeds. Invasive non-native vegetation is a serious concern in many remaining wooded areas as bush honeysuckle, euonymus, Japanese honeysuckle, privet, and other non-native plants have been introduced and are increasing without management controls.

Long-term fire suppression has resulted in some sites with an increased in fire in-tolerant species such as sugar maple, red maple, white ash, and tulip poplar. Saplings of sugar maple, red maple, hickory, tulip poplar, and white ash are often noticeable in the understory on these sites. Oak regeneration is usually low. Without management inputs such as prescribed fire or timber stand improvement work, oak regeneration may be very limited.


# State and transition model

## Ecosystem states




- T1A** - Large scale disturbance
- T1C** - Clearing of site; agricultural production - forage
- T1B** - Clearing of site; agricultural production -row crops.
- R2A** - Restoration inputs such as planting, brush control, prescribed fire, and timber stand improvement.
- T2B** - Clearing; agricultural production - forage
- T2C** - Clearing; agricultural production - row crops
- T3B** - Abandonment of agricultural practices
- T3A** - Site preparation and tillage, seeding, weed control, cropland management
- T4A** - Transition site to forage production; seeding; weed/brush control; pasture management

## State 1 submodel, plant communities




- 1.1.A** - Fire suppression
- 1.2.A** - Natural fire regime

## State 2 submodel, plant communities



## State 3 submodel, plant communities





## State 1

### Reference State

Two communities are associated with this reference site and the community phases are based on natural fire regime or lack thereof. The historic reference site was a mature oak-hickory forest. Dominant species include white oak (*Quercus alba*), northern red oak (*Quercus rubra*), and shagbark hickory (*Carya ovata*). Sites with a higher slopes and exposed aspect often include more southern red oak (*Quercus falcata*) and black oak (*Quercus velutina*). The sub-canopy includes flowering dogwood (*Cornus florida*). The shrub layer consists of Virginia creeper (*Parthenocissus quinquefolia*) and in some areas, blueberries (*Vaccinium* spp.). The herbaceous layer will include ticktrefoil (*Desmodium* spp.), bedstraws (*Galium* spp.), and Christmas fern (*Polystichum acrostichoides*). The shrub and herbaceous layers will be influenced by slope and aspect, so density and composition of the understory will vary on these sites. Historically these sites were influenced by reoccurring wildfires, wind damage, ice storms, and grazing by native species.

#### Dominant plant species

- white oak (*Quercus alba*), tree
- shagbark hickory (*Carya ovata*), tree
- flowering dogwood (*Cornus florida*), shrub
- Virginia creeper (*Parthenocissus quinquefolia*), other herbaceous
- clustered blacksnakeroot (*Sanicula odorata*), other herbaceous
- Christmas fern (*Polystichum acrostichoides*), other herbaceous

## Community 1.1

### Reference community

The overstory in this phase is dominated by oak and hickory species including white oak, northern red oak, shagbark hickory, mockernut hickory, southern red oak, and black oak. The driest sites may include southern red oak, pignut hickory, scarlet oak and/or post oak. According to LANDFIRE, historical fire frequency was approximately every 1.5 -3 decades. Reoccurring fires would have maintained an oak-dominated forest along with a more open understory.

#### Dominant plant species

- white oak (*Quercus alba*), tree
- shagbark hickory (*Carya ovata*), tree
- northern red oak (*Quercus rubra*), tree
- flowering dogwood (*Cornus florida*), shrub
- black cherry (*Prunus serotina*), shrub
- sassafras (*Sassafras albidum*), shrub
- sanicle (*Sanicula*), other herbaceous
- ticktrefoil (*Desmodium*), other herbaceous
- Virginia creeper (*Parthenocissus quinquefolia*), other herbaceous

## Community 1.2

### Fire suppressed community

This community would be characterized by an increase in fire-intolerant species such as maple and hickory. The understory would especially have a noticeable increase in density of maple seedlings and saplings. The shrub layer would be denser than community 1.1.

### **Dominant plant species**

- sugar maple (*Acer saccharum*), tree
- red maple (*Acer rubrum*), tree
- ash (*Fraxinus*), tree
- tuliptree (*Liriodendron tulipifera*), tree
- sugar maple (*Acer saccharum*), shrub

### **Pathway 1.1.A**

#### **Community 1.1 to 1.2**

Suppression of the natural fire regime will result in the site slowly transitioning to a more mesic mixed deciduous community with an increase in maple, ash, and/or poplar species. Seed sources and the severity/length of disturbance will influence the species composition of this community.

### **Pathway 1.2.A**

#### **Community 1.2 to 1.1**

These communities developed under a historic, natural fire regime which encouraged oak reproduction and reduced competition from more fire-intolerant species such as sugar maple and white ash.

## **State 2**

### **Disturbed State**

Most remaining Bedrock Backslope Forest sites have been altered due to a long-term absence of fire, clearing, grazing, selective harvest and/or invasive species. Trees on site, will depending on the type, length and severity of disturbances. However, sites that have had a long-term absence of fire will display the following characteristics: an increase in fire -intolerant species, an increase in shrub density, an increase in leaf-litter buildup, and an increase in shade-tolerant understory species. Diversity of species may also be reduced, especially if there has been an introduction of non-native species such as bush honeysuckle (*Lonicera maackii*). Many of these sites are eventually transitioned to Pastureland (State 3) or cropland (State 4) is common, especially on lower slope sites.

### **Dominant plant species**

- sugar maple (*Acer saccharum*), tree
- white ash (*Fraxinus americana*), tree
- red maple (*Acer rubrum*), tree
- tuliptree (*Liriodendron tulipifera*), tree
- sugar maple (*Acer saccharum*), shrub

### **Community 2.1**

#### **Disturbed community**

This is a disturbed, successional community that includes a variety of fast-growing trees such as maples and ashes. Other species may include maple, ash, sweetgum and poplar. Shrub and understory species will depend on the type, severity and length of disturbances, available seed sources, and management inputs, if present.

### **Dominant plant species**

- red maple (*Acer rubrum*), tree
- silver maple (*Acer saccharinum*), tree
- white ash (*Fraxinus americana*), tree

## **State 3**

### **Pastureland/ Forage State**

A portion of these sites have been converted to pastureland or forage production. Species selection will depend upon the objectives and goals of the landowner; however, commonly planted grasses include tall fescue

(*Schedonorus arundinaceus*), brome (*Bromus* spp.), white clover (*Trifolium repens*) and red clover (*Trifolium pratense*). Species health and productivity are determined by the management and long-term overgrazing on some sites has caused soil erosion and compaction.

#### **Dominant plant species**

- tall fescue (*Schedonorus arundinaceus*), grass
- brome (*Bromus*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- white clover (*Trifolium repens*), other herbaceous
- red clover (*Trifolium pratense*), other herbaceous

### **Community 3.1**

#### **Pastureland/Forage community**

These sites are managed for forage production and often include tall fescue (*Schedonorus arundinaceus*), brome (*Bromus* spp.), white clover (*Trifolium repens*) and red clover (*Trifolium pratense*). Selection of species will depend on the landowner's objectives.

#### **Dominant plant species**

- tall fescue (*Schedonorus arundinaceus*), grass
- brome (*Bromus*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- red clover (*Trifolium pratense*), other herbaceous
- white clover (*Trifolium repens*), other herbaceous

### **State 4**

#### **Cropland State**

This is a state is only appropriate for sites on lower slopes. Soil erosion potential and water quality issues on higher slope sites makes them more appropriate for forestland or well managed grazing. Common crops include corn (*Zea mays*), soybeans (*Glycine max*), and occasionally winter wheat (*Triticum aestivum*). Some landowners choose to convert sites to cool season grasses for a period before resuming cropland production. A return to the historical Reference State from State 4 is unlikely, if not impossible.

#### **Dominant plant species**

- corn (*Zea mays*), other herbaceous
- soybean (*Glycine*), other herbaceous

### **Community 4.1**

#### **Cropland community**

This community is characterized by the management and production of row crop agriculture. Common species include corn, soybean and wheat. Many other crops are suitable for these sites, and species selection will depend upon the landowners goals and objectives.

#### **Dominant plant species**

- soybean (*Glycine max*), other herbaceous
- corn (*Zea mays*), other herbaceous

### **Transition T1A**

#### **State 1 to 2**

Severe disturbances, such as clearing or selective harvesting (oak/hickory removal) , will transition this site to State 2.

## **Transition T1C**

### **State 1 to 3**

Site is transitioned to an agricultural site focused on forage production. Management inputs would include clearing, site preparation, seeding and weed/brush control.

## **Transition T1B**

### **State 1 to 4**

Site is transitioned to an agricultural site focused on row crop production. Management inputs would include clearing, site preparation, seeding and weed control. Hydrological modifications are often installed to aid in drainage.

## **Restoration pathway R2A**

### **State 2 to 1**

Restoration would require long-term management inputs including planting of desired species, weed control, brush control, timber stand improvement, and prescribed fire.

## **Transition T2B**

### **State 2 to 3**

Site is cleared and forage/pasture production is initiated. Management inputs would include tree/shrub removal, site preparation, seeding, and weed/brush control.

## **Transition T2C**

### **State 2 to 4**

Site is cleared and row crop production is initiated. Management inputs would include tree/shrub removal, site preparation, tillage, seeding, and weed control.

## **Transition T3B**

### **State 3 to 2**

Site is abandoned and slowly would transition to a wooded state dominated by deciduous trees. Species on site would depend on the severity and length of disturbance and available seed sources.

## **Transition T3A**

### **State 3 to 4**

Management inputs that transition a site from pasture or forage production to a site that is utilized for row crop production.

## **Transition T4A**

### **State 4 to 3**

Management inputs to transition a site from cropland production to a state of pasture/forage production.

## **Additional community tables**

### **Inventory data references**

A Provisional Ecological Site Description (PESD) describes ecological potential and ecosystem dynamics of land areas and their potential management. Ecological sites are linked to soil survey map unit components, which allows for mapping of ecological sites. A PESD with a provisional status represents the lowest tier of documentation that is releasable to the public. No field level data have been collected as part of this PESD. It is expected that a PESD will continue to be refined through field verification and field sampling.

Reference and alternative state concepts, including the state-and-transition model and vegetative communities are not yet well-documented and will require field sampling for verification.

This document is provisional.

## Other references

Brinson, M. M. 1993. A hydrogeomorphic classification for wetlands. Technical Report WRP-DE-4, U.S. Army Corps of Engineers, Engineer Waterways Experiment Station, Vicksburg, MS.

Braun, E. Lucy. 2001. Deciduous forests of eastern North America. Caldwell, N.J.: Blackburn Press.

Cleland, D. T., J. A. Freeouf, J. E. Keys, G. J. Nowacki, C. Carpenter, and W. H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Conterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pp.

Comer P. J., Faber-Langendoen D, Evans R, Gawler S. C, Josse C, Kittel G, Menard S, Pyne M, Reid M, Schulz K, Snow K, and Teague J. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia.

Cowardin, L.M., V. Carter, F. C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep water habitats of the United States. U.S. Dept. of Interior, Fish & Wildlife Service, Office of Biological Services, Washington DC. FWS/OBS-79/31. 142 pp.

Homoya, M. A., Abrell, D. B., Aldrich, J. R., & Post, T. W. (1985). The Natural Regions of Indiana. Indiana Academy of Science, 94, 245-269

Jackson, Marion T. 1997. The Natural heritage of Indiana. Bloomington: Indiana University Press, published in association with the Indiana Department of Natural Resources and the Indiana Academy of Science.

LANDFIRE (Landfire National Vegetation Dynamics Database). 2009. Landfire National Vegetation Dynamics Models. Landfire Project, USDA Forest Service, U.S. Department of Interior. (<http://www.LANDFIRE.gov/index.php>: accessed 2020).

Mohlenbrock, R. H. and D. M. Ladd. 1978. Distribution of Illinois Vascular Plants. Southern Illinois Univ. Press, Carbondale and Edwardsville, Ill. 282 pp.

Mohlenbrock, R. H. 2003. Vascular Flora of Illinois, 3rd edition. Carbondale, Illinois: Southern Illinois University Press. 736 pp.

National Cooperative Soil Survey (NCSS). National Cooperative Soil Characterization Database. Available online: <https://ncsslabdatamart.sc.egov.usda.gov/>. Accessed: 2020.

NatureServe. 2018. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available <http://explorer.natureserve.org>. (Association Detail Report: CEG002427) (Accessed: 2020)

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Web Soil Survey (SSS NRCS WSS). Available online at the following link: <https://websoilsurvey.sc.egov.usda.gov/>. Accessed 2020.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Official Soil Series Descriptions (SSS NRCS OSD). Available online. Accessed 2020. <https://soilseries.sc.egov.usda.gov/osdname.aspx>

United States Department of Agriculture, Natural Resources Conservation Service (USDA – NRCS). 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. USDA

Handbook 296. 682 pp.

United States Department of Agriculture, Natural Resources Conservation Service. 2022. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture, Agriculture Handbook 296.

USDA, NRCS. 2018. The PLANTS Database (<http://plants.usda.gov>, 1 March 2018). National Plant Data Team, Greensboro, NC 27401-4901 USA.

Voigt, J. W., and R. H. Mohlenbrock. 1964. Plant communities of southern Illinois. Southern Illinois University Press, Carbondale. 202 pp.

Whitaker, John O., Charles J. Amlaner, Marion T. Jackson, George R. Parker, and Peter Evans Scott. 2012. Habitats and ecological communities of Indiana presettlement to present. Bloomington: Indiana University Press.

## Contributors

Ralph Tucker, Soil Survey Office Leader, USDA-NRCS, Missouri  
Dena Anderson, Resource Soil Scientist, USDA-NRCS, Indiana  
Zack Weber, Soil Scientist, USDA-NRCS, Illinois  
Anita Arends, Ecological Site Specialist, USDA-NRCS, Illinois

## Approval

Suzanne Mayne-Kinney, 12/30/2024

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	01/08/2025
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
- 
5. **Number of gullies and erosion associated with gullies:**
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:**
- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

- 
16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-