

## **Ecological site F115XC008IL Loess Exposed Backslope Woodland**

Last updated: 12/30/2024  
Accessed: 01/09/2025

---

### **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### **MLRA notes**

Major Land Resource Area (MLRA): 115X—Central Mississippi Valley Wooded Slopes

This MLRA is characterized by deeply dissected, loess-covered hills bordering well defined valleys of the Illinois, Mississippi, Missouri, Ohio, and Wabash Rivers and their tributaries. It is used to produce cash crops and livestock. About one-third of the area is forested, mostly on the steeper slopes. This area is in Illinois (50 percent), Missouri (36 percent), Indiana (13 percent), and Iowa (1 percent) in two separate areas. It makes up about 25,084 square miles (64,967 square kilometers).

Most of this area is in the Till Plains section and the Dissected Till Plains section of the Central Lowland province of the Interior Plains. The Springfield-Salem plateaus section of the Ozarks Plateaus province of the Interior Highlands occurs along the Missouri River and the Mississippi River south of the confluence with the Missouri River. The nearly level to very steep uplands are dissected by both large and small tributaries of the Illinois, Mississippi, Missouri, Ohio, and Wabash Rivers. The Ohio River flows along the southernmost boundary of this area in Indiana. Well defined valleys with broad flood plains and numerous stream terraces are along the major streams and rivers. The flood plains along the smaller streams are narrow. Broad summits are nearly level to undulating. Karst topography is common in some parts along the Missouri and Mississippi Rivers and their tributaries. Well-developed karst areas have hundreds of sinkholes, caves, springs, and losing streams. In the St. Louis area, many of the karst features have been obliterated by urban development.

Elevation ranges from 90 feet (20 meters) on the southernmost flood plains to 1,030 feet (320 meters) on the highest ridges. Local relief is mainly 10 to 50 feet (3 to 15 meters) but can be 50 to 150 feet (15 to 45 meters) in the steep, deeply dissected hills bordering rivers and streams. The bluffs along the major rivers are generally 200 to 350 feet (60 to 105 meters) above the valley floor.

The uplands in this MLRA are covered almost entirely with Peoria Loess. The loess can be more than 7 feet (2 meters) thick on stable summits. On the steeper slopes, it is thin or does not occur. In Illinois, the loess is underlain mostly by Illinoian-age till that commonly contains a paleosol. Pre-Illinoian-age till is in parts of this MLRA in Iowa and Missouri and to a minor extent in the western part of Illinois. Wisconsin-age outwash, alluvial deposits, and sandy eolian material are on some of the stream terraces and on dunes along the major tributaries. The loess and glacial deposits are underlain by several bedrock systems. Pennsylvanian and Mississippian bedrock are the most extensive. To a lesser extent are Silurian, Devonian, Cretaceous, and Ordovician bedrock. Karst areas have formed where limestone is near the surface, mostly in the southern part of the MLRA along the Mississippi River and some of its major tributaries. Bedrock outcrops are common on the bluffs along the Mississippi, Ohio, and Wabash Rivers and their major tributaries and at the base of some steep slopes along minor streams and drainageways.

The annual precipitation ranges from 35 to 49 inches (880 to 1,250 millimeters) with a mean of 41 inches (1,050 millimeters). The annual temperature ranges from 48 to 58 degrees F (8.6 to 14.3 degrees C) with a mean of 54 degrees F (12.3 degrees C). The freeze-free period ranges from 150 to 220 days with a mean of 195 days.

**Soils** The dominant soil orders are Alfisols and, to a lesser extent, Entisols and Mollisols. The soils in the area have

a mesic soil temperature regime, an aquic or udic soil moisture regime, and mixed or smectitic mineralogy. They are shallow to very deep, excessively drained to poorly drained, and loamy, silty, or clayey.

The soils on uplands in this area support natural hardwoods. Oak, hickory, and sugar maple are the dominant species. Big bluestem, little bluestem, and scattered oak and eastern redcedar grow on some sites. The soils on flood plains support mixed forest vegetation, mainly American elm, eastern cottonwood, river birch, green ash, silver maple, sweetgum, American sycamore, pin oak, pecan, and willow. Sedge and grass meadows and scattered trees are on some low-lying sites. (United States Department of Agriculture, Natural Resources Conservation Service, 2022)

## **LRU notes**

The Central Mississippi Valley Wooded Slopes, Northern part (Land Resource Unit (LRU) (115XC) encompasses the Wyaconda River Dissected Till Plains, Mississippi River Hills, and Mississippi River Alluvial Plain (Schwegman et al. 1973; Nelson 2010). It spans three states – Illinois (73 percent), Iowa (6 percent), and Missouri (21 percent) – comprising about 13,650 square miles (Figure 1). The elevation ranges from 420 feet above sea level (ASL) along the Mississippi River floodplains to 885 feet on the upland ridges. Local relief varies from 10 to 20 feet but can be as high as 50 to 100 feet along drainageways and streams and the bluffs on the major rivers reaching 250 feet above valley floors. Wisconsin-aged loess covers the uplands, while Illinoian glacial drift lies directly below. The loess and drift deposits are underlain by several bedrock systems, including the Cretaceous, Pennsylvania, Mississippian, Silurian, Devonian, and Ordovician Systems. Wisconsin outwash deposits and sandy eolian material occur along stream terraces of major tributaries (USDA-NRCS 2006).

The vegetation across the region has undergone drastic changes over time. At the end of the last glacial episode – the Wisconsin glacial episode – the evolution of vegetation began with the development of tundra habitats, followed by a phase of spruce and fir forests, and eventually spruce-pine forests. Not until approximately 9,000 years ago did the climate undergo a warming trend which prompted the development of deciduous forests dominated by oak and hickory. As the climate continued to warm and dry, prairies began to develop approximately 8,300 years ago. Another shift in climate that resulted in an increase in moisture prompted the emergence of savanna-like habitats from 8,000 to 5,000 years before present (Taft et al. 2009). During the most recent climatic shifts, forested ecosystems maintained footholds on steep valley sides and wet floodplains. Due to the physiography of the MLRA, forests were the dominant ecosystems and were affected by such natural disturbances as droughts, wind, lightning, and occasional fire (Taft et al. 2009).

## **Classification relationships**

Major Land Resource Area (MLRA) (USDA-NRCS, 2022):  
115X–Central Mississippi Valley Wooded Slopes

USFS Subregions: Central Dissected Till Plains (251C)Section; Western Mississippi River Hills (251Ce), Mississippi River and Illinois Alluvial Plains (251Cf), Eastern Mississippi River Hills (251Ci), Galesburg Dissected Till Plain (251Cj), and Wyaconda River Dissected Till Plain (251Cm) Subsections (Cleland et al. 2007)

U.S. EPA Level IV Ecoregion: Upper Mississippi River Alluvial Plain (72d), River Hills (72f), and Western Dissected Illinoian Till Plain (72i) (USEPA 2013)

National Vegetation Classification – Ecological Systems: North-Central Interior Dry-Mesic Oak Forest and Woodland (CES202.046) (NatureServe 2018)

National Vegetation Classification – Plant Associations: *Quercus alba* – *Quercus rubra* – *Carya ovata* Glaciated Forest (CEGL002068) (Nature Serve 2018)

Biophysical Settings: North-Central Interior Dry-Mesic Oak Forest and Woodland (BpS 4913100)

Illinois Natural Areas Inventory: Dry-mesic upland forest (White and Madany 1978)

Missouri Terrestrial Natural Communities: Dry-mesic loess/glacial till woodland (Nelson 2010)

Ecological site concept

Loess Exposed Backslope Woodlands are located within the green areas on the map. They occur on south and west-facing upland backslopes. The soils are Alfisols and Entisols that are somewhat poorly to well drained and very deep, formed in loess and loess-covered substrates.

The historic pre-European settlement vegetation on this ecological site was dominated by an open oak-hickory woodland with a continuous understory. White oak (*Quercus alba* L.) and shagbark hickory (*Carya ovata* (Mill.) K. Koch) are the dominant species in the tree canopy. Black oak (*Quercus velutina* Lam.) and mockernut hickory (*Carya tomentosa* (Lam.) Nutt.) can be common canopy associates. Hairy woodland brome (*Bromus pubescens* Muhl. ex Willd.) and elmleaf goldenrod (*Solidago ulmifolia* Muhl. ex Willd.) are the dominant herbaceous species. Forbs typical of an undisturbed plant community associated with this ecological site include Culver's root (*Veronicastrum virginicum* (L.) Farw.) and smooth blue aster (*Symphotrichum laeve* (L.) Á. Löve & D. Löve var. *laeve*) (Ladd and Thomas 2015). Fire is the primary disturbance factor that maintains this ecological site, while drought, windthrow, and grazing are secondary factors (LANDFIRE 2009).

Associated sites

F115XC005IL	<b>Loess Upland Forest</b> Loess and loess-covered substrates on uplands including Atlas, Baylis, Bunkum, Caseyville, Creal, Derinda, Dodge, Fayette, Fishhook, Hickory, Kendall, Keomah, Keswick, Menfro, Metea, Navlys, Rozetta, Seaton, Stookey, Stronghurst, Sylvan, Thebes, Timula, Ursa, and Winfield soils. Slope is less than 18 percent and has a water table greater than 12 inches.
F115XC007IL	<b>Loess Protected Backslope Forest</b> Loess and loess-covered substrates on south and west-facing backslopes including Atlas, Baylis, Fayette, Hennepin, Hickory, Keswick, Menfro, Seaton, Stookey, Sylvan, Timula, and Ursa soils. This site is greater than 18 percent slope and on north and east aspects.

Similar sites


F115XC007IL	<b>Loess Protected Backslope Forest</b> This site is greater than 18 percent slope and on north and east aspects.
-------------	--

Table 1. Dominant plant species


Tree	(1) <i>Quercus alba</i> (2) <i>Carya ovata</i>
Shrub	Not specified
Herbaceous	(1) <i>Bromus pubescens</i> (2) <i>Solidago ulmifolia</i>

Physiographic features

Loess Exposed Backslope Woodlands occur on south and west facing upland backslopes. They are situated on elevations ranging from approximately 341 to 1738 feet above sea level (ASL). The site does not experience flooding but rather generates runoff to adjacent, downslope ecological sites.



**Figure 1. Location of Loess Exposed Backslope Woodland ecological site within LRU 115XC.**



**Figure 2. Representative block diagram of Loess Exposed Backslope Woodland and associated ecological sites.**

**Table 2. Representative physiographic features**

Hillslope profile	(1) Backslope
Slope shape across	(1) Convex
Slope shape up-down	(1) Convex
Landforms	(1) Upland
Runoff class	High to very high
Elevation	104–530 m
Slope	15–70%
Water table depth	46–203 cm
Aspect	W, SE, S, SW

### Climatic features

The Central Mississippi Valley Wooded Slopes, Northern Part falls into the humid subtropical (Cfa) and hot-summer humid continental climate (Dfa) Köppen-Geiger climate classifications (Peel et al. 2007). The two main factors that drive the climate of the MLRA are latitude and weather systems. Latitude, and the subsequent reflection of solar input, determines air temperatures and seasonal variations. Solar energy varies across the seasons, with summer receiving three to four times as much energy as opposed to winter. Weather systems (air masses and cyclonic storms) are responsible for daily fluctuations of weather conditions. High-pressure systems are responsible for settled weather patterns where sun and clear skies dominate. In fall, winter, and spring, the polar jet stream is responsible for the creation and movement of low-pressure systems. The clouds, winds, and precipitation

associated with a low-pressure system regularly follow high-pressure systems every few days (Angel n.d.).

The soil temperature regime of LRU 115XC is classified as mesic, where the mean annual soil temperature is between 46 and 59°F (USDA-NRCS 2006). Temperature and precipitation occur along a north-south gradient, where temperature and precipitation increase the further south one travels. The average freeze-free period of this ecological site is about 182 days, while the frost-free period is about 146 days. The majority of the precipitation occurs as rainfall in the form of convective thunderstorms during the growing season. Average annual precipitation is 39 inches, which includes rainfall plus the water equivalent from snowfall. The average annual low and high temperatures are 42 and 63°F, respectively.

Table 3. Representative climatic features

Frost-free period (characteristic range)	139-150 days
Freeze-free period (characteristic range)	169-199 days
Precipitation total (characteristic range)	965-1,041 mm
Frost-free period (actual range)	135-162 days
Freeze-free period (actual range)	167-204 days
Precipitation total (actual range)	914-1,041 mm
Frost-free period (average)	146 days
Freeze-free period (average)	182 days
Precipitation total (average)	991 mm




Figure 3. Monthly precipitation range







Figure 4. Monthly minimum temperature range




**Figure 5. Monthly maximum temperature range**



**Figure 6. Monthly average minimum and maximum temperature**



**Figure 7. Annual precipitation pattern**



**Figure 8. Annual average temperature pattern**

## Climate stations used

- (1) MORRISON [USC00115833], Morrison, IL
- (2) NEW BOSTON DAM 17 [USC00116080], Wapello, IL
- (3) LA HARPE [USC00114823], La Harpe, IL

- (4) HAVANA [USC00113940], Lewistown, IL
- (5) PITTSFIELD #2 [USC00116837], Pittsfield, IL
- (6) ST CHARLES CO AP [USW00053904], Portage des Sioux, MO

## Influencing water features

Loess Exposed Backslope Woodland are not influenced by wetland or riparian water features. Precipitation is the main source of water for this ecological site. Infiltration is very slow to moderate (Hydrologic Groups B, C, and D), and surface runoff is high to very high. Surface runoff contributes some water to downslope ecological sites.




Figure 9. Hydrologic cycling in Loess Exposed Backslope Woodland ecological site.

## Soil features

Soils of Loess Exposed Backslope Woodlands are classified as Aerlic Chromic Vertic Epiaqualfs, Aquertic Chromic Hapludalfs, Chromic Vertic Hapludalfs, Typic Hapludalfs, Typic Paleudalfs, Typic Eutrudepts with very slow to moderate infiltration and high to very high runoff potential. The soil series associated with this site includes Atlas, Baylis, Fayette, Hennepin, Hickory, Keswick, Menfro, Seaton, Stookey, Sylvan, Timula, and Ursa. The parent material is loess and loess-covered substrates, and the soils are somewhat poorly to well-drained and very deep. Soil pH classes are very strongly acid to moderately alkaline. A shallow, densic material may be noted as a rooting restriction are for some soils of this ecological site.




Figure 10. Profile sketches of soil series associated with Loess Exposed Backslope Woodland.

Table 4. Representative soil features

Parent material	(1) Loess
-----------------	-----------

Surface texture	(1) Silty clay loam (2) Silty clay (3) Silt loam
Family particle size	(1) Fine-silty
Drainage class	Somewhat poorly drained to well drained
Permeability class	Very slow to moderately slow
Depth to restrictive layer	203 cm
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (Depth not specified)	7.62–20.32 cm
Calcium carbonate equivalent (Depth not specified)	0–40%
Electrical conductivity (Depth not specified)	0–2 mmhos/cm
Soil reaction (1:1 water) (Depth not specified)	4.5–8.4
Subsurface fragment volume <=3" (Depth not specified)	3–18%
Subsurface fragment volume >3" (Depth not specified)	0–16%

## Ecological dynamics

The information in this Ecological Site Description, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

The MLRA lies within the tallgrass prairie ecosystem of the Midwest, but a variety of environmental and edaphic factors resulted in a landscape that historically supported upland hardwood forests, lowland mixed forests, and scattered grass and sedge meadows. Loess Exposed Backslope Woodlands form an aspect of this vegetative continuum. This ecological site occurs on south and west facing upland backslopes on somewhat poorly to well drained soils. Species characteristic of this ecological site consist of an open canopy of oak and hickory with herbaceous vegetation.

Fire is a critical factor that maintains Loess Exposed Backslope Woodlands. Fire typically consisted of low- to moderate-severity surface fires every 15 to 25 years (LANDFIRE 2009). Ignition sources included summertime lightning strikes from convective storms and bimodal, human ignitions during the spring and fall seasons. Native Americans regularly set fires to improve sight lines for hunting, drive large game, improve grazing and browsing habitat, agricultural clearing, and enhance vital ethnobotanical plants (Barrett 1980; LANDFIRE 2009).

Drought and storm damage have also played a role in shaping this ecological site. The periodic episodes of reduced soil moisture in conjunction with the somewhat poorly to well-drained soils have favored the proliferation of plant species tolerant of such conditions. Drought can also slow the growth of plants and result in dieback of certain species. Damage to trees from wind and ice storms can vary from minor, patchy effects of individual trees to stand effects that temporarily affect community structure and species richness and diversity (Irland 2000; Peterson 2000). When coupled with fire, periods of drought and catastrophic storm damage can greatly delay the establishment and maturation of woody vegetation (Pyne et al. 1996).


Today, Loess Exposed Backslope Woodlands have been reduced from their pre-settlement extent. Moderate slopes have been converted to cropland, while steeper slopes have been converted to forage land. Remnants that do exist




have experienced long-term fire suppression and overbrowsing resulting in significant changes to the forest structure. A return to the historic plant community may not be possible following extensive land modification, but long-term conservation agriculture or woodland reconstruction efforts can help to restore some biotic diversity and ecological function. The state-and-transition model that follows provides a detailed description of each state, community phase, pathway, and transition. This model is based on available experimental research, field observations, literature reviews, professional consensus, and interpretations.

## State and transition model

### Ecosystem states



### States 2 and 5 (additional transitions)



**T1A** - Long-term fire suppression and/or land abandonment

**T1B** - Cultural treatments are implemented to increase forage quality and yield

**T1C** - Agricultural conversion via tillage, seeding, and non-selective herbicide

**T2A** - Cultural treatments are implemented to increase forage quality and yield

**T2B** - Agricultural conversion via tillage, seeding, and non-selective herbicide

**T3A** - Long-term fire suppression and/or land abandonment

**T3B** - Agricultural conversion via tillage, seeding, and non-selective herbicide

**R3A** - Site preparation, tree planting, non-native species control, and native seeding

**T4A** - Long-term fire suppression and/or land abandonment

**T4B** - Cultural treatments are implemented to increase forage quality and yield


**R4A** - Site preparation, tree planting, non-native species control, and native seeding

**T5A** - Long-term fire suppression and/or land abandonment

**T5B** - Cultural treatments are implemented to increase forage quality and yield

**T5C** - Agricultural conversion via tillage, seeding, and non-selective herbicide


### State 1 submodel, plant communities



**1.1A** - Fire return interval greater than 25 years

**1.2A** - Replacement fire every 20 years


#### State 2 submodel, plant communities



**2.1A** - Continued fire suppression

**2.2A** - Severe disturbance event

#### State 3 submodel, plant communities



**3.1A** - Mechanical harvesting is replaced with domestic livestock grazing

**3.1B** - Mechanical harvesting is replaced with domestic livestock. Forage plants are overutilized.


**3.2A** - Domestic livestock grazing is replaced by mechanical harvesting

**3.2B** - Grazing; overutilization of forage plants

**3.3B** - Mechanical harvesting

**P3.3A** - Implementation of grazing; forage to animal balance


#### State 4 submodel, plant communities



**4.1A** - Less tillage, residue management

- 4.1B - Less tillage, residue management, and implementation of cover cropping
- 4.2A - Intensive tillage, remove residue, and reinitiate monoculture row cropping
- 4.2B - Implementation of cover cropping
- 4.3B - Intensive tillage, remove residue, and reinitiate monoculture row cropping
- 4.3A - Remove cover cropping

#### State 5 submodel, plant communities



- 5.1A - Invasive species control and implementation of disturbance regimes
- 5.2A - Drought or improper timing/use of management actions

## State 1 Reference State

The reference plant community is categorized as an oak-hickory thicket, dominated by deciduous trees and shade-tolerant herbaceous vegetation. The one community phase within the reference state is dependent on recurring fire intervals. The severity and intensity of fire alters species composition, cover, and extent, while regular fire intervals keep the canopy from succeeding to mesophytic, fire-intolerant species. Drought and catastrophic storm damage have more localized impacts in the reference phases, but do contribute to overall species composition, diversity, cover, and productivity

### Dominant plant species

- white oak (*Quercus alba*), tree
- shagbark hickory (*Carya ovata*), tree
- hairy woodland brome (*Bromus pubescens*), grass
- elmleaf goldenrod (*Solidago ulmifolia*), other herbaceous

## Community 1.1

### White Oak - Shagbark Hickory / Hairy woodland Brome - Elmleaf Goldenrod

Sites in this reference community phase are an open canopy woodland. White oak and shagbark hickory are the dominant tree species, but black oak and mockernut hickory are common canopy associates. Trees are large (21 to 33 inches DBH) and cover ranges from 21 to 60 percent (LANDFIRE 2009). The herbaceous layer is mostly a continuous mix of graminoids and forbs including hairy woodland brome, eastern bottlebrush grass (*Elymus hystrix* L. var. *hystrix*), elmleaf goldenrod, and hairy sunflower (*Helianthus hirsutus* Raf.) (Nelson 2010). Surface fires occurring approximately every 20 years will maintain this phase, but beyond 25 years the community will shift to phase 1.2 (LANDFIRE 2009).

### Dominant plant species

- white oak (*Quercus alba*), tree
- shagbark hickory (*Carya ovata*), tree
- hairy woodland brome (*Bromus pubescens*), grass
- elmleaf goldenrod (*Solidago ulmifolia*), other herbaceous

## Community 1.2

### White Oak – Shagbark Hickory/Fragrant Sumac/Hairy Woodland Brome – Elmleaf Goldenrod

This community phase represents natural succession as a result of an extended fire return interval. The lack of disturbance allows the overstory to shift. Tree size class remains steady, but canopy coverage ranges from 61 to 80

percent shifting the site to a closed canopy woodland (LANDFIRE 2009). The shrub layer becomes more prominent during this phase with species such as fragrant sumac (*Rhus aromatica* Aiton) (Nelson 2010). Heliophytic herbs remain common in the understory but shade-tolerant species begin to develop as well. Periodic surface fires will maintain this phase, but surface fires occurring approximately every 20 years will shift the community back to phase 1.1 (LANDFIRE 2009).

#### **Dominant plant species**

- white oak (*Quercus alba*), tree
- shagbark hickory (*Carya ovata*), tree
- fragrant sumac (*Rhus aromatica*), shrub
- hairy woodland brome (*Bromus pubescens*), grass
- elmleaf goldenrod (*Solidago ulmifolia*), other herbaceous

#### **Pathway 1.1A**

##### **Community 1.1 to 1.2**

Fire return interval greater than 25 years.

#### **Pathway 1.2A**

##### **Community 1.2 to 1.1**

Low to moderate intensity surface fires occur every 15 to 25 years.

### **State 2**

#### **Fire Suppressed State**

Fire suppression can transition the reference plant community from an oak forest to an oak-maple mesophytic forest. As the natural fire regime is removed from the landscape, encroachment and dominance by shade-tolerant, fire-intolerant species ensues. This results in a positive feedback loop of mesophication whereby plant community succession continuously creates cool, damp shaded conditions that perpetuate a closed canopy ecosystem (Nowacki and Abrams 2008). Succession to this forested state can occur in as little as 50 years from the last fire (LANDFIRE 2009).

#### **Dominant plant species**

- white oak (*Quercus alba*), tree
- sugar maple (*Acer saccharum*), tree
- honeysuckle (*Lonicera*), shrub
- common buckthorn (*Rhamnus cathartica*), shrub
- mayapple (*Podophyllum peltatum*), other herbaceous
- garlic mustard (*Alliaria petiolata*), other herbaceous

#### **Community 2.1**

##### **White Oak – Sugar Maple/Honeysuckle – Common Buckthorn/Mayapple – Garlic Mustard**

This community phase represents the early stages of long-term fire suppression and overbrowsing. Mature oaks and hickories are still present, but the more shade tolerant sugar maple (*Acer saccharum* Marshall) and American basswood begin to co-dominate. The tree canopy closes to 100 percent cover and basal area increases (LANDFIRE 2009). Non-native shrubs, such as honeysuckle (*Lonicera* L.) and common buckthorn (*Rhamnus cathartica* L.), can rapidly colonize. The herbaceous layer continues to support shade-tolerant species, but diversity is reduced as the fully closed canopy results in favorable conditions mostly by spring ephemerals. Grazing pressure alters species composition, allowing plants such as mayapple to increase as it is commonly avoided by deer (Gubanyi et al. 2008; Rawbinski 2008). The non-native garlic mustard (*Alliaria petiolata* (M. Bieb) begins to invade and spread rapidly due to its ability to suppress re-establishment of native species (Stinson et al. 2006; Nelson 2010).

#### **Dominant plant species**

- white oak (*Quercus alba*), tree
- sugar maple (*Acer saccharum*), tree

## Community 2.2

### Sugar Maple – American Basswood/Honeysuckle – Common Buckthorn/Garlic Mustard - litter

Sites falling into this community phase have a well-established, fire-intolerant canopy dominated by sugar maple and American basswood. Oak seedlings are virtually absent from the understory due to the lack of available light. Without recurring fire, downed woody debris and leaf litter are frequently encountered on the forest floor.

#### Dominant plant species

- sugar maple (*Acer saccharum*), tree
- American basswood (*Tilia americana*), tree
- honeysuckle (*Lonicera*), shrub
- common buckthorn (*Rhamnus cathartica*), shrub
- garlic mustard (*Alliaria petiolata*), other herbaceous
- Litter, herbaceous (*Litter, herbaceous*), other herbaceous

## Pathway 2.1A

### Community 2.1 to 2.2

Continued fire suppression.

## Pathway 2.2A

### Community 2.2 to 2.1

Pathway 2.2A – Severe disturbance event such as a replacement fire, severe drought, or windstorm.

## State 3

### Pasture State

The pasture state occurs when the reference state is converted to a farming system that emphasizes domestic livestock production known as grassland agriculture. Fire suppression, periodic cultural treatments (e.g., clipping, drainage, soil amendment applications, planting new species and/or cultivars, mechanical harvesting) and grazing by domesticated livestock transition and maintain this state (USDA-NRCS 2003). Early settlers seeded non-native species, such as smooth brome (*Bromus inermis* Leyss.) tall fescue (*Festuca arundinacea*), and Kentucky bluegrass (*Poa pratensis* L.), to help extend the grazing season. Over time, as lands were continuously harvested or grazed by herds of cattle, the non-native species were able to spread and expand across the landscape, reducing the native species diversity and ecological function. This state is most common on the steeply sloping sites.

#### Dominant plant species

- smooth brome (*Bromus inermis*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- tall fescue (*Schedonorus arundinaceus*), grass
- timothy (*Phleum pratense*), grass
- alfalfa (*Medicago*), other herbaceous
- clover (*Trifolium*), other herbaceous

## Community 3.1

### Hayfield

Sites in this community phase consist of forage plants that are planted and mechanically harvested. Mechanical harvesting removes much of the aboveground biomass and nutrients that feed the soil microorganisms (Franzluebbers et al. 2000; USDA-NRCS 2003). As a result, soil biology is reduced leading to decreases in nutrient uptake by plants, soil organic matter, and soil aggregation. Frequent biomass removal can also reduce the site's

carbon sequestration capacity (Skinner 2008).

### **Dominant plant species**

- brome (*Bromus*), grass
- tall fescue (*Schedonorus arundinaceus*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- timothy (*Phleum pratense*), grass
- clover (*Trifolium*), other herbaceous

## **Community 3.2**

### **Smooth brome -Tall fescue - Clover Pasture**

This community is characterized by seeded cool-season grass and forbs. Species will depend upon landowner goals and objectives and may include many different grasses and forbs. Common species include smooth brome (*Bromus inermis*), tall fescue (*Festuca arundinacea*), Kentucky bluegrass (*Poa pratensis*), timothy (*Phleum pratense*), red clover (*Trifolium pratense*) and white clover (*Trifolium repens* L.). Management inputs include control of weeds and brush. These sites are managed to ensure a proper forage/animal balance. Plants are not overutilized and have adequate rest and recovery.

### **Dominant plant species**

- smooth brome (*Bromus inermis*), grass
- tall fescue (*Schedonorus arundinaceus*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- white clover (*Trifolium repens*), other herbaceous
- red clover (*Trifolium pratense*), other herbaceous

## **Community 3.3**

### **Broomsedge -Thistle-Smooth brome -Tall Fescue Pasture**

Over utilization of the pasture will result in a shift to include more undesirable species such as thistle (*Cirsium* spp.), broomsedge (*Andropogon virginicus* L.), ironweed (*Vernonia gigantea*), buttercup (*Ranunculus* spp.), ragweed (*Ambrosia* spp.) and blackberries (*Rubus* spp.). Many different woody and weed species may be present depending on seed sources and level of soil disturbance. This community reflects an improper forage-to-animal balance which will negatively impact forage productivity and reproduction, soil health, and water quality. Ecological resiliency is compromised under these conditions.

### **Dominant plant species**

- broomsedge bluestem (*Andropogon virginicus*), grass
- tall fescue (*Schedonorus arundinaceus*), grass
- Kentucky bluegrass (*Poa pratensis*), grass
- thistle (*Cirsium*), other herbaceous
- buttercup (*Ranunculus*), other herbaceous
- ragweed (*Ambrosia*), other herbaceous

## **Pathway 3.1A**

### **Community 3.1 to 3.2**

Mechanical harvesting is replaced with domestic livestock grazing.

## **Pathway 3.1B**

### **Community 3.1 to 3.3**

Mechanical harvesting is replaced with domestic livestock grazing.

## **Pathway 3.2A**

## **Community 3.2 to 3.1**

Domestic livestock are removed and mechanical harvesting is implemented.

## **Pathway 3.2B**

### **Community 3.2 to 3.3**

Grazing of livestock with overutilization of the forage plants.

## **Pathway 3.3B**

### **Community 3.3 to 3.1**

Domestic livestock are removed, and mechanical harvesting is implemented.

## **Pathway P3.3A**

### **Community 3.3 to 3.2**

Forage plants are not overutilized and the site has a proper forage-to-animal balance

## **State 4**

### **Cropland State**

The continuous use of tillage, row-crop planting, and chemicals (i.e., herbicides, fertilizers, etc.) has effectively eliminated the reference community and many of its natural ecological functions in favor of crop production. Corn and soybeans are the dominant crops for the site, and oats (*Avena L.*) and alfalfa (*Medicago sativa L.*) may be rotated periodically. These areas are likely to remain in crop production for the foreseeable future.

#### **Dominant plant species**

- corn (*Zea mays*), other herbaceous
- soybean (*Glycine max*), other herbaceous

## **Community 4.1**

### **Conventional Tillage Field**

Sites in this community phase typically consist of monoculture row-cropping maintained by conventional tillage practices. They are cropped in either continuous corn or corn-soybean rotations. The frequent use of deep tillage, low crop diversity, and bare soil conditions during the non-growing season negatively impacts soil health. Under these practices, soil aggregation is reduced or destroyed, soil organic matter is reduced, erosion and runoff are increased, and infiltration is decreased, which can ultimately lead to undesirable changes in the hydrology of the watershed (Tomer et al. 2005).

#### **Dominant plant species**

- corn (*Zea mays*), other herbaceous
- soybean (*Glycine max*), other herbaceous

## **Community 4.2**

### **Conservation Tillage Field**

This community phase is characterized by rotational crop production that utilizes various conservation tillage methods to promote soil health and reduce erosion. Conservation tillage methods include strip-till, ridge-till, vertical-till, or no-till planting systems. Strip-till keeps seedbed preparation to narrow bands less than one-third the width of the row where crop residue and soil consolidation are left undisturbed in-between seedbed areas. Strip-till planting may be completed in the fall and nutrient application either occurs simultaneously or at the time of planting. Ridge-till uses specialized equipment to create ridges in the seedbed and vegetative residue is left on the surface in between the ridges. Weeds are controlled with herbicides and/or cultivation, seedbed ridges are rebuilt during cultivation, and soils are left undisturbed from harvest to planting. Vertical-till systems employ machinery that lightly tills the soil and cuts up crop residue, mixing some of the residue into the top few inches of the soil while leaving a

large portion on the surface. No-till management is the most conservative, disturbing soils only at the time of planting and fertilizer application. Compared to conventional tillage systems, conservation tillage methods can improve soil ecosystem function by reducing soil erosion, increasing organic matter and water availability, improving water quality, and reducing soil compaction.

#### **Dominant plant species**

- corn (*Zea mays*), other herbaceous
- soybean (*Glycine max*), other herbaceous

### **Community 4.3**

#### **Conservation Tillage Field/ Cover Crop Field**

This community phase applies conservation tillage methods as described above as well as adds cover crop practices. Cover crops typically include nitrogen-fixing species (e.g., legumes), small grains (e.g., rye, wheat, oats), or forage covers (e.g., turnips, radishes, rapeseed). The addition of cover crops not only adds plant diversity but also promotes soil health by reducing soil erosion, limiting nitrogen leaching, suppressing weeds, increasing soil organic matter, and improving the overall soil ecosystem. In the case of small grain cover crops, surface cover and water infiltration are increased, while forage covers can be used to graze livestock or support local wildlife. Of the three community phases for this state, this phase promotes the greatest soil sustainability and improves ecological functioning within a cropland system.

#### **Dominant plant species**

- rye (*Secale*), grass
- wheat (*Triticum*), grass
- oat (*Avena*), grass
- corn (*Zea mays*), other herbaceous
- soybean (*Glycine max*), other herbaceous
- radish (*Raphanus*), other herbaceous

### **Pathway 4.1A**

#### **Community 4.1 to 4.2**

Tillage operations are greatly reduced, crop rotation occurs on a regular interval, and crop residue remains on the soil surface.

### **Pathway 4.1B**

#### **Community 4.1 to 4.3**

Tillage operations are greatly reduced or eliminated, crop rotation occurs on a regular interval, crop residue remains on the soil surface, and cover crops are planted following crop harvest.

### **Pathway 4.2A**

#### **Community 4.2 to 4.1**

Intensive tillage is utilized, and monoculture row-cropping is established.

### **Pathway 4.2B**

#### **Community 4.2 to 4.3**

Cover crops are implemented to minimize soil erosion.

### **Pathway 4.3B**

#### **Community 4.3 to 4.1**

Intensive tillage is utilized, cover crops practices are abandoned, monoculture row-cropping is established, and crop rotation is reduced or eliminated.



## **Pathway 4.3A**

### **Community 4.3 to 4.2**

Cover crop practices are abandoned.

## **State 5**

### **Reconstructed Oak Woodland State**

The combination of natural and anthropogenic disturbances occurring today has resulted in numerous forest health issues, and restoration back to the historic reference condition may not be possible. Forests are being stressed by non-native diseases and pests, habitat fragmentation, changes in soil conditions, and overabundant deer populations on top of naturally-occurring disturbances (severe weather and native pests) (IFDC 2018). However, these habitats provide multiple ecosystem services including carbon sequestration; clean air and water; soil conservation; biodiversity support; wildlife habitat; timber, fiber, and fuel products; as well as a variety of cultural activities (e.g., hiking, camping, hunting) (Millennium Ecosystem Assessment 2005; IFDC 2018). Therefore, conservation of forests and woodlands should still be pursued. Forest reconstructions are an important tool for repairing natural ecological functioning and providing habitat protection for numerous species associated with Loess Exposed Backslope Forests. Therefore, ecological restoration should aim to aid the recovery of degraded, damaged, or destroyed ecosystems. A successful restoration will have the ability to structurally and functionally sustain itself, demonstrate resilience to the ranges of stress and disturbance, and create and maintain positive biotic and abiotic interactions (SER 2002). The reconstructed oak forest state is the result of a long-term commitment involving a multi-step, adaptive management process.

#### **Dominant plant species**

- oak (*Quercus*), tree

## **Community 5.1**

### **Early Successional Oak Woodland**

This community phase represents the early community assembly from woodland reconstruction. It is highly dependent on the current condition of the woodland based on past and current land management actions, invasive species, and proximity to land populated with non-native pests and diseases. Therefore, no two sites will have the same early successional composition. Technical forestry assistance should be sought to develop suitable conservation management plans.

#### **Dominant plant species**

- oak (*Quercus*), tree

## **Community 5.2**

### **Late Successional Oak Woodland**

Appropriately timed management practices (e.g., prescribed fire, hazardous fuels management, forest stand improvement, continuing integrated pest management) applied to the early successional community phase can help increase the stand maturity, pushing the site into a late successional community phase over time. A late successional reconstructed forest will have an uneven-aged canopy and a well-developed shrub layer and understory.

#### **Dominant plant species**

- oak (*Quercus*), tree

## **Pathway 5.1A**

### **Community 5.1 to 5.2**

Management plan implemented successfully.

## **Pathway 5.2A**

### **Community 5.2 to 5.1**

Reconstruction experiences a setback from extreme weather event or improper timing of management actions.

#### **Transition T1A**

##### **State 1 to 2**

Transition 1A – Long-term fire suppression in excess of 50 years transitions the site to the fire-suppressed state (2).

#### **Transition T1B**

##### **State 1 to 3**

Transition 1B – Cultural treatments to enhance forage quality and yield transitions the site to the pasture state (3).

#### **Transition T1C**

##### **State 1 to 4**

Agricultural conversion via tillage, seeding, and non-selective herbicide

#### **Transition T2A**

##### **State 2 to 3**

Transition 2A – Cultural treatments to enhance forage quality and yield transitions the site to the pasture state (3).

#### **Transition T2B**

##### **State 2 to 4**

Transition 2B – Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (4).

#### **Transition T3A**

##### **State 3 to 2**

Transition 3A – Land abandonment transitions the site to the fire-suppressed state (2).

#### **Restoration pathway T3B**

##### **State 3 to 4**

Transition 3B – Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (4).

#### **Restoration pathway R3A**

##### **State 3 to 5**

Restoration 3A – Site preparation, tree planting, invasive species control, and seeding native species transition this site to the reconstructed oak woodland state (5).

#### **Transition T4A**

##### **State 4 to 2**

Transition 4A – Land abandonment transitions the site to the fire-suppressed state (2).

#### **Transition T4B**

##### **State 4 to 3**

Transition 4B – Cultural treatments to enhance forage quality and yield transitions the site to the pasture state (3).

## **Restoration pathway R4A**

### **State 4 to 5**

Restoration 4A – Site preparation, tree planting, invasive species control, and seeding native species transition this site to the reconstructed oak woodland state (5).

## **Transition T5A**

### **State 5 to 2**

Transition 5A – Fire suppression and removal of active management transitions this site to the fire-suppressed state (2).

## **Transition T5B**

### **State 5 to 3**

Transition 5B – Cultural treatments to enhance forage quality and yield transition the site to the pasture state (3).

## **Transition T5C**

### **State 5 to 4**

Transition 5C – Tillage, seeding of agricultural crops, and non-selective herbicide transition this site to the cropland state (4).

## **Additional community tables**

### **Inventory data references**

No field plots were available for this site. A review of the scientific literature and professional experience were used to approximate the plant communities for this provisional ecological site. Information for the state-and-transition model was obtained from the same sources. All community phases are considered provisional based on these plots and the sources identified in this ecological site description.

### **Other references**

Angel, J. No date. Climate of Illinois Narrative. Illinois State Water Survey, Prairie Research Institute, University of Illinois at Urbana-Champaign. Available at <https://www.isws.illinois.edu/statecli/General/Illinois-climate-narrative.htm>. Accessed 8 November 2018.

Barrett, S.W. 1984. Indians and fire. *Western Wildlands*. Spring: 17-20.

Bharati, L., K.-H. Lee, T.M. Isenhardt, and R.C. Schultz. 2002. Soil-water infiltration under crops, pasture, and established riparian buffer in Midwestern USA. *Agroforestry Systems* 56: 249-257.

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. *Ecological Subregions: Sections and Subsections of the Conterminous United States*. USDA Forest Service, General Technical Report WO-76. Washington, DC. 92 pps.

Flickinger, A. 2010. *Iowa Forests Today: An Assessment of the Issues and Strategies for Conserving and Managing Iowa's Forests*. Iowa Department of Natural Resources. 329 pps.

Franzluebbers, A.J., J.A. Stuedemann, H.H. Schomberg, and S.R. Wilkinson. 2000. Soil organic C and N pools under long-term pasture management in the Southern Piedmont USA. *Soil Biology and Biochemistry* 32:469-478.

Gubanyi, J., J. Savidge, S.E. Hygnstrom, K. VerCauteren, G.W. Garabrandt, and S. Korte. 2008. Deer impact on vegetation in natural areas in southeastern Nebraska. USDA National Wildlife Research Center – Staff Publications. 913. Available at [http://digitalcommons.unl.edu/icwdm\\_usdanwrc/913](http://digitalcommons.unl.edu/icwdm_usdanwrc/913). (Accessed 6 April 2017).

- Illinois Forestry Development Council (IFDC). 2018. Illinois Forest Action Plan: A Statewide Forest Resource Assessment and Strategy, Version 4.1. Illinois Forestry Development Council and Illinois Department of Natural Resources. 80 pps.
- Irland, L.C. 2000. Ice storms and forest impacts. *The Science of the Total Environment* 262:231-242.
- LANDFIRE. 2009. Biophysical Setting 4213100 North-Central Interior Dry-Mesic Oak Forest and Woodland. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.
- Leake, J., D. Johnson, D. Donnelly, G. Muckle, L. Boddy, and D. Read. 2004. Networks of power and influence: the role of mycorrhizal mycelium in controlling plant communities and agroecosystem functioning. *Canadian Journal of Botany* 82: 1016-1045.
- Millennium Ecosystem Assessment. 2005. Ecosystems and Human Well-Being: Current States and Trends. World Resources Institute. Island Press, Washington, D.C. 948 pages.
- NatureServe. 2018. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1 NatureServe, Arlington, VA. Available at <http://explorer.natureserve.org>. (Accessed 4 December 2019).
- Nelson, P. 2010. The Terrestrial Natural Communities of Missouri. Missouri Department of Natural Resources, Missouri Natural Areas Committee. 550 pps.
- Nowacki, G.J. and M.D. Abrams. 2008. The demise of fire and “mesophication” of forests in the eastern United States. *BioScience* 58: 123-138.
- Peel, M.C., B.L. Finlayson, and T.A. McMahon. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11: 1633-1644.
- Peterson, C.J. 2000. Catastrophic wind damage to North American forests and the potential impact of climate change. *The Science of the Total Environment* 262: 287-311.
- Pyne, S.J., P.L. Andrews, and R.D. Laven. 1996. Introduction to Wildland Fire, Second Edition. John Wiley and Sons, Inc. New York, New York. 808 pps.
- Rawbinski, T.J. 2008. Impacts of White-tailed Deer Overabundance in Forest Ecosystems: An Overview. U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. Newton Square, PA, USA. Available at [https://www.na.fs.fed.us/fhp/special\\_interests/White-tailed\\_deer.pdf](https://www.na.fs.fed.us/fhp/special_interests/White-tailed_deer.pdf) (Accessed 17 April 2017).
- Schwegman, J.E., G.B. Fell, M. Hutchinson, G. Paulson, W.M. Shepherd, and J. White. 1973. Comprehensive Plan for the Illinois Nature Preserves System, Part 2 The Natural Divisions of Illinois. Illinois Nature Preserves Commission, Rockford, IL. 32 pps.
- Skinner, R.H. 2008. High biomass removal limits carbon sequestration potential of mature temperate pastures. *Journal for Environmental Quality* 37: 1319-1326.
- Society for Ecological Restoration [SER] Science & Policy Working Group. 2002. The SER Primer on Ecological Restoration. Available at: <http://www.ser.org/>. (Accessed 28 February 2017).
- Stinson, K.A., S.A. Campbell, J.R. Powell, B.E. Wolfe, R.M. Callaway, G.C. Thelen, S.G. Hallett, D. Prati, and J.N. Klironomos. 2006. Invasive plant suppresses the growth of native tree seedlings by disrupting belowground mutualisms. *PLOS Biology* 4: 0727-0731.
- Taft, J.B., G.S. Wilhelm, D.M. Ladd, and L.A. Masters. 1997. Floristic Quality Assessment for vegetation in Illinois, a method for assessing vegetation integrity. *Erigenia* 15: 3-95.
- Taft, J.B., R.C. Anderson, L.R. Iverson, and W.C. Handel. 2009. Chapter 4: Vegetation ecology and change in terrestrial ecosystems. In: C.A. Taylor, J.B. Taft, and C.E. Warwick (eds.). *Canaries in the Catbird Seat: The Past,*

Present, and Future of Biological Resources in a Changing Environment. Illinois Natural Heritage Survey Special Publication 30, Prairie Research Institute, University of Illinois at Urbana-Champaign. 306 pps.

Teague, W.R., S.L. Dowhower, S.A. Baker, N. Haile, P.B. DeLaune, and D.M. Conover. 2011. Grazing management impacts on vegetation, soil biota and soil chemical, physical and hydrological properties in tall grass prairie. *Agriculture, Ecosystems and Environment* 141: 310-322.

Tomer, M.D., D.W. Meek, and L.A. Kramer. 2005. Agricultural practices influence flow regimes of headwater streams in western Iowa. *Journal of Environmental Quality* 34:1547-1558.

Undersander, D., B. Albert, D. Cosgrove, D. Johnson, and P. Peterson. 2002. *Pastures for Profit: A Guide to Rotational Grazing* (A3529). University of Wisconsin-Extension and University of Minnesota Extension Service. 43 pps.

United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2003. *National Range and Pasture Handbook, Revision 1*. Grazing Lands Technology Institute. 214 pps.

United States Department of Agriculture – Natural Resource Conservation Service (USDA-NRCS). 2006. *Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin*. U.S. Department of Agriculture Handbook 296. 682 pps.

United States Department of Agriculture, Natural Resources Conservation Service. 2022. *Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin*. U.S. Department of Agriculture, Agriculture Handbook 296.

U.S. Environmental Protection Agency [EPA]. 2013. *Level III and Level IV Ecoregions of the Continental United States*. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1:3,000,000. Available at <http://www.epa.gov/eco-research/level-iii-andiv-ecoregions-continental-united-states>. (Accessed 1 March 2017).

VerCauteren, K. and S.E. Hygnstrom. 2011. Managing white-tailed deer: Midwest North America. *Papers in Natural Resources*. Paper 380. Available at <http://digitalcommons.unl.edu/natrespapers/380>. (Accessed 17 April 2017).

White, J. and M.H. Madany. 1978. Classification of natural communities in Illinois. In: J. White. *Illinois Natural Areas Inventory Technical Report*. Illinois Natural Areas Inventory, Department of Landscape Architecture, University of Illinois at Urbana/Champaign. 426 pps.

## **Contributors**

Lisa Kluesner  
Rick Francen

## **Approval**

Suzanne Mayne-Kinney, 12/30/2024

## **Acknowledgments**

This project could not have been completed without the dedication and commitment from a variety of staff members. Team members supported the project by serving on the technical team, assisting with the development of state and community phases of the state-and-transition model, providing peer review and technical editing, and conducting quality control and quality assurance reviews.

List of primary contributors and reviewers.

Organization	Name	Title	Location
--------------	------	-------	----------

Iowa Department of Natural Resources	Kevin Andersen	State Private Lands Biologist	Fairfield, IA
--------------------------------------	----------------	-------------------------------	---------------

Natural Resources Conservation Service	Patrick Chase	State Soil Scientist	Des Moines, IA
--	---------------	----------------------	----------------

Ron Collman State Soil Scientist Champaign, IL  
Tonie Endres Senior Regional Soil Scientist Indianapolis, IN  
Rick Francen Soil Scientist Springfield, IL  
Lisa Kluesner Ecological Site Specialist Waverly, IA  
Jorge, Lugo-Camacho State Soil Scientist Columbia, MO  
Kevin Norwood Soil Survey Regional Director Indianapolis, IN  
Stanley Sipp Resource Inventory Specialist Champaign, IL  
Jason Steele Area Resource Soil Scientist Fairfield, IA  
Chris Tecklenberg Acting Regional Ecological Site Specialist Hutchinson, KS  
Doug Wallace ACES Ecologist Columbia, MO

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Lisa Kluesner
Contact for lead author	
Date	01/09/2025
Approved by	Suzanne Mayne-Kinney
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 
17. **Perennial plant reproductive capability:**
-