

## Ecological site NX117X01Y034 Shallow Upland

Last updated: 9/22/2023  
Accessed: 05/03/2024

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 117X–Boston Mountains

This MLRA is about 6,850 square miles (17,775 square kilometers) and is in Arkansas (82 percent) and Oklahoma (18 percent). The Ozark National Forest makes up a significant portion of the MLRA.

This area is predominantly in the Boston Mountains Section of the Ozark Plateaus Province of the Interior Highlands. This MLRA marks the southern extent of the Ozarks, a deeply eroded plateau. Ridges of the MLRA are narrow and rolling, and valley sides are steep. Elevation ranges from 570 feet (170 meters) to 1,860 feet (560 meters). Some lower and higher areas are on valley floors and crests of ridges, respectively.

Level to slightly tilted shale, sandstone, and siltstone strata of geologic formations from the Pennsylvanian Subperiod underlie most of the MLRA. These formations include the Atoka Formation, the Bloyd Shale, and the Cane Hill and Prairie Grove Members of the Hale Formation. The Pitkin Limestone, Fayetteville Shale, and Batesville Sandstone Formations from the Mississippian Subperiod underlie parts of the northern edge of the MLRA. Alluvial deposits are in river valleys and consist of an unconsolidated mixture of clay, silt, sand, and gravel.

The MLRA has a thermic soil temperature regime and a udic soil moisture regime. The dominant soil orders are Ultisols and Inceptisols. The soils are loamy and have mixed or siliceous mineralogy. They are shallow to very deep and are well drained. Hapludults and Dystrudepts formed in residuum on hills, plateaus, and mountains. Paleudults, on terraces and hills, formed in old alluvium over residuum or colluvium over residuum.

### Ecological site concept

The Shallow Upland Ecological Site is on hills and mountains, specifically hillslopes and mountain slopes. The soils associated with this site are less than 20 inches (50 cm) deep, are well drained, and formed in residuum and colluvium from sandstone. Slope ranges from 4 to 40 percent, and elevation ranges from 500 to 2,500 feet. The soils have less than 35 percent clay in the particle-size control section.

### Associated sites

NX117X01Y033	<b>Loamy Upland</b> This site is on mountains and hills, specifically mountain slopes and hillslopes. The soils associated with the site are greater than 20 inches deep.
--------------	--

### Similar sites

NX117X01Y031	<b>Clayey Upland</b> This site is on mountains and hills, specifically hillslopes and mountain slopes. The soils associated with the site have 35 to 60 percent clay in the particle-size control section, and there is a layer of clay accumulation within eight inches of the soil surface.
--------------	--

**Table 1. Dominant plant species**

Tree	(1) <i>Quercus</i> (2) <i>Cornus</i>
Shrub	(1) <i>Vaccinium</i> (2) <i>Hamamelis</i>
Herbaceous	(1) <i>Schizachyrium scoparium</i> (2) <i>Sorghastrum nutans</i>

## Legacy ID

R117XY034AR

## Physiographic features

This ecological site is on hills and mountains, specifically hillslopes and mountain slopes. Slope ranges from 4 to 40 percent, and elevation ranges from 500 to 2,500 feet (150 to 762 meters). The runoff class is high or very high. The site is not subject to ponding or flooding.

**Table 2. Representative physiographic features**

Landforms	(1) Mountains > Mountain slope (2) Hills > Hillslope
Runoff class	High to very high
Flooding frequency	None
Ponding frequency	None
Elevation	152–853 m
Slope	4–40%
Aspect	Aspect is not a significant factor

## Climatic features

Hot summers, cool winters, and mild spring and fall temperatures are typical for the ecological site. The mean annual precipitation is 51 inches. The average frost-free period is 165 days, and the average freeze-free period is 191 days. The highest precipitation occurs in May (6.2 inches), and the lowest occurs in January (3.0 inches). The warmest month of the year is August (91°F average high), and the coolest is January (25°F average low). Thunderstorms and heat waves are common and occur frequently during the summer months.


Occasionally, catastrophic storm events, such as tornados, ice storms, floods, and hailstorms, will occur. According to the Oklahoma Water Resource Board, drought occurs every 5 to 10 years (Oklahoma Water Resources Board, 2022). The EPA predicts droughts will become more severe throughout Arkansas due to longer periods without rain and an increase in very hot days (EPA, 2016).

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/>. The Deer, Fayetteville, Huntsville, Mountainburg, and Mountain View climate stations provided general climate data. Site-specific climate data is available through the National Weather Service.


**Table 3. Representative climatic features**

Frost-free period (characteristic range)	160-179 days
Freeze-free period (characteristic range)	191-204 days
Precipitation total (characteristic range)	1,219-1,346 mm
Frost-free period (actual range)	159-182 days


Freeze-free period (actual range)	186-207 days
Precipitation total (actual range)	1,194-1,397 mm
Frost-free period (average)	171 days
Freeze-free period (average)	197 days
Precipitation total (average)	1,295 mm




**Figure 1. Monthly precipitation range**




**Figure 2. Monthly minimum temperature range**




**Figure 3. Monthly maximum temperature range**



**Figure 4. Monthly average minimum and maximum temperature**



**Figure 5. Annual precipitation pattern**



**Figure 6. Annual average temperature pattern**

### Climate stations used

- (1) MTN VIEW [USC00035046], Mountain View, AR
- (2) DEER [USC00031900], Deer, AR
- (3) HUNTSVILLE 1 SSW [USC00033544], Huntsville, AR
- (4) FAYETTEVILLE DRAKE FLD [USW00093993], Fayetteville, AR
- (5) MOUNTAINBURG 2 NE [USC00035018], Mountainburg, AR

### Influencing water features

Water features do not significantly influence this ecological site.

### Wetland description

Wetlands do not significantly influence this ecological site.

### Soil features

The soils associated with this ecological site formed in residuum from sandstone, siltstone, and shale. The soils are

shallow, are well drained, and have a moderately rapid permeability class. The surface texture is commonly gravelly or stony fine sandy loam. There is a restrictive layer less than 20 inches (50cm) from the surface. The soils have less than 35 percent clay and greater than 35 percent rock fragment by volume in the particle-size control section.

The soil series associated with this site are the Hector, Mountainburg, and Montevallo Series.

**Table 4. Representative soil features**

Parent material	(1) Residuum—sandstone and shale (2) Siltstone
Surface texture	(1) Gravelly fine sandy loam (2) Stony sandy loam
Drainage class	Well drained
Permeability class	Moderately rapid
Depth to restrictive layer	51 cm
Soil depth	25–51 cm
Surface fragment cover ≤3"	9–15%
Surface fragment cover >3"	15–28%
Available water capacity (Depth not specified)	1.52–4.57 cm
Soil reaction (1:1 water) (Depth not specified)	4.5–6
Subsurface fragment volume ≤3" (Depth not specified)	16–29%
Subsurface fragment volume >3" (Depth not specified)	16–41%

## Ecological dynamics

The Reference State for this ecological site consists of a hardwood forest characterized by mixed hardwood species and a vegetative ground cover. The characteristic tree species for this state are hardwoods such as oak, beech, and hickory (Foti, 2004). Native grass species such as big bluestem, switchgrass, little bluestem, and Indiangrass grow in canopy openings and throughout the understory (Owen, 1858).

Fire significantly influences this ecological site. Historically, the average fire-return interval was likely between 3 and 25 years (Guyette and Spetich, 2003; Hallgren et al., 2012). Some of these wildfires occurred naturally through lightning strikes, but human activities probably caused most of these fires (DeSantis et al., 2010). Native species evolved with and responded well to fire (Spetich and He, 2008; Engle and Bidwell, 2001). Fires on similar upland ecological sites today are likely of moderate to low severity due to forested conditions and lower amounts of ground vegetation (Carey, 1992).


Climate-related events, such as hailstorms, tornados, thunderstorms, and extreme precipitation, occur on this site. Hailstorms can reduce canopy size, increase litter deposition, and increase tree bark removal. When paired with other disturbances, such as fire, the effects on tree species were much greater than in areas not affected by the hailstorm (Gower et al., 2015). Tornados can change plant community compositions in savanna ecosystems, favoring hardwoods and eliminating softwoods (Liu et al., 1997). Lightning storms greatly affect ecosystems. Lightning storms generally occur during summer months but can occur during any season. If a fire starts with a lightning strike, the effects on the ecosystem vary depending on the season (Hiers et al., 2000).

Grazing and farming are feasible for this ecological site. Changes to the ecological dynamics are proportional to the intensity of livestock grazing and can accelerate through overgrazing (Angerer et al., 2016). For example, desirable grasses and forbs repeatedly grazed by livestock become weak and can die, and less desirable species may replace preferable ones (Smith, 1940).

The state-and-transition model for the Shallow Upland Ecological Site consists of three identified states: Reference, Encroached, and Pasture. Because of sparse data availability, the model only explored basic principles and included only a small number of species. Further data collection from this ecological site would provide a greater understanding of ecological form and function and of resource consumption and distribution.

## State and transition model

### Ecosystem states



**T1A** - Woody species encroachment


**T1B** - Tree removal Mechanical and chemical suppression of woody vegetation Tillage Introduction of annual or perennial forage species

**R2A** - Natural regeneration and disturbance regimes

**T2A** - Tree removal Mechanical and chemical suppression of woody vegetation Tillage Introduction of annual or perennial forage species

**T3B** - Lack of management Abandonment


### State 1 submodel, plant communities



**1.1A** - Absence of fire Woody species growth Excessive grazing pressure

**1.2A** - Wildfire or other disturbance that reduces woody canopy


### State 2 submodel, plant communities



**2.1A** - Fire suppression

**2.2A** - Excessive fire Mechanical tree removal.

### State 3 submodel, plant communities



**3.1A** - Fire suppression

**3.2A** - Tree removal Brush management

**3.2B** - Fire suppression

**3.3A** - Tree removal Brush management

**3.3B** - Tree removal Brush management

## State 1 Reference

The Reference State represents the natural range of variability for the ecological site without major human influences. The main drivers for community pathways within the Reference State are fire frequency intervals between 3 and 25 years (Guyette and Spetich, 2003; Hallgren et al., 2012), climate effects (decadal scale), insect or disease presence or establishment, and wildlife grazing or browsing. Fire is the main feedback mechanism within this state, and fire-tolerant species dominate the ecological site. Fire intervals suppress vegetation growth; wildlife grazing or browsing reduces the amount of grass available. Reduced grass availability lessens fire intensity and causes wildlife migration.

**Characteristics and indicators.** The Reference State consists of a hardwood forest characterized by mixed hardwoods and sporadic gaps where other vegetation grows. Hardwood species include oak, beech, and hickory (Foti, 2004). Native grass species, such as big bluestem, switchgrass, little bluestem, and Indiangrass, grow in canopy openings and throughout the understory (Arkansas Geological Survey, 2005).

### Dominant plant species

- oak (*Quercus*), tree
- beech (*Fagus*), tree
- hybrid hickory (*Carya*), tree
- big bluestem (*Andropogon gerardii*), grass
- little bluestem (*Schizachyrium*), grass
- Indiangrass (*Sorghastrum*), grass
- switchgrass (*Panicum virgatum*), grass

## Community 1.1 Post Oak/Little Bluestem

Oak trees and warm-season, perennial tallgrasses dominate this community phase. Dominant grasses are little bluestem, big bluestem, Indiangrass, and switchgrass. Common trees include post oak, blackjack oak, white oak, and red oak.

## Community 1.2

## Post Oak-Eastern Redcedar (At-Risk Community)

This community phase has a moderately closed canopy and an understory of tallgrasses and midgrasses. The absence of fire allows post oak, blackjack oak, and eastern redcedar densities to increase. Competition from the denser canopy leads to a reduction in the herbaceous understory.

### Pathway 1.1A Community 1.1 to 1.2

The main drivers for this community pathway are the absence of fire and the growth of woody species. Excessive grazing pressure may also accompany this transition.

### Pathway 1.2A Community 1.2 to 1.1

The main drivers for this community pathway are periodic wildfires that reduce the amount of woody vegetation. Drought and outbreaks of insects, diseases, or both may also result in a reduced woody canopy.

#### Conservation practices

Prescribed Burning
Prescribed Grazing

## State 2 Encroached

The main drivers for community pathways within the Encroached State are the absence of wildfire, seed dispersal by wildlife, climate effects (decadal scale), and canopy density. The main feedback mechanism for community pathways in this state is the dominance of woody species. Woody species shade and outcompete herbaceous species. Fire intensity diminishes, and woody species take control of nutrient and water cycling.

**Characteristics and indicators.** The Encroached State consists of many woody species (especially eastern redcedar) and significant canopy closure. Time and fire intensity determine the community phases and species abundance and variation. As the woody canopy increases and the Encroached State develops, the hydrology of the ecological site alters. The denser canopy intercepts most precipitation and changes hydrological patterns to favor woody species over herbaceous species. Understory species, therefore, have less available water for growth and must compete with an extensive overstory root system (Zou et al., 2018).

#### Dominant plant species

- eastern redcedar (*Juniperus virginiana*), tree
- oak (*Quercus*), tree
- hybrid hickory (*Carya*), tree
- beech (*Fagus*), tree

### Community 2.1 Post Oak-Eastern Redcedar

This community phase consists of oak, hickory, beech, and eastern redcedar. The canopy cover increases, thereby causing a reduction in the herbaceous ground cover. Eastern redcedar increases in size and quantity.

### Community 2.2 Eastern Redcedar-Post Oak

Eastern redcedar dominates this community phase. Oak, hickory, and beech trees may be present. Oak, hickory, and beech trees experience reduced vigor and reproductive capacity due to shading and competition from eastern redcedar.



## **Pathway 2.1A**

### **Community 2.1 to 2.2**

The absence of wildfire is the main driver for this community pathway and contributes to natural regeneration and increased woody vegetation growth.

## **Pathway 2.2A**

### **Community 2.2 to 2.1**

The main driver for this community pathway is a reduction in canopy cover. As canopy cover decreases, more sunlight reaches the ground, and herbaceous vegetation increases. Increased herbaceous vegetation can lead to fires that reduce woody vegetation. Excessive fire and the mechanical removal of trees contribute to reduced canopy cover.

## **State 3**

### **Pasture**

The Pasture State consists of introduced herbaceous species planted to maximize livestock forage production. The main drivers for community pathways within the Pasture State are the mechanical disturbance of the soil and seed planting, climate effects (decadal scale), seed dispersal, and wildlife and livestock grazing or browsing. The main feedback mechanism for community pathways in this state is the use of mechanical equipment and chemicals to increase forage. Fertilizer inputs and brush management are essential for maintaining high productivity. Wildlife or livestock grazing or browsing reduces the amount of available forage.

**Characteristics and indicators.** The Pasture State consists of species grown for specific management goals, mainly livestock grazing. Common pasture species include buffalograss, western wheatgrass, little bluestem, sideoats grama, Bermudagrass, and bahiagrass. The quality and quantity of forbs, grasses, and legume species within this state depend on the level of management inputs (seeding, weed management, and land use). Both warm-season and cool-season grasses are suitable for this ecological site.

#### **Dominant plant species**

- Bermudagrass (*Cynodon*), grass
- sideoats grama (*Bouteloua curtipendula*), grass
- bahiagrass (*Paspalum notatum*), grass

## **Community 3.1**

### **Bermudagrass**

This community phase consists of herbaceous species planted to maximize forage production for grazing livestock.

## **Community 3.2**

### **Bermudagrass/ Post Oak-Eastern Redcedar**

This community phase consists of herbaceous species planted to maximize forage production for grazing livestock. Improper pasture management allows for the growth of woody species, such as eastern redcedar and oaks.

## **Community 3.3**

### **Eastern Redcedar-American Elm-Post Oak/ Bermudagrass**

This community phase consists of herbaceous species planted to maximize forage production for grazing livestock. Improper pasture management over time allows the growth of woody species, such as eastern redcedar, oaks, and American elm. Woody species encroach on the pasture and dominate resources previously used by forage species.

## **Pathway 3.1A**

### **Community 3.1 to 3.2**

The main drivers for this community pathway are the absence of fire, improper management, and natural

regeneration.

### **Pathway 3.2A** **Community 3.2 to 3.1**

The main driver for this community pathway is the removal or reduction of woody species.

### **Pathway 3.2B** **Community 3.2 to 3.3**

The main drivers for this community pathway are the absence of fire, improper management, and natural regeneration.

### **Pathway 3.3A** **Community 3.3 to 3.1**

The main driver for this community pathway is the removal or reduction of woody species.

### **Pathway 3.3B** **Community 3.3 to 3.2**

The main driver for this community pathway is the removal or reduction of woody species.

### **Transition T1A** **State 1 to 2**

The main trigger for this transition is the absence of wildfire, which allows woody species to grow and compete for nutrients, water, and sunlight. The main slow variable for this transition is the increased competition for sunlight, nutrients, and moisture resources. Increased overstory competition reduces the vigor and reproductive capacity of the herbaceous understory. The main threshold for this transition is a shift in nutrient cycling (from grass and leaf dominance to leaf and needle dominance). Increased woody canopy cover alters hydrologic cycles, potentially reducing runoff and infiltration and increasing precipitation interception by woody species.

### **Transition T1B** **State 1 to 3**

The main triggers for this transition are tree removal, mechanical and chemical suppression of woody vegetation, tillage, and the introduction of annual or perennial forage species. The main slow variables for this transition are increased production and management of forage species. The main thresholds for this transition are changes in soil properties, such as structure, organic matter, and nutrient cycling, and changes in type and frequency of disturbance.

#### **Conservation practices**

Brush Management
Prescribed Burning
Land Clearing
Prescribed Grazing

### **Restoration pathway R2A** **State 2 to 1**

Restoration efforts for this pathway begin with mechanical and chemical treatment of undesirable woody vegetation and seeding native species. A grazing management plan and the reintroduction of historical disturbance regimes must accompany these initial treatments. Returning to a historical fire interval through prescribed burning helps to suppress woody vegetation and manage invasive species.

## Conservation practices

Brush Management
Prescribed Burning

## Transition T2A State 2 to 3

The main triggers for this transition are tree removal, mechanical and chemical suppression of woody vegetation, tillage, and the introduction of annual or perennial forage species. The main slow variables for this transition are increased production and management of forage species. The main thresholds for this transition are changes in soil properties, such as structure, organic matter, and nutrient cycling, and changes in type and frequency of disturbance.

## Conservation practices

Brush Management
Prescribed Burning
Land Clearing
Prescribed Grazing

## Transition T3B State 3 to 2

The main trigger for this transition is the lack of management or abandonment. The main slow variables for this transition are increased establishment and increased size of woody species. The main threshold for this transition is woody species dominance. Woody species dominate ecological processes, causing shading and increased competition for soil moisture, nutrients, and sunlight. Woody species dominance reduces the vigor and reproductive capacity of herbaceous species in the understory.

## Additional community tables

### Animal community

Common wildlife species in this MLRA include whitetail deer, coyote, red fox, gray fox, bobcat, beaver, raccoon, opossum, skunk, muskrat, mink, cottontail, fox squirrel, gray squirrel, bobwhite quail, and mourning dove.

### Hydrological functions

The following are estimated withdrawals of freshwater by use in this MLRA:

Public supply—surface water, 24.4%; ground water, 5.1%  
Livestock—surface water, 8.1%; ground water, 0.6%  
Irrigation—surface water, 0.0%; ground water, 0.0%  
Other—surface water, 61.8%; ground water, 0.0%

Total withdrawals average 95 million gallons per day (360 million liters per day). About 6 percent is from groundwater sources, and 94 percent is from surface-water sources. The moderately high precipitation is adequate for crops and pasture. Large reservoirs on a few major streams supply water for municipal purposes, aid in flood control, and provide recreation opportunities. The surface-water quality is generally good, and the water is suitable for most uses. Shallow wells are the principal sources of water for domestic use. Deep wells are necessary for obtaining moderate to large quantities of ground water. Water from the Ozark aquifer system in the northern half of this MLRA is suitable for drinking.

### Recreational uses

In this MLRA, mountain biking, camping, fishing, hiking, horseback riding, hunting, mineral prospecting, nature viewing, off-highway vehicle riding, and water activities are available where permitted on public land, and where allowed on private land. The Ozark National Forest makes up a significant portion of this MLRA.

## **Wood products**

Public and private timberland cover large areas throughout this MLRA. Loblolly pine is the most popular species to harvest and provides timber for lumber, pulpwood, posts, and poles. Hardwood species provide timber for lumber, flooring, and pulpwood.

## **Other products**

Poultry production is a major industry throughout the MLRA. Small grains, soybeans, and hay are major crops.

## **References**

- Angerer, J., W. Fox, and J. Wolfe. 2016. Land Degradation in Rangeland Ecosystems. Biological and Environmental hazards, Risks, and Disasters. Academic Press.
- Carey, J. 1992. *Quercus stellata*, Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. Fire Sciences Laboratory.
- DeSantis, R.D., S.W. Hallgreen, and D.W. Stahle. 2010. Historic Fire Regime of an Upland Oak Forest in South Central North America. Fire Ecology. USDA Forest Service, Northern Research Station, Saint Paul, Minnesota.
- Engle, D. and T. Bidwell. 2001. The response of central North American prairies to seasonal fire. Range Management 54:2–10.
- Foti, T. 2004. Upland Hardwood Forests and Related Communities of the Arkansas Ozarks in the Early 19th Century. Southern Research Station. U.S. Department of Agriculture, Asheville, NC. 21–29.
- Gower, K., J. Fontaine, C. Birnbaum, and N. Enright. 2015. Sequential Disturbance Effects of Hailstorms and Fire on Vegetation in a Mediterranean-Type Ecosystem. Ecosystems 18:1121–1134.
- Guyette, R.P. and M. A. Spetich. 2003. Fire History of Oak-Pine Forests in the Lower Boston Mountains, Arkansas, USA. Forest Ecology and Management. Elsevier. 463–474.
- Hallgren, S.W., DeSantis, R. D., and J.A. Burton. 2012. Fire and vegetation Dynamics in the Cross Timbers Forests of South-Central North America. Proceedings of the 4th Fire in Eastern Oak Forests Conference. USDA Forest Service General Technical Report NRS-P-102, Springfield, Missouri. 52–66.
- Hiers, K., R. Wyatt, and R. Mitchell. 2000. The effects of fire regime on legume reproduction in longleaf pine savannas: is a season selective?. Oecologia 125:521–530.
- Liu, C., J. Glitzenstein, P. Harcombe, and R. Knox. 1997. Tornado and fire effects on tree species composition in a savanna in the Big Thicket National Preserve, southeast Texas, USA. Forest Ecology and Management 91:279–289.
- Oklahoma Water Resource Board. September 2012 (Date accessed). Hydrologic Drought of Water Year 2011: A

Historical Context.

Owen, D. 1858. First Geological Reconnaissance Of The Northern Counties Of Arkansas Made During The Years 1857 And 1858. Johnson & Yerkes, State Printers, Little Rock, AR.

Smith, C. 1940. The Effects of Overgrazing and Erosion Upon the Biota of the Mixed-Grass Prairie of Oklahoma. Ecology. Wiley. 381–397.

Spetich, M. and H. He. 2008. Oak decline in the Boston Mountains, Arkansas, USA: Spatial and temporal patterns under two fire regimes. Forest Ecology and Management 254:454–462.

United States Environmental Protection Agency. 2016. What climate change means for Arkansas. Report EPA 430-F-16-006. U.S. Environmental Protection Agency.

Zou, C., D. Twidwell, and C. Bielski. 2018. Impact of Eastern Redcedar Proliferation on Water Resources in the Great Plains USA- Current State of Knowledge.

### **Other references**

Arkansas Soil Survey  
Ouachita National Forest  
Arkansas State Parks  
The Nature Conservancy  
US Fish and Wildlife Service  
Encyclopedia of Arkansas  
United States Forest Service Southern Research Station  
NatureServe  
Oklahoma Water Resource Board  
National Centers for Environmental Information  
University of Arkansas  
Oklahoma State University  
Arkansas Department of Forestry  
Oklahoma Department of Forestry

### **Contributors**

Trevor Crandall, Ecological Site Specialist

### **Approval**

Bryan Christensen, 9/22/2023

### **Acknowledgments**

Larry Gray  
Elizabeth Gray  
Erin Hourihan

### **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be

known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/03/2024
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**  
\_\_\_\_\_
2. **Presence of water flow patterns:**  
\_\_\_\_\_
3. **Number and height of erosional pedestals or terracettes:**  
\_\_\_\_\_
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**  
\_\_\_\_\_
5. **Number of gullies and erosion associated with gullies:**  
\_\_\_\_\_
6. **Extent of wind scoured, blowouts and/or depositional areas:**  
\_\_\_\_\_
7. **Amount of litter movement (describe size and distance expected to travel):**  
\_\_\_\_\_
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**  
\_\_\_\_\_
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**  
\_\_\_\_\_
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**  
\_\_\_\_\_
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be**

mistaken for compaction on this site):

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 

14. **Average percent litter cover (%) and depth ( in):**
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 

17. **Perennial plant reproductive capability:**
-