

Ecological site F126XY008OH Tread

Accessed: 05/12/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 126X–Central Allegheny Plateau

These sites generally occur on terrace treads with soils that are well drained or drier. The MAAT > 45 degree F.

This ecosite is found in Alleghany plateau in MLRA 126. Steep slopes are dominant, but level to gently rolling plateau remnants exist. The area is dominantly forest, containing large blocks of state forest, game lands, and national forest. Less than one-tenth of the MLRA consists of urban areas

Classification relationships

This site crosswalks to Landfire biophysical setting (BpS) Northeastern Interior Dry-Mesic Oak Forest

NatureServe's description (2007) for the equivalent ecological system CES202.592 -- Northeastern Interior Dry-Mesic Oak Forest notes this system is found from central New England west through OH and PA and south to VA.

Component Associations

Association Unique ID Association Name

CEGL002059 *Quercus alba* - *Quercus rubra* - *Quercus prinus* - *Acer saccharum* / *Lindera benzoin* Forest

CEGL002067 *Quercus alba* - *Quercus rubra* - *Carya* (alba, ovata) / *Cornus florida* Acidic Forest

CEGL002068 *Quercus alba* - *Quercus rubra* - *Carya ovata* Glaciated Forest

CEGL004793 *Quercus muehlenbergii* - *Quercus* (alba, rubra) - *Carya cordiformis* / *Viburnum prunifolium* Forest

CEGL006057 *Quercus prinus* - *Quercus rubra* / *Hamamelis virginiana* Forest

CEGL006216 *Quercus alba* - *Carya glabra* - *Fraxinus americana* / *Cercis canadensis* / *Muhlenbergia sobolifera* - *Elymus hystrix* Forest

CEGL006241 *Quercus bicolor* / *Vaccinium corymbosum* / *Carex stipata* Forest

CEGL006293 *Pinus strobus* - *Quercus* (rubra, velutina) - *Fagus grandifolia* Forest

CEGL006301 *Quercus rubra* - *Carya* (glabra, ovata) / *Ostrya virginiana* / *Carex lucorum* Forest

CEGL006304 *Liriodendron tulipifera* - *Pinus strobus* - *Tsuga canadensis* - *Quercus* (rubra, alba) / *Polystichum acrostichoides* Forest

CEGL006336 *Quercus* (alba, rubra, velutina) - *Carya* spp. / *Viburnum acerifolium* Forest

CEGL006454 *Quercus* (rubra, velutina, alba) - *Betula lenta* - (*Pinus strobus*) Forest

CEGL006921 *Fagus grandifolia* - *Betula lenta* - *Quercus* (alba, rubra) / *Carpinus caroliniana* Forest

CEGL008475 *Quercus alba* - *Quercus rubra* - *Carya alba* / *Cornus florida* / *Vaccinium stamineum* / *Desmodium nudiflorum* Piedmont Forest

CEGL008514 *Quercus rubra* - *Quercus prinus* - *Carya ovalis* / (*Cercis canadensis*) / *Solidago* (caesia, curtisii) Forest

CEGL008516 *Quercus prinus* - *Quercus rubra* - *Carya ovalis* / *Carex pensylvanica* - (*Calamagrostis porteri*) Forest

CEGL008517 *Quercus rubra* - *Acer saccharum* / *Ostrya virginiana* / *Cardamine concatenata* Forest

CEGL008518 *Quercus rubra* - *Carya* (ovata, ovalis) - *Fraxinus americana* / *Actaea racemosa* - *Hydrophyllum virginianum* Forest

CEGL008528 *Tilia americana* - *Fraxinus americana* / *Acer pensylvanicum* - *Ostrya virginiana* / *Parthenocissus quinquefolia* - *Impatiens pallida* Woodland

See also NatureServe's description (2007):

Allegheny-Cumberland Dry Oak Forest and Woodland CES202.359

Component Associations

Association Unique ID Association Name

CEGL004761 *Pinus echinata* - *Quercus prinus* - *Quercus* (*coccinea*, *velutina*) Forest

CEGL004786 *Quercus prinus* - *Carya* (*alba*, *glabra*, *ovata*) / *Juniperus virginiana* var. *virginiana* Forest

CEGL005023 *Quercus prinus* - *Quercus* (*alba*, *coccinea*, *velutina*) / *Viburnum acerifolium* - (*Kalmia latifolia*) Forest

CEGL006557 *Pinus rigida* - *Quercus coccinea* / *Vaccinium angustifolium* Woodland

CEGL007119 *Pinus virginiana* - *Pinus* (*rigida*, *echinata*) - (*Quercus prinus*) / *Vaccinium pallidum* Forest

CEGL007231 *Quercus alba* - *Quercus velutina* - *Carya* (*ovata*, *alba*, *glabra*) - *Pinus* sp. Forest

CEGL007240 *Quercus alba* - *Quercus rubra* - *Carya ovata* / *Cercis canadensis* - *Juniperus virginiana* var. *virginiana* Forest

CEGL007244 *Quercus falcata* - *Quercus alba* - *Carya alba* / *Oxydendrum arboreum* / *Vaccinium stamineum* Forest

CEGL007247 *Quercus falcata* - *Quercus* (*coccinea*, *stellata*) / *Vaccinium* (*pallidum*, *stamineum*) Forest

CEGL007261 *Quercus prinus* - *Carya* spp. - *Quercus velutina* / *Vaccinium arboreum* / *Iris verna* var. *smalliana* Forest

CEGL007269 *Quercus prinus* - *Quercus rubra* - *Carya* (*ovata*, *glabra*) - *Pinus virginiana* Forest

CEGL007517 *Pinus strobus* - *Quercus alba* - (*Carya alba*) / *Gaylussacia ursina* Forest

CEGL007519 *Pinus strobus* - *Quercus* (*coccinea*, *pinus*) / (*Gaylussacia ursina*, *Vaccinium stamineum*) Forest

CEGL007700 *Quercus prinus* - *Quercus* spp. / *Vaccinium arboreum* - (*Kalmia latifolia*, *Styrax grandifolius*) Forest

CEGL007795 *Quercus alba* - *Carya alba* - (*Quercus velutina*) / *Desmodium nudiflorum* - (*Carex picta*) Forest

CEGL008406 *Quercus stellata* - *Pinus virginiana* / (*Schizachyrium scoparium*, *Piptochaetium avenaceum*) Woodland

CEGL008430 *Quercus alba* - (*Quercus prinus*) / (*Hydrangea quercifolia*) - *Viburnum acerifolium* / *Carex picta* - *Piptochaetium avenaceum* Forest

CEGL008431 *Quercus prinus* - (*Quercus coccinea*) / *Carya pallida* / *Vaccinium arboreum* - *Vaccinium pallidum* Forest

CEGL008443 *Quercus alba* - *Quercus stellata* / *Ostrya virginiana* - *Acer barbatum* / *Chasmanthium sessiliflorum* Forest

CEGL008521 *Quercus alba* - *Quercus* (*coccinea*, *velutina*, *pinus*) / *Gaylussacia baccata* Forest

CEGL008567 *Quercus alba* - *Quercus falcata* / *Vaccinium* (*arboreum*, *hirsutum*, *pallidum*) Forest

Ecological site concept

From Landfire <http://www.landfire.gov/index.php>:

This ecological section comprises part of the Appalachian Plateaus geomorphic province (USDA Forest Service 1995). It is a maturely dissected plateau characterized by high hills, sharp ridges, and narrow valleys. Bedrock is overlain by Quaternary residuum on the ridges and hilltops, colluvium on the slopes, and either or both alluvium and Pleistocene lacustrine materials in the valleys. Udalfs, Udults, and Ochrepts dominate, in combination with mesic soil temperature regime, an udic soil moisture regime, and mixed or illitic mineralogy. Soils formed in parent materials are divided into five groups: residual material, which developed in place by the weathering of underlying bedrock; colluvial material which weathered from bedrock strata transported by water and gravity to the lower slopes; alluvium, lacustrine sediments and outwash deposited by water; and loess deposited by wind. Precipitation averages 35 to 45 in (900 to 1,150 mm); it occurs mainly during summer, winter, and spring. Rain on snow is common during winter and early spring. Summers are dry with low humidity. Temperature averages 52 degrees F (11 degrees C). The growing season is 120 to 180 days. This ecological section is characterized by a relatively high density of streams, with gradients ranging from high, steep headwaters streams to low gradient rivers that flow into the OH River.

NatureServe (2007) provides the following description pertaining to this system type. These oak-dominated forests are one of the matrix forest systems in the northeastern and north-central US. Occurring in dry-mesic settings, they are typically closed-canopy forests, though there may be areas of patchy-canopy woodlands. They cover large expanses at low to mid elevations, where the topography is flat to gently rolling, occasionally steep. Soils are acidic and relatively infertile but not strongly xeric.

Table 1. Dominant plant species

Tree	(1) <i>Quercus Alba</i>
Shrub	(1) <i>Cornus Florida</i>
Herbaceous	Not specified

Physiographic features

Climatic features

Influencing water features

Soil features

Ecological dynamics

Information contained in this section was adapted from several sources. The information presented is representative of very complex vegetation communities. Key indicator plants, animals and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The reference plant community is not necessarily the management goal. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

From Landfire <http://www.landfire.gov/index.php>:

This ecological section comprises part of the Appalachian Plateaus geomorphic province. It is a maturely dissected plateau characterized by high hills, sharp ridges, and narrow valleys. Bedrock is overlain by Quaternary residuum on the ridges and hilltops, colluvium on the slopes, and either or both alluvium and Pleistocene lacustrine materials in the valleys. Udalfs, Udufts, and Ochrepts dominate, in combination with mesic soil temperature regime, an udic soil moisture regime, and mixed or illitic mineralogy. Soils formed in parent materials are divided into five groups: residual material, which developed in place by the weathering of underlying bedrock; colluvial material which weathered from bedrock strata transported by water and gravity to the lower slopes; alluvium, lacustrine sediments and outwash deposited by water; and loess deposited by wind. Precipitation averages 35 to 45in (900 to 1,150mm0150mm); it occurs mainly during summer, winter, and spring. Rain on snow is common during winter and early spring. Summers are dry with low humidity. Temperature averages 52 degrees F (11 degrees C). The growing season is 120 to 180 days. This ecological section is characterized by a relatively high density of streams, with gradients ranging from high, steep headwaters streams to low gradient rivers that flow into the OH River.

NatureServe (2007) provides the following description pertaining to this system type. These oak-dominated forests are one of the matrix forest systems in the northeastern and north-central US. Occurring in dry-mesic settings, they are typically closed-canopy forests, though there may be areas of patchy-canopy woodlands. They cover large expanses at low to mid elevations, where the topography is flat to gently rolling, occasionally steep. Soils are acidic and relatively infertile but not strongly xeric.

Vegetation Description

The vegetation consisted of forests dominated by oaks of dry-mesic conditions, especially white oak (*Quercus alba*) and red oak (*Quercus rubra*), and, on drier sites, chestnut oak (*Quercus prinus*), black oak (*Quercus velutina*), and scarlet oak (*Quercus coccinea*). Scarlet oak is absent at the northern edge of the system range. Along with oaks are varying amounts of hickory (*Carya* spp.), red maple (*Acer rubrum*), and other species such as white pine (*Pinus strobus*) and white ash (*Fraxinus americana*). American chestnut (*Castanea dentata*) was a prominent tree in these forests before chestnut blight eradicated it as a canopy constituent (NatureServe 2007). Common shrubs include mountain laurel (*Kalmia* spp.), greenbriar (*Smilax* spp.), blueberries (*Vaccinium* spp.), and huckleberries (*Gaylussacia* spp.). In the Ridge and Valley region, bear oak is an important shrub component. Herbs, forbs, and ferns are usually sparse to moderate in density. Areas experiencing frequent fire had a greater abundance of

grasses and sedges.

State and transition model




Figure 1. State and Transition Diagram




Figure 2. Legend

State 1
Agricultural Land

Land managed for agricultural production of crops and livestock.

Community 1.1

Pasture (Grasses, Forbes) or Cropland (Grasses, Forbes)

This community phase may contain a wide variety of plants depending on the level of management. In pasture circumstances that are managed tall fescue, bluegrass and white clover will dominate the vegetation canopy. Without management such as prescribed grazing, nutrient management and weed control, less desirable forage species and weeds will invade.

State 2

Post Settlement Forest

This forest vegetation community is the result of removing fire from the landscape, and poor logging techniques (high grading). Following European settlement fire as a management tool was abandoned and mesophication of the vegetation took place.

Community 2.1

Tree/Shrub/Herb

Maple (Red and Sugar)/Striped Maple Forest best captures the nature of this vegetation state. The dominate overstory canopy consist of Maples with minor canopy coverage of Tulip tree. The shrub layer contains Stripped maple, greenbrier and blackberry. The herb layer will contain flowering forbes (asters and goldenrod), ferns (woodferns) and grasses (native fescue, povertygrass).

State 3

Reference Pre-Contact Forest

The reference state can be represented by several communities within the Allegheny-Cumberland Dry Oak Forest and Woodland(Natureserve, 2007). Forest overstory canopies are oak dominated and generally closed canopy.

Community 3.1

Tree/Shrub/Herb

Chestnut Oak-White Oak/Mountain Laurel Forest best captures the nature of this vegetation state. The dominate overstory canopy consist of Oaks with minor canopy coverage of Hickory. The shrub layer contains mountain laurel, greenbrier and ericads. The herb layer will contain flowering forbes (asters and goldenrod), ferns (woodferns) and grasses (native fescue, povertygrass).

Transition T2A

State 1 to 3

The site agricultural management is abandoned and forest regrowth occurs through natural succession or tree planting.

Restoration pathway R3A

State 2 to 1

Remove undesirable species using herbicides, cutting or prescribed fire. Plant desired species if absent from the site.

Transition T3A

State 2 to 2

The site is logged and managed for agricultural land.

Transition T1A

State 3 to 2

The site is logged and managed for agricultural land.

Transition T1B

State 3 to 3

The site is logged and fire is suppressed allowing mesophication to occur.

Additional community tables

Other references

Landfire <http://www.landfire.gov/index.php>

NatureServe. 2007. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA, U.S.A. Data current as of 15 April 2007.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

-
7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
