

Ecological site F134XY013MO Loamy Footslope Forest

Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 134X–Southern Mississippi Valley Loess

The Southern Mississippi Valley Loess (outlined in red on the map; northern portion only) is a relatively narrow strip of the coastal plain bordering the Mississippi River valley, that is blanketed with loess. The northern part of this MLRA, discussed here, is locally referred to as Crowley's Ridge. Elevation ranges from about 300 feet on the footslopes to nearly 600 feet on the highest ridges. Loess caps the summits and upper slopes, and Pliocene-aged sand and gravel deposits of the coastal plain influence soils on lower, steeper slopes.

Classification relationships

Terrestrial Natural Community Type in Missouri (Nelson, 2010):

The reference state for this ecological site is most similar to a Mesic Loess/Glacial Till Forest.

Missouri Department of Conservation Forest and Woodland Communities (Missouri Department of Conservation, 2006):

The reference state for this ecological site is most similar to an Oak-Mixed Hardwood Mesic Forest.

National Vegetation Classification System Vegetation Association (NatureServe, 2010):

The reference state for this ecological site is most similar to an *Acer saccharum* - *Quercus rubra* - *Carya cordiformis* / *Asimina triloba* Forest (CEGL002060).

Geographic relationship to the Missouri Ecological Classification System (Nigh & Schroeder, 2002):
This Ecological Site occurs in the Crowley's Ridge Subsection, and in the Benton Loess Woodland/Forest Hills Land Type Association of the Ozark Outer Border Subsection.

Ecological site concept

Loamy Footslope Forests are within the green areas on the map (Missouri portion only; distributions farther south are currently under review). These sites are locally extensive on footslopes of Crowley's Ridge mostly in Scott county, Missouri, and in adjacent lowland areas in southern Cape Girardeau county, Missouri. Soils are very deep, typically with loamy surfaces and loamy or clayey subsoils. The reference plant community is forest with an overstory dominated by a variety of trees including white oak, sugar maple, northern red oak, bitternut hickory, American elm, walnut and Kentucky coffee tree, an understory dominated by pawpaw, spicebush, leatherwood, and Ohio buckeye, and a rich herbaceous ground flora.

Table 1. Dominant plant species

Tree	(1) <i>Quercus alba</i> (2) <i>Quercus rubra</i>
Shrub	(1) <i>Asimina triloba</i> (2) <i>Lindera benzoin</i>
Herbaceous	(1) <i>Erigenia bulbosa</i> (2) <i>Cardamine concatenata</i>

Physiographic features

This site is on footslopes and stream terraces with slopes of 0 to 8%. The site receives runoff from adjacent upland sites. Some areas flood on rare occasions.

The adjacent figure (adapted from Butler, 1985) shows the typical landscape position of this ecological site, and landscape relationships with other ecological sites. It is within the area labeled "3" on the figure, on footslopes and stream terraces along upland drainageways. Fragipan Upland Woodland sites and Deep Loess Backslope sites are typically upslope.




Figure 2. Typical landscape relationships for this ecological

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Stream terrace
Flooding frequency	None
Ponding frequency	None
Slope	0–8%

Water table depth	61–183 cm
Aspect	Aspect is not a significant factor

Climatic features

Table 3. Representative climatic features

Frost-free period (characteristic range)	164-169 days
Freeze-free period (characteristic range)	198-199 days
Precipitation total (characteristic range)	1,194-1,245 mm
Frost-free period (actual range)	163-170 days
Freeze-free period (actual range)	197-200 days
Precipitation total (actual range)	1,194-1,245 mm
Frost-free period (average)	167 days
Freeze-free period (average)	199 days
Precipitation total (average)	1,219 mm

Climate stations used

- (1) CAPE GIRARDEAU MUNI AP [USW00003935], Chaffee, MO
- (2) JACKSON [USC00234226], Jackson, MO

Influencing water features

Soil features

These soils have no rooting restriction. The soils were formed under forest vegetation, and have thin, light-colored surface horizons. Parent material is alluvium in some areas, loess in some areas, and marine sediments in some areas. Surface horizons are loam or silt loam. Subsurface horizons are loamy or clayey. Some soils are slightly affected by seasonal wetness. Soil series associated with this site include Adler, Farrenburg, Lilbourn, Memphis, and Shadygrove.

Table 4. Representative soil features

Surface texture	(1) Silt loam (2) Fine sandy loam (3) Loam
Family particle size	(1) Loamy
Drainage class	Moderately well drained to well drained
Permeability class	Very slow to moderately slow
Surface fragment cover <=3"	0–5%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	15.24–20.32 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0

Soil reaction (1:1 water) (0-101.6cm)	4.5–7.3
Subsurface fragment volume <=3" (Depth not specified)	0–1%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

State and transition model

Ecosystem states

1. Mesic Bottomland Forest

State 1 submodel, plant communities

1.1. Sugar Maple-Red Oak-Bitternut Hickory/PawPaw Forest

State 1 Mesic Bottomland Forest

Community 1.1 Sugar Maple-Red Oak-Bitternut Hickory/PawPaw Forest

Additional community tables

Table 5. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (M)	Canopy Cover (%)	Diameter (Cm)	Basal Area (Square M/Hectare)
Tree							
sugar maple	ACSA3	<i>Acer saccharum</i>	Native	–	–	–	–
northern red oak	QURU	<i>Quercus rubra</i>	Native	–	–	–	–
bitternut hickory	CACO15	<i>Carya cordiformis</i>	Native	–	–	–	–
Kentucky coffeetree	GYDI	<i>Gymnocladus dioicus</i>	Native	–	–	–	–
black walnut	JUNI	<i>Juglans nigra</i>	Native	–	–	–	–
white oak	QUAL	<i>Quercus alba</i>	Native	–	–	–	–
American elm	ULAM	<i>Ulmus americana</i>	Native	–	–	–	–
slippery elm	ULRU	<i>Ulmus rubra</i>	Native	–	–	–	–

Table 6. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (M)	Canopy Cover (%)
Forb/Herb					
Adam and Eve	APHY	<i>Aplectrum hyemale</i>	Native	—	—
green dragon	ARDR3	<i>Arisaema dracontium</i>	Native	—	—
spring blue eyed Mary	COVE2	<i>Collinsia verna</i>	Native	—	—
lowland bladderfern	CYPR4	<i>Cystopteris protrusa</i>	Native	—	—
common persimmon	DIVI5	<i>Diospyros virginiana</i>	Native	—	—
Shumard's oak	QUSH	<i>Quercus shumardii</i>	—	—	—
American basswood	TIAM	<i>Tilia americana</i>	Native	—	—
zigzag spiderwort	TRSU2	<i>Tradescantia subaspera</i>	Native	—	—
Missouri violet	VIMI3	<i>Viola missouriensis</i>	Native	—	—
striped cream violet	VIST3	<i>Viola striata</i>	Native	—	—
white fawnlily	ERAL9	<i>Erythronium albidum</i>	Native	—	—
common cowparsnip	HEMA80	<i>Heracleum maximum</i>	Native	—	—
zigzag iris	IRBR2	<i>Iris brevicaulis</i>	Native	—	—
butternut	JUCI	<i>Juglans cinerea</i>	Native	—	—
northern spicebush	LIBE3	<i>Lindera benzoin</i>	Native	—	—
Virginia bluebells	MEVI3	<i>Mertensia virginica</i>	Native	—	—
Shrub/Subshrub					
Greek valerian	PORE2	<i>Polemonium reptans</i>	Native	—	—
Tree					
pawpaw	ASTR	<i>Asimina triloba</i>	Native	—	—

Other references

MDC, 2010. Missouri Forest and Woodland Community Profiles. Missouri Department of Conservation, Jefferson City, Missouri.

NatureServe, 2010. Vegetation Associations of Missouri (revised). NatureServe, St. Paul, Minnesota.

Nelson, Paul W. 2010. The Terrestrial Natural Communities of Missouri. Missouri Department of Conservation, Jefferson City, Missouri.

Nigh, Timothy A., & Walter A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City, Missouri.

Contributors

Fred Young

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	

Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-