

# Ecological site F140XY026PA

## Dry Till Uplands

Last updated: 5/20/2020  
Accessed: 04/27/2024

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 140X–Glaciated Allegheny Plateau and Catskill Mountains

This area is primarily in the Southern New York Section of the Appalachian Plateaus Province of the Appalachian Highlands. The top of the dissected plateau in this MLRA is broad and is nearly level to moderately sloping. The narrow valleys have steep walls and smooth floors. The Catskills in the east have steep slopes. Elevation is typically 650 to 1,000 feet on valley floors; 1,650 to 2,000 feet on the plateau surface; and 3,600 feet or more in parts of the Catskills.

The average annual precipitation in most of this area is 30 to 45 inches. Rainfall occurs as high-intensity, convective thunderstorms during the summer, but most of the precipitation in this area occurs as snow. The average annual temperature is 40 to 50 degrees F.

The dominant soil order in this MLRA is Inceptisols. The soils in the area dominantly have a mesic soil temperature regime, an aquic or udic soil moisture regime, and mixed mineralogy. Frigid soils are found within the higher elevations.

This area supports forest vegetation, particularly hardwood species. Beech-birch-maple and elm-ash-red maple are the potential forest types. The extent of oak species increases from east to west, particularly in areas of shallow and dry soils. In some areas conifers, such as white pine, are important. Aspen, hemlock, northern white-cedar, and black ash grow on the wetter soils. In some parts of the area, sugar maple has potential economic significance. Some of the major wildlife species in this area are white-tailed deer, cottontail, turkey, pheasant, and grouse.

### Classification relationships

USDA NRCS:

LRR: R - Northeastern Forage and Forest Region

MLRA 140 - Glaciated Allegheny Plateau and Catskills Mountains

NY Natural Heritage Program Plant Community Classification:

Allegheny Oak Forest

PA Natural Heritage Program Plant Community Classification:

Dry oak - heath forest

### Ecological site concept

Landform/Landscape Position:

The site occurs on benches, hills, hillslopes, ridges and ground moraines. Slopes range from 0 to 35 percent.

Soils:

The soils consists of moderately deep, well drained and somewhat excessively drained, loamy-skeletal soils that formed in glacial till derived mostly from sandstone, siltstone, and shale. Representative soil is Oquaga.

#### Vegetation:

The reference plant community is an oak dominated forest (Dry oak - heath forest, PA Natural Heritage Program). Black oak and scarlet oak are often the most common on the site. Other associated trees include chestnut oak, white oak, red oak, pignut hickory, and eastern white pine. Black huckleberry and lowbush blueberry are common shrubs. In openings following tree falls or other natural disturbances poverty oatgrass and little bluestem may occur with early successional trees such as eastern red cedar and black birch. Invasive exotic plants such as Japanese barberry, multiflora rose, winged euonymus, and shrub honeysuckles can occur in disturbed sites.

**Table 1. Dominant plant species**

Tree	(1) <i>Quercus rubra</i> (2) <i>Carya tomentosa</i>
Shrub	(1) <i>Viburnum recognitum</i> (2) <i>Amelanchier laevis</i>
Herbaceous	(1) <i>Uvularia sessilifolia</i> (2) <i>Maianthemum racemosum ssp. racemosum</i>

### Physiographic features

The site occurs on benches, hills, hillslopes, ridges and ground moraines. Slopes range from 0 to 35 percent.

**Table 2. Representative physiographic features**


Landforms	(1) Bench (2) Hill (3) Hillside (4) Ridge
Slope	0–35%

### Climatic features


Mean annual precipitation is 43 inches and evenly distributed throughout the year. Most of the rainfall occurs as high intensity, convective thunderstorms during the summer. Snowfall is common from late in autumn to early spring. Average frost-free and freeze-free days are 122 and 153, respectively.

**Table 3. Representative climatic features**


Frost-free period (characteristic range)	110-134 days
Freeze-free period (characteristic range)	136-168 days
Precipitation total (characteristic range)	38-49 in
Frost-free period (actual range)	101-136 days
Freeze-free period (actual range)	136-168 days
Precipitation total (actual range)	36-51 in
Frost-free period (average)	122 days
Freeze-free period (average)	154 days
Precipitation total (average)	43 in




**Figure 1. Monthly precipitation range**



**Figure 2. Monthly minimum temperature range**



**Figure 3. Monthly maximum temperature range**



**Figure 4. Monthly average minimum and maximum temperature**




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

### Climate stations used

- (1) BINGHAMTON [USW00004725], Johnson City, NY
- (2) STROUDSBURG [USC00368596], East Stroudsburg, PA
- (3) TOWANDA 1 S [USC00368905], Towanda, PA
- (4) MONTROSE [USC00365915], Montrose, PA
- (5) CORNING [USC00301787], Corning, NY
- (6) ROCK HILL 3 SW [USC00307210], Rock Hill, NY
- (7) CANTON [USC00361212], Canton, PA

### Influencing water features

#### Soil features

The site consists of moderately deep, well drained and somewhat excessively drained, loamy-skeletal soils that formed in glacial till derived mostly from sandstone, siltstone, and shale. Reaction ranges from extremely acid to moderately acid throughout the soil. Representative soil is Oquaga.

Table 4. Representative soil features

Parent material	(1) Till–sandstone and shale (2) Till–sandstone and siltstone
Surface texture	(1) Very channery, channery silt loam (2) Very channery, channery loam (3) Very channery, channery sandy loam
Family particle size	(1) Loamy-skeletal
Drainage class	Well drained to somewhat excessively drained
Depth to restrictive layer	24–35 in
Surface fragment cover >3"	2–40%


Soil reaction (1:1 water) (Depth not specified)	3.6–6
Subsurface fragment volume <=3" (Depth not specified)	20–40%
Subsurface fragment volume >3" (Depth not specified)	5–70%

## Ecological dynamics


The reference plant community is an oak dominated forest (Dry oak - heath forest, PA Natural Heritage Program). Black oak and scarlet oak are often the most common on the site. Other associated trees include chestnut oak, white oak, red oak, pignut hickory, and eastern white pine. Black huckleberry and lowbush blueberry are common shrubs. In openings following tree falls or other natural disturbances poverty oatgrass and little bluestem may occur with early successional trees such as eastern red cedar and black birch. Invasive exotic plants such as Japanese barberry, multiflora rose, winged euonymus, and shrub honeysuckles can occur in disturbed sites.

## State and transition model


### Ecosystem states




### State 1 submodel, plant communities




### State 2 submodel, plant communities



### State 3 submodel, plant communities



### State 4 submodel, plant communities



## State 1

### Reference - Minimally Managed

Reference is an oak-pine forest. Natural disturbances such as wind and ice storms, tree fall, insect damage will create openings for an early successional plant community or young forest. This forest may have at one time been cleared or plowed during colonial times.

**Characteristics and indicators.** Soil may have evidence of an historic plow layer (Ap horizon).

**Resilience management.** Ensure that regenerating trees and shrubs are not heavily browsed by deer that they cannot replace overstory trees. Deer have been shown to have negative effects on forest understories (New York Natural Heritage Program, 2020). Avoid cutting old-growth forests.

### Community 1.1

#### Oak-Pine Forest

The reference community is oak-pine forest/woodland characterized by black oak, chestnut oak, eastern white pine, pitch pine, sassafras, black birch, American beech, witch-hazel, black huckleberry, lowbush blueberry, mountain laurel, maple-leaved viburnum, and Pennsylvania sedge.

### Community 1.2

#### Young Forest

More open canopy, black birch is a common early successional tree.

### Pathway P

#### Community 1.1 to 1.2

Natural disturbances - wind/ice storm, tree fall, and insect damage.

### Conservation practices



### Pathway P

#### Community 1.2 to 1.1

Time (succession).

## State 2

## **Managed Timber**

The state is characterized by active logging. Composition of forest stands will vary based on management objectives.

### **Community 2.1 Managed Timber**

#### **State 3 Minimally Managed Forest with Invasive Species.**

Invasive species such as Japanese barberry, bush honeysuckle, multiflora rose, garlic mustard, and stiltgrass are common in the understory.

### **Community 3.1 Oak-pine forest with invasive species**

#### **State 4 Pasture/Grassland**

Forest has been cleared and grasses and forbs have been introduced for livestock grazing, hay production, and/or wildlife.

### **Community 4.1 Introduced grasses and forbs**

#### **Community 4.2 Woody plant encroachment**

#### **Pathway P Community 4.1 to 4.2**

Abandonment (lack of mowing or fire suppression)

#### **Pathway P Community 4.2 to 4.1**

Mowing, prescribed fire, and/or brush management.

#### **Conservation practices**

Brush Management
------------------

#### **Transition T State 1 to 2**

Timber harvest; logging.

#### **Transition T State 1 to 3**

Introduction of invasive species usually after disturbance.

#### **Transition T State 1 to 4**

Land use conversion.

## **Restoration pathway R**

### **State 2 to 1**

Time (succession). Forest stand improvement, restoration.

## **Transition T**

### **State 2 to 3**

Introduction of invasive species. Lack of timber management.

## **Transition T**

### **State 2 to 4**

Land use conversion

## **Restoration pathway R**

### **State 3 to 1**

Brush management, invasive species management.

## **Transition T**

### **State 3 to 2**

Timber management/harvest, logging.

## **Transition T**

### **State 3 to 4**

Land use conversion.

## **Restoration pathway R**

### **State 4 to 1**

Abandonment, Time (succession), forest restoration.

## **Restoration pathway R**

### **State 4 to 3**

Abandonment, time (sucession) and introduction of invasive species.

## **Additional community tables**

### **Inventory data references**

Site Development and Testing Plan:

Future work to validate the vegetation information in this provisional ecological site description is needed. This will include field activities to collect low and medium intensity sampling and analysis of that data. Field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final approved level document. Reviews of the project plan are to be conducted by the Ecological Site Technical Team.

### **Other references**

Edinger, G.J., Evans, D.J., Gebauer, S., Howard, T.G., Hunt, D.M., and A.M. Olivero, A.M. (eds.). 2014. Ecological Communities of New York State, Second Edition: A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of

Environmental Conservation, Albany, NY.

Zimmerman, E., T. Davis, G. Podniesinski, M. Furedi, J. McPherson, S. Seymour, B. Eichelberger, N. Dewar, J. Wagner, and J. Fike (editors). 2012. Terrestrial and Palustrine Plant Communities of Pennsylvania, 2nd Edition. Pennsylvania Natural Heritage Program, Pennsylvania Department of Conservation and Natural Resources, Harrisburg, Pennsylvania.

## Approval

Nels Barrett, 5/20/2020

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/27/2024
Approved by	Nels Barrett
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 
17. **Perennial plant reproductive capability:**
-