

# Ecological site F142XA019NY

## Acidic Moist Till Frigid

Last updated: 5/22/2020  
Accessed: 04/23/2024

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 142X–St. Lawrence-Champlain Plain

This MLRA is a glaciated area of low relief dominated by broad expanses of nearly level, sandy deltas and shallow lacustrine basins or plains punctuated by low hills of glacial till. Rivers and streams have cut relatively deep but narrow valleys across the plain. Elevation ranges from 80 to 1,000 feet, increasing gradually from the St. Lawrence River southward and from Lake Champlain to the east and west. Local relief generally is less than 30 feet, but glacial till ridges, till plains, and some outwash terraces rise 15 to 80 feet above the adjacent plains.

This area has been glaciated, and a thin mantle of till covers most of the bedrock. Extensive areas of sandy glacial outwash and eolian deposits also occur. Some glacial lake sediments have been deposited above glacial moraines. These deposits are thickest in the valleys and thinnest on the ridges and highlands. During the later stages of the Wisconsin glacial period, seawater entered the Champlain Valley and deposited marine sediments that were later covered by freshwater sediments. The marine deposits are unique to the area.

This area supports hardwoods. The beech-birch-sugar maple forest type is the dominant climax forest type on uplands. Associated with this type are basswood, American elm, maple species, white ash, black cherry, and white pine. The aspen-birch type, earlier in succession, is economically important. Such species as eastern hemlock, red maple, American elm, and spruce are on wet soils.

Some of the major wildlife species in this area are white-tailed deer, red fox, raccoon, beaver, woodchuck, muskrat, cottontail, ruffed grouse, and woodcock.

### LRU notes

Land Resource Unit (LRU): Frigid Soil Temperature Regime

The upper St. Lawrence and Champlain Valleys are characterized with soils in the frigid soil temperature regime (mean annual soil temperature greater than 32°F but less than 46°F and with a difference between mean summer and mean winter soil temperatures greater than 41°F at 20 inches below the surface or at a densic, lithic, or paralithic contact, whichever is shallower).

The Frigid Soil Temperature Regime (STR) will have shorter growing season than the lower St. Lawrence and Champlain Valleys which are characterized with soils in the mesic STR. Species more tolerant of colder year round temperatures would also be evident in the Frigid LRU.

### Classification relationships

NRCS:

Land Resource Region: R - Northeastern Forage and Forest Region

MLRA: 142 - St. Lawrence-Champlain Plain

LRU: A/02 - Frigid Mean Annual Soil Temperature

USFS:

Domain: 200 - Humid Temperate

Division: 210 - Warm Continental

Province: 211 - Northeastern Mixed Forest

Section: 211E - St. Lawrence and Champlain Valley

Subsections: 211Ea - St. Lawrence Glacial Marine Plain

EPA:

Level I: 8 - Eastern Temperate Forests

Level II: 8.1 - Mixed Wood Plains

Level III: 83 - Eastern Great Lakes Lowlands

Level IV: 83d - St. Lawrence Lowlands

83e - Upper St. Lawrence Valley

## Ecological site concept

Landform/Landscape Position:

The site occurs on till plains, hills, ridges, benches. Slopes range from 0 to 15 percent.

Soils:

The site consists of moderately deep and very deep, moderately well drained and somewhat poorly drained coarse-loamy soils that formed in glacial till derived mainly from sandstone and crystalline rock. Soils have an acidic base status throughout. Representative soils are Chazy, Moira, Schroon, and Topknot.

Vegetation:

Based on existing information and known soil/vegetation relationships of the area, the reference plant community is considered to be a Hemlock-Northern Hardwood (Edinger et al. 2014).

Table 1. Dominant plant species

Tree	(1) <i>Tsuga canadensis</i> (2) <i>Acer saccharum</i>
Shrub	(1) <i>Rhododendron maximum</i> (2) <i>Viburnum acerifolium</i>
Herbaceous	(1) <i>Dryopteris intermedia</i> (2) <i>Huperzia lucidula</i>

## Physiographic features

The site occurs on till plains, hills, ridges, benches. Slopes range from 0 to 15 percent.

Table 2. Representative physiographic features


Landforms	(1) Till plain (2) Bench (3) Hill (4) Ridge
Slope	0–15%

## Climatic features


Mean annual precipitation is 37 inches and evenly distributed throughout the year. Most of the rainfall occurs as high intensity, convective thunderstorms during the summer. Snowfall is heavy from late in autumn to early spring. The average temperature in winter is 18°F and in summer it is 66°F. Average frost-free and freeze-free days are 126 and 153, respectively.

**Table 3. Representative climatic features**


Frost-free period (average)	126 days
Freeze-free period (average)	153 days
Precipitation total (average)	940 mm



**Figure 1. Monthly precipitation range**



**Figure 2. Monthly average minimum and maximum temperature**



**Figure 3. Annual precipitation pattern**

### Climate stations used

- (1) OGDENSBURG 4 NE [USC00306164], Ogdensburg, NY
- (2) MASSENA INTL AP [USW00094725], Massena, NY
- (3) LAWRENCEVILLE 3 SW [USC00304647], Nicholville, NY
- (4) MALONE [USC00304996], Malone, NY
- (5) PLATTSBURGH AFB [USC00306659], Plattsburgh, NY

### Influencing water features

### Soil features

The site consists of moderately deep and very deep, moderately well drained and somewhat poorly drained coarse-loamy soils that formed in glacial till derived mainly from sandstone and crystalline rock. Soils have an acidic base status throughout. Representative soils are Chazy, Moira, Schroon, and Topknot.

**Table 4. Representative soil features**


Parent material	(1) Till–sandstone
Family particle size	(1) Coarse-loamy (2) Loamy
Drainage class	Somewhat poorly drained to moderately well drained
Surface fragment cover >3"	2%
Subsurface fragment volume <=3" (Depth not specified)	5–24%
Subsurface fragment volume >3" (Depth not specified)	2–14%

## Ecological dynamics


Based on existing information and known soil/vegetation relationships of the area, the reference plant community is considered to be a Hemlock-Northern Hardwood (Edinger et al. 2014).

## State and transition model


### Ecosystem states




### State 1 submodel, plant communities



### State 2 submodel, plant communities



**State 3 submodel, plant communities**



**State 1**

**Mixed Hardwood Forest - Reference**

Minimally managed hemlock-northern hardwood forest. Natural disturbances primarily from weather events, insect damage, tree fall create pockets of young forest communities (early successional).

**Community 1.1**

**Closed Canopy Forest**

Eastern hemlock, eastern white pine red pine, sugar maple, paper birch are some characteristic trees.

**Community 1.2**

**Young Forest (early successional)**

**Pathway P**

**Community 1.1 to 1.2**

**Conservation practices**

Early Successional Habitat Development/Management

**Pathway P**

**Community 1.2 to 1.1**

**State 2**

**Semi-Natural**

Timber harvest common, invasive species such as bush honeysuckle, oriental bittersweet, Japanese barberry, and multiflora rose may be common in disturbed areas.

**Community 2.1**

**Closed Canopy Forest**

**Community 2.2**

**Young Forest (early successional)**

**State 3**

**Pasture**

Usually in areas of low relief, forest has been converted to pastureland/grassland for either livestock grazing, hay production, or wildlife habitat.

**Community 3.1**

**Introduced Grasses and Forbs**

Grasses and forbs introduces for livestock grazing, hay production, or wildlife habitat.

## Community 3.2 Shrub Encroached

### Pathway P Community 3.1 to 3.2

Lack of mowing, grazing, or fire.

### Pathway P Community 3.2 to 3.1

Brush management and/or fire.

#### Conservation practices

Brush Management
Prescribed Burning

### Transition T State 1 to 2

Human disturbances (roads, timber harvest,) affect species composition and allow the potential for non-native and invasive species to establish and persist.

### Transition T State 1 to 3

#### Conservation practices

Land Clearing
Forage and Biomass Planting

### Restoration pathway R State 2 to 1

#### Conservation practices

Brush Management
Forest Stand Improvement
Forest Management Plan - Written
Forest stand improvement for habitat and soil quality
Herbaceous Weed Control

### Restoration pathway R State 3 to 1

#### Conservation practices

Native Plant Community Restoration and Management
---

## Additional community tables

### Inventory data references

Site Development and Testing Plan:

Future work to validate the vegetation information in this provisional ecological site description is needed. This will include field activities to collect low and medium intensity sampling and analysis of that data. Field reviews should be done by soil scientists and vegetation specialists. A final field review, peer review, quality control, and quality assurance reviews of the ESD will be needed to produce the final approved level document. Reviews of the project plan are to be conducted by the Ecological Site Technical Team.

## Other references

Edinger, G.J., Evans, D.J., Gebauer, S., Howard, T.G., Hunt, D.M., and A.M. Olivero, A.M. (eds.). 2014. Ecological Communities of New York State, Second Edition: A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation, Albany, NY.

Thompson E. H., Sorenson E. R. 2000. Wetland, Woodland, Wildland: A Guide to the Natural Communities of Vermont. Vermont Department of Fish and Wildlife and The Nature Conservancy. University Press of New England, Hanover and London.

## Approval

Nels Barrett, 5/22/2020

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/23/2024
Approved by	Nels Barrett
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if**



**their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

**17. Perennial plant reproductive capability:**

---