

# Ecological site F142XB007VT

## Wet Clayplain Depression

Accessed: 05/20/2024

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 142X–St. Lawrence-Champlain Plain

The St. Lawrence-Champlain Plain MLRA is a glaciated area of low relief dominated by broad expanses of nearly level, sandy deltas and shallow lacustrine basins or plains punctuated by low hills of glacial till. Rivers and streams have cut relatively deep but narrow valleys across the plain. Elevation ranges from 80 to 1,000 feet (25 to 305 meters), increasing gradually from the St. Lawrence River southward and from Lake Champlain to the east and west. Local relief generally is less than 30 feet (10 meters), but glacial till ridges, till plains, and some outwash terraces rise 15 to 80 feet (5 to 25 meters) above the adjacent plains.

### Classification relationships

USDA-NRCS  
LRR: D-Northeastern Forage and Forest Region  
MLRA: 142-St. Lawrence-Champlain Plain.  
LRU: TBD

EPA (Ecoregions)  
Level III: 83-Eastern Great Lakes Lowlands  
Level IV: 83b-Champlain Lowlands

USDA-USFS  
Province: Laurentian Mixed Forest  
Section: 212E--St. Lawrence and Champlain Valley  
Sub-section: 212Ec Champlain Glacial Lake and Marine Plains

### Ecological site concept

Wet clayplain forests occur throughout the MLRA on poorly and very poorly drained glacial lake plains. Much of this Ecological Site has been cleared of forest vegetation and is currently used as hayfield and pasture. This Ecological Site is often adjacent to or embedded within Mesic Clayplain sites. These sites have poor surface drainage; creating shallow rooted trees that are often uprooted by heavy winds. The resultant microtopography of tip-up mounds and depressions is characteristic. Regularly occurring trees include green ash, Eastern hemlock, American elm, red maple, and on some sites, northern white cedar. Invasive shrubs are often common including bush honeysuckles, European buckthorn, common buckthorn, and Japanese barberry. Other shrubs and sapling trees such as arrowwood, American hornbeam, green ash and slippery elm also occur. Herbaceous cover can be variable in composition, reflecting the wetness of individual sites. These include sensitive fern, New York fern, graceful sedge, barren strawberry, white avens, jewelweed, and others.

Table 1. Dominant plant species

Tree	(1) <i>Fraxinus pennsylvanica</i> (2) <i>Quercus bicolor</i>
Shrub	Not specified
Herbaceous	Not specified

## Physiographic features

The site occurs on level to gently sloping glacial lake plains, terraces, and depressions.




Figure 1. Wet Clayplain ES (Covington soils)

Table 2. Representative physiographic features


Landforms	(1) Lake plain (2) Lake terrace (3) Depression
Elevation	90–1,000 ft
Slope	1–5%
Water table depth	9 in
Aspect	Aspect is not a significant factor

## Climatic features


The Lake Champlain Valley has a humid climate with cold winters and warm summers. Average annual precipitation ranges from 750 to 900 mm (30- 35 inches) with peaks in summer (July) and autumn (October). Snowfall averages 1020 to 1520 mm (40 to 60 in) in the Champlain Valley. Average monthly temperature for the warmest month (July) is 28 °C (82.4 °F) and average monthly temperature for the coldest month (January) is -11.9 °C (10.6 °F). The frost free-season is 140 days. The area falls within USDA Hardiness zones 5a (and to a small extent, 5b). The Lake influences the climate of the Valley by moderating temperatures enough with offshore winds to delay the onset of fall, and conversely, the advance of spring. Climate change scenarios for the Northeastern US predict an overall increase in temperature, particularly in the winter and predict a slight increase in annual precipitation.

Table 3. Representative climatic features


Frost-free period (average)	140 days
Freeze-free period (average)	160 days
Precipitation total (average)	39 in



**Figure 2. Monthly precipitation range**



**Figure 3. Monthly average minimum and maximum temperature**



**Figure 4. Annual precipitation pattern**

## Climate stations used

- (1) SOUTH HERO [USC00437607], South Hero, VT
- (2) BURLINGTON INTL AP [USW00014742], South Burlington, VT
- (3) WHITEHALL [USC00309389], Whitehall, NY
- (4) SALISBURY 2 N [USC00437098], Salisbury, VT

## Influencing water features

### Soil features

The site consists of very deep, poorly drained soils that formed in calcareous glacio-lacustrine clays on glacial lake plains. Saturated hydraulic conductivity is low to moderately high in the mineral surface layer and the subsoil, and moderately low to very low in the substratum.

Soil pH increases with soil depth. The surface horizon is considered neutral while the substratum ranges from slightly alkaline to moderately alkaline. With the exception of phosphorous, soil macronutrients such as nitrogen, calcium, potassium, magnesium, and sulfur are increased in moderately alkaline soils, such as the Covington soil

component.



Figure 6. Covington soil, Pedon 2015VT013001

Table 4. Representative soil features

Parent material	(1) Glaciolacustrine deposits–limestone
Surface texture	(1) Clay (2) Silty clay (3) Clay loam
Family particle size	(1) Clayey
Drainage class	Poorly drained
Permeability class	Very slow
Soil depth	72 in
Surface fragment cover ≤3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	4–9 in
Calcium carbonate equivalent (0-40in)	2–8%
Electrical conductivity (0-40in)	0 mmhos/cm
Sodium adsorption ratio (0-40in)	0
Subsurface fragment volume ≤3" (Depth not specified)	0–2%
Subsurface fragment volume >3" (Depth not specified)	0%

## Ecological dynamics

In the reference condition, this site is characterized by mature trees, well-developed strata, and relatively high species diversity owing to the high fertility of the soils. In spite of being wet, because soils are fertile, parts of this site has been drained and lost to agriculture. More research is needed to determine the extent in the nearby eastern Ontario plain and adjacent Canada. Drainage and complete removal of all vegetation and tilling of soils converts the site for agricultural crops, usually corn. Cessation of agriculture allows for the establishment of a sedge – bulrush wet meadow, characterized by sedges, bulrushes, and other graminoids and scattered forbs. Invasion by non-native grasses and forbs, as wells as scattered shrubs also occurs. If unmanaged, early successional native and non-native trees and shrubs become established. These are characterized by wind- or bird-dispersed seeds. If

no active management is employed, the site converts gradually to early successional forest comprising young trees, shrubs, and herbs in a patchy mosaic, and is highly variable among locations. If invasive species are present - and in most cases they are - only manual removal, followed by close monitoring for newly established invasive plants, and judicious planting of native trees and shrubs can convert the site to a semblance of the reference state.

## State and transition model




Figure 7. 142XY002 - Wet Clayplain

Code	Activity, Process
1.1A	Minor disturbances, tree removal or windthrow, soil disturbance, plant replacements
T1A, T3A, T4A	Removal of all vegetation, soil disturbance and planting of crops
3.2A	Manual removal of invasives, shrub removal
1.2A, R2A	Natural replacement or succession
T1B, T3B	Invaded replacement or succession
R3A	Cessation of agriculture, natural replacement or succession
R4A	Manual removal of invasives and planting of native species

Figure 8. 142XY002 - Weta Clayplain

## State 1 Reference

This state is characterized by a mature forest characterized by green ash and swamp white oak. Associated trees may include slippery elm, basswood, eastern hemlock, red maple, shagbark hickory, and white pine. Northern white cedar is sometimes present. The tree canopy is generally closed to partially open. Shrub and herb layers are of variable cover; American hornbeam and southern arrowwood form an open shrub layer. Common herbaceous species include sensitive fern, jack-in-the-pulpit, jewelweed, slender sedge, enchanter's nightshade, and parasol whitetop aster.

## Community 1.1

### Green ash – Swamp white oak / Graceful sedge Wet Clayplain Forest (*Fraxinus pennsylvanica* – *Quercus*



Figure 9. Wet clayplain, Robinsons Point 2015VT013001

This wet, deciduous forest occurs in wet, fertile clay and silt plains. The closed canopy is typically green ash (*Fraxinus pennsylvanica*) and swamp white oak (*Quercus bicolor*). The shrub layer is quite variable and dominated by musclewood aka American hornbeam (*Carpinus caroliniana*). Several wetland plants are found in the understory including the graceful sedge (*Carex gracillima*), Inland sedge (*Carex interior*), and jewelweed (*Impatiens capensis*).

**Forest overstory.** The closed canopy is Green ash, and Swamp white oak admixed with slippery elm and red maple. Occasional trees include white pine and eastern hemlock. Northern white cedar is sometimes present.

**Forest understory.** Forest understory includes a variously developed shrub layer of musclewood aka American hornbeam.

The herb layer is rich with ferns, sedges, and herbs - characteristically the graceful sedge, interior sedge, and jewelweed.

Table 5. Soil surface cover

Tree basal cover	0%
Shrub/vine/liana basal cover	0-2%
Grass/grasslike basal cover	1-5%
Forb basal cover	2-5%
Non-vascular plants	1-20%
Biological crusts	0%
Litter	64-88%
Surface fragments >0.25" and ≤3"	0%
Surface fragments >3"	0-1%
Bedrock	0%
Water	0-1%
Bare ground	0%

Table 6. Woody ground cover

Downed wood, fine-small (<0.40" diameter; 1-hour fuels)	4-30%
Downed wood, fine-medium (0.40-0.99" diameter; 10-hour fuels)	20-30%
Downed wood, fine-large (1.00-2.99" diameter; 100-hour fuels)	13-22%

Downed wood, coarse-small (3.00-8.99" diameter; 1,000-hour fuels)	1-6%
Downed wood, coarse-large (>9.00" diameter; 10,000-hour fuels)	—
Tree snags** (hard***)	—
Tree snags** (soft***)	—
Tree snag count** (hard***)	
Tree snag count** (hard***)	

\* **Decomposition Classes:** N - no or little integration with the soil surface; I - partial to nearly full integration with the soil surface.

\*\* >10.16cm diameter at 1.3716m above ground and >1.8288m height--if less diameter OR height use applicable down wood type; for pinyon and juniper, use 0.3048m above ground.

\*\*\* Hard - tree is dead with most or all of bark intact; Soft - most of bark has sloughed off.

**Table 7. Canopy structure (% cover)**

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	—	—
>0.5 <= 1	—	—	—	—
>1 <= 2	—	—	—	—
>2 <= 4.5	—	1-1%	5-20%	20-65%
>4.5 <= 13	—	20-50%	—	—
>13 <= 40	5-10%	—	—	—
>40 <= 80	65-70%	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

## Community 1.2

### White pine - green ash - swamp white oak Wet Clayplain Forest

This phase is characterized by patches of younger forest, often characterized by greater cover of white pine or other early successional trees such as trembling aspen or big-tooth aspen. Species composition of shrub and herbaceous layers is similar to Community Phase 1.1.

**Forest overstory.** The closed canopy is Green ash, and Swamp white oak admixed with slippery elm and red maple. Occasional trees include white pine and eastern hemlock. Northern white cedar is sometimes present.

**Forest understory.** Forest understory includes a variously developed shrub layer of musclewood aka American hornbeam.

The herb layer is rich with ferns, sedges, and herbs - characteristically the graceful sedge, interior sedge, and jewelweed.

## Pathway 1.1A

### Community 1.1 to 1.2

This transition results from minor disturbances such as windthrows, ice storms, soil disturbance, light tree harvest, and plant replacements.

### Conservation practices

Brush Management
Forage Harvest Management
Tree/Shrub Establishment
Forest Management Plan - Applied

## Pathway 1.2A

### Community 1.2 to 1.1

Natural replacement (succession) or native planting

#### Conservation practices

Tree/Shrub Establishment
--------------------------

## State 2

### Converted Agricultural State

This state bears no resemblance to the reference state or earlier successional states. All native vegetation is removed and soils are severely disturbed by tilling. Corn is a typical agricultural crop on this converted site.

## Community 2.1

### Corn

Corn in a typical agricultural crop on this converted site.

## State 3

### Fallow / Abandoned State

This states exists after agricultural abandonment. Dominated by graminoids (sedges, rushes, and grasses), this state can persist indefinitely if actively managed.

## Community 3.1

### Sedge - bulrush wet meadow

This phase is characterized by a treeless wet meadow that may occur after the immediate cessation of agriculture. Dominated by Bulrushes (*Scirpus* spp.) and other graminoids. Kiviat and Stevens (2001) describe a Wet Clay Meadow type associated with the Hudson – Vergennes complex of the Hudson River Estuary Corridor in eastern New York. The vegetation is characterized by goldenrods, false foxglove, fox sedge, Bush's sedge, as well as the shrubs gray dogwood, silky dogwood, northern arrowwood, meadowsweet, and alder. Additional sedges include *Carex lasiocarpa*, *C. granularis*, *C. annectens*, *C. cristatella*, *C. hirsutella*, and *C. scoparia*.

Table 8. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	—	—
>0.5 <= 1	—	—	—	—
>1 <= 2	—	—	—	—
>2 <= 4.5	—	—	—	—
>4.5 <= 13	—	—	100-100%	—
>13 <= 40	—	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

## State 4

### Invaded State

The tree canopy is very open, characterized by young trees, and an absence of late-successional species such as



swamp white oak, hickories, basswood, or eastern hemlock that otherwise characterize the reference state.

## Community 4.1

### Green ash – Slippery elm / Common buckthorn Wet Clayplain Forest

This phase is highly variable in composition, but is characterized by a high cover of exotic invasive species. Common plants include common buckthorn, Morrow's honeysuckle, Tartarian honeysuckle, European cranberry-bush, common barberry, and creeping Jenny. Early successional native trees are often high in cover, including big-tooth aspen, trembling aspen, and eastern red-cedar.

**Forest overstory.** This ecological site can be highly variable in composition. The open canopy is green ash and slippery elm occur sporadically in the open canopy with smaller early successional trees like bigtooth aspen trembling aspen and red cedar in more abundance.

**Forest understory.** Forest understory includes a variously developed shrub layer of mainly invasive plants, including the common buckthorn, honeysuckles.

Table 9. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	—	—
>0.5 <= 1	—	—	—	—
>1 <= 2	—	—	—	—
>2 <= 4.5	—	—	—	60-60%
>4.5 <= 13	—	70-70%	—	—
>13 <= 40	—	—	—	—
>40 <= 80	20-40%	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

## Transition T1A

### State 1 to 2

Removal of all vegetation, soil disturbance, and planting of crops

## Transition T1A

### State 1 to 2

Complete removal of all vegetation, soil disturbance, and planting of crops.

## Transition T1B

### State 1 to 4

Disturbances leading to the establishment of invaded plants

## Restoration pathway R2A

### State 2 to 3

Restoration occurs following cessation of active agriculture, when natural succession or active planting allows establishment of early successional trees shrubs, and herbs.

## Conservation practices

Tree/Shrub Establishment
--------------------------

**Restoration pathway R3A**  
**State 3 to 1**

Natural replacement (succession), selective plantings

**Conservation practices**

Tree/Shrub Establishment
Native Plant Community Restoration and Management

**Transition T3A**  
**State 3 to 2**

Removal of all vegetation, soil disturbance, planting of crops

**Transition T3B**  
**State 3 to 4**

This transition occurs when minor disturbance allow for the establishment of invasive plants.

**Restoration pathway R4A**  
**State 4 to 1**

Active management to removeinvasive plants and plant native plants.

**Conservation practices**

Brush Management
Native Plant Community Restoration and Management
Invasive Plant Species Control

**Transition T4A**  
**State 4 to 2**

Complete removal of all vegetation , soil disturbance, and planting of crops.

**Additional community tables**

Table 10. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)	Diameter (In)	Basal Area (Square Ft/Acre)
<b>Tree</b>							
red maple	ACRU	<i>Acer rubrum</i>	Native	33–66	10–25	–	–
green ash	FRPE	<i>Fraxinus pennsylvanica</i>	Native	33–82	10–25	–	–
arborvitae	THOC2	<i>Thuja occidentalis</i>	Native	33–66	0–25	–	–
eastern hemlock	TSCA	<i>Tsuga canadensis</i>	Native	33–66	5–10	–	–
American elm	ULAM	<i>Ulmus americana</i>	Native	33–66	5–10	–	–
American basswood	TIAM	<i>Tilia americana</i>	Native	33–66	2–5	–	–
eastern white pine	PIST	<i>Pinus strobus</i>	Native	33–66	2–5	–	–
American beech	FAGR	<i>Fagus grandifolia</i>	Native	33–66	2–5	–	–
hophornbeam	OSVI	<i>Ostrya virginiana</i>	Native	–	2–5	–	–
shagbark hickory	CAOV2	<i>Carya ovata</i>	Native	33–66	2–5	–	–
slippery elm	ULRU	<i>Ulmus rubra</i>	Native	33–66	2–5	–	–

**Table 11. Community 1.1 forest understory composition**

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
<b>Grass/grass-like (Graminoids)</b>					
inland sedge	CAIN11	<i>Carex interior</i>	Native	0.3–1.6	2–5
graceful sedge	CAGR2	<i>Carex gracillima</i>	Native	0.3–1.6	2–5
New England sedge	CANO4	<i>Carex novae-angliae</i>	Native	0.3–1.6	1–2
whitegrass	LEVI2	<i>Leersia virginica</i>	Native	–	–
<b>Forb/Herb</b>					
jewelweed	IMCA	<i>Impatiens capensis</i>	Native	0.3–1.6	2–5
eastern poison ivy	TORA2	<i>Toxicodendron radicans</i>	Native	0.3–1.6	1–2
white avens	GECA7	<i>Geum canadense</i>	Native	0.3–1.6	1–2
Appalachian barren strawberry	WAFR	<i>Waldsteinia fragarioides</i>	Native	0.3–1.6	1–2
waxflower shinleaf	PYEL	<i>Pyrola elliptica</i>	Native	0.3–1.6	1–2
broadleaf enchanter's nightshade	CILUC	<i>Circaea lutetiana ssp. canadensis</i>	Native	0.3–1.6	0.5–1
Jack in the pulpit	ARTR	<i>Arisaema triphyllum</i>	Native	0.3–1.6	0.5–1
parasol whitetop	DOUM2	<i>Doellingeria umbellata</i>	Native	0.3–1.6	0.5–1
littleleaf buttercup	RAAB	<i>Ranunculus abortivus</i>	Native	0.3–1.6	0.5–1
calico aster	SYLA4	<i>Symphyotrichum lateriflorum</i>	Native	0.3–1.6	0.5–1
white trillium	TRGR4	<i>Trillium grandiflorum</i>	Native	0.3–1.6	0.5–1
alpine violet	VILA10	<i>Viola labradorica</i>	Native	0.3–1.6	0.5–1
blisterwort	RARE2	<i>Ranunculus recurvatus</i>	Native	–	–
wild sarsaparilla	ARNU2	<i>Aralia nudicaulis</i>	Native	–	–
wrinkleleaf goldenrod	SORU2	<i>Solidago rugosa</i>	Native	–	–
<b>Fern/fern ally</b>					
sensitive fern	ONSE	<i>Onoclea sensibilis</i>	Native	0.3–1.6	1–2
eastern marsh fern	THPA	<i>Thelypteris palustris</i>	Native	0.3–1.6	1–2
bulblet bladderfern	CYBU3	<i>Cystopteris bulbifera</i>	Native	0.3–1.6	1–2
New York fern	THNO	<i>Thelypteris noveboracensis</i>	Native	0.3–1.6	1–2
common ladyfern	ATFI	<i>Athyrium filix-femina</i>	Native	0.3–1.6	1–2
spinulose woodfern	DRCA11	<i>Dryopteris carthusiana</i>	Native	0.3–1.6	0.5–1
mountain woodfern	DRCA3	<i>Dryopteris campyloptera</i>	Native	–	–
<b>Shrub/Subshrub</b>					
Morrow's honeysuckle	LOMO2	<i>Lonicera morrowii</i>	Introduced	6.5–16	5–10
Tatarian honeysuckle	LOTA	<i>Lonicera tatarica</i>	Introduced	0.3–1.6	2–5
southern arrowwood	VIDE	<i>Viburnum dentatum</i>	Native	0.3–1.6	1–2
European cranberrybush	VIOP	<i>Viburnum opulus</i>	Native	0.3–1.6	0.5–1
common winterberry	ILVE	<i>Ilex verticillata</i>	Native	–	–
<b>Tree</b>					
American hornbeam	CACA18	<i>Carpinus caroliniana</i>	Native	6.5–16	10–25
green ash	FRPE	<i>Fraxinus pennsylvanica</i>	Native	6.5–16	2–5
eastern hemlock	TSCA	<i>Tsuga canadensis</i>	Native	6.5–16	1–2

Table 12. Community 2.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
<b>Grass/grass-like (Graminoids)</b>					
corn	ZEMA	<i>Zea mays</i>	Introduced	—	—

Table 13. Community 3.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
<b>Grass/grass-like (Graminoids)</b>					
rufous bulrush	SCPE4	<i>Scirpus pendulus</i>	Native	1.6–5	—
blackgirdle bulrush	SCAT4	<i>Scirpus atrocinctus</i>	Native	1.6–5	—
green bulrush	SCAT2	<i>Scirpus atrovirens</i>	Native	1.6–5	—
red clover	TRPR2	<i>Trifolium pratense</i>	Introduced	—	—
timothy	PHPR3	<i>Phleum pratense</i>	Introduced	1.6–5	—

Table 14. Community 4.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)	Diameter (In)	Basal Area (Square Ft/Acre)
<b>Tree</b>							
green ash	FRPE	<i>Fraxinus pennsylvanica</i>	Native	33–50	20–40	—	—
slippery elm	ULRU	<i>Ulmus rubra</i>	Native	33–50	20–40	—	—

Table 15. Community 4.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
<b>Grass/grass-like (Graminoids)</b>					
graceful sedge	CAGR2	<i>Carex gracillima</i>	Native	1.6–3.3	—
fox sedge	CAVU2	<i>Carex vulpinoidea</i>	Native	1.6–3.3	—
<b>Forb/Herb</b>					
Canada goldenrod	SOCA6	<i>Solidago canadensis</i>	Native	1.6–3.3	—
Canada goldenrod	SOAL6	<i>Solidago altissima</i>	Native	1.6–3.3	—
candle anemone	ANCY	<i>Anemone cylindrica</i>	—	1.6–3.3	—
broadleaf enchanter's nightshade	CILUC	<i>Circaea lutetiana ssp. canadensis</i>	Native	1.6–3.3	—
<b>Shrub/Subshrub</b>					
Morrow's honeysuckle	LOMO2	<i>Lonicera morrowii</i>	Introduced	3.3–9.8	—
European cranberrybush	VIOP	<i>Viburnum opulus</i>	Introduced	3.3–9.8	—
common buckthorn	RHCA3	<i>Rhamnus cathartica</i>	Introduced	3.3–9.8	—
<b>Tree</b>					
sugar maple	ACSA3	<i>Acer saccharum</i>	Native	3.3–9.8	—
arborvitae	THOC2	<i>Thuja occidentalis</i>	Native	3.3–9.8	—
eastern redcedar	JUVI	<i>Juniperus virginiana</i>	Native	3.3–9.8	—
<b>Vine/Liana</b>					
Virginia creeper	PAQU2	<i>Parthenocissus quinquefolia</i>	Native	1.6–3.3	—

## Animal community

Source: Thompson and Sorenson 2000) Mammals characteristic of this site include gray squirrel, eastern chipmunk, raccoon, and white-tailed deer. Beaver also frequent this site where it is associated with standing water.

Common birds include wood thrush, eastern wood peewee, ovenbird, northern oriole, and downy woodpecker. Amphibians associated with this site include blue-spotted salamander, American toad, wood frog, and green treefrog. Where vernal pools occur within this site, invertebrates such as caddis flies, predaceous diving beetles, and horsehair worms can be found. Animals associated with Kiviat and Stevens (2001) Wet Clay Meadow: included amphibians American toad and spotted salamander. Associated birds included Virginia rail, American woodcock, alder flycatcher, sedge wren, vesper sparrow, Henslow's sparrow, and orchard oriole. The Baltimore checkerspot is a potentially occurring butterfly.

## **Hydrological functions**

The poorly drained soils cause surface water to pool in low-lying areas, but no water features are associated with this site.

## **Recreational uses**

This site has limited use for recreation other than occasional hiking.

## **Wood products**

This site has limited use for wood products except as a by-product of clearing for agriculture.

## **Other products**

This site has deep, fertile soils that are heavily used for agriculture.

## **Other information**

Conservation Status of clayplain forest is high.

## **Inventory data references**

The data contained in this document is derived from the analysis of field inventories (relevé plots and reconnaissance notes collected by NatureServe with the assistance of MLRA Soil Survey Office 12-TOL, 12-PAS, 12-AMH and the VT NRCS State Office. Five high intensity plots were conducted for the reference state. This information was supplemented by plot data from Vermont Natural Heritage Program.

## **Other references**

### **REFERENCES**

Kiviat, E. and G. Stevens. 2001. Biodiversity assessment manual for the Hudson River Estuary Corridor. Hudsonia Ltd. New York State Department of Environmental Conservation, Albany, New York. 508 pp.

Thompson, E. H., and E. R. Sorenson. 2000. Wetland, woodland, wildland: A guide to the natural communities of Vermont. The Nature Conservancy and the Vermont Department of Fish and Wildlife. University Press of New England, Hanover, NH. 456 pp.

Edinger, G.J., D.J. Evans, S. Gebauer, T.G. Howard, D.M. Hunt, and A.M. Olivero (editors). 2014. Ecological Communities of New York State. Second Edition. A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation, Albany, NY.

## **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be

known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/20/2024
Approved by	Nels Barrett
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**  

---
2. **Presence of water flow patterns:**  

---
3. **Number and height of erosional pedestals or terracettes:**  

---
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**  

---
5. **Number of gullies and erosion associated with gullies:**  

---
6. **Extent of wind scoured, blowouts and/or depositional areas:**  

---
7. **Amount of litter movement (describe size and distance expected to travel):**  

---
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**  

---
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**  

---
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**  

---
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be**

mistaken for compaction on this site):

---

12. **Functional/Structural Groups** (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence** (include which functional groups are expected to show mortality or decadence):
- 

14. **Average percent litter cover (%) and depth ( in):**
- 

15. **Expected annual annual-production** (this is TOTAL above-ground annual-production, not just forage annual-production):
- 

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-