

Ecological site F143XY902ME

Alpine Ridge Krummholz

Last updated: 10/07/2024
Accessed: 12/04/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 143X–Northeastern Mountains

MLRA 143, known as the Northeastern Mountains, covers approximately 23 million acres of mountains, hills, and valleys in northern Maine, New Hampshire, Vermont, New York, and Massachusetts. The area is sparsely populated, with less than five percent of the land area developed for agriculture, residential, and urban development. About 90 percent of the area is forested, most of which is actively managed for timber. Elevations are mostly between 1,000 to 4,000 feet, with a few isolated peaks more than 5,000 feet above sea level. The present day mountains are but remnants of a much larger ancient range that has been eroding for approximately 500 million years. Bedrock consists of mostly very old metamorphic rock (gneiss, schist, slate, marble, quartzite, etc.) with younger intrusions of igneous rock (e.g. granite and granodiorite) from the Triassic and Cretaceous periods. MLRA 143 differs somewhat geologically from its neighboring MLRAs (142, 144A, 144B, 145, and 146), which have greater amounts of nutrient-rich sedimentary rock. Compared to MLRA 143, they are all lower in elevation, with longer growing seasons large areas that were once submerged by the ocean following glaciation.

The characteristic landforms and soils of northern New England were derived from the massive continental ice sheet that engulfed the region during North America's most recent glaciation. Mighty glaciers, embedded with sediment and rock fragments, scoured bedrock and compacted mineral beds in a steady march south and east toward the Atlantic Ocean. The softer sedimentary rocks were pulverized into fine silts and clays under the immense weight of ice a mile thick, while the more resistant igneous and metamorphic rocks were sculpted into steep mountains and hills or plucked and dragged along the base of the glacier. With a warming climate, the ice retreated northward, depositing a thin layer of unsorted glacial till sediment atop the newly-exposed bedrock and compacted mineral beds. Deeper mounds of unsorted till formed small hills, kames, moraines and drumlins. Enormous chunks of ice detached as the glacier retreated, melting slowly in place and forming many kettle lakes and basins where water and fine sediments collect. Raging torrents of glacial meltwater dissected much of the barren landscape, entraining coarse and fine sediments, carving river valleys, and leaving well-sorted deposits of mostly sand and gravel along the watercourse. By 10,000 years ago the ice sheet had fully receded from MLRA 143. Silty floodplains developed along perennial rivers, many of which occupy the same channels that once gushed with sediment-rich glacial meltwater. Over time, wet basins accumulated fine sediment, some dried out, and still others became acidified by organic matter inputs from colonizing vegetation.

In terms of climate, MLRA 143 is distinguished from neighboring MLRAs by a shorter growing season and the occurrence of cryic soil temperature regimes at high elevations. The majority of MLRA 143 averages 32 to 44 inches of precipitation annually with a five to six month growing season and frigid winter temperatures. However, the higher elevations may receive up to double the annual precipitation of the lower elevations, and have a three to four month growing season with extremely cold winters. As the northernmost MLRA in the region with the coldest temperatures and shortest growing season, the Northeastern Mountains have less overall tree diversity, fewer pine and oak trees, and more abundant spruce and fir trees than neighboring MLRAs.

Classification relationships

This site occurs in Ecological Site Group 9 (Alpine/Subalpine) of MLRA 143 (The Northeastern Mountains), in the Northeastern Forage and Forest Region (Land Resource Region R).

The Northeastern Forage and Forest LRR includes all of Maine, New Hampshire, Vermont, Rhode Island, and Connecticut, as well as large portions of Massachusetts, New York, New Jersey, Pennsylvania, and Ohio. Its southern boundary marks the extent of the Wisconsin ice sheet, which engulfed the entire LRR as recently as 10,000 to 15,000 years ago. Erosional and depositional processes associated with glaciation created many of the topographic patterns that distinguish MLRAs within the Northeastern region. Harder granitic and metamorphic bedrock to the north were more resistant to glacial erosion, resulting in the relatively nutrient poor mountains of MLRA 143; whereas nutrient-rich sedimentary bedrock of MLRAs 139, 140, and 146 resulted in relatively flat, fertile landscapes ideal for cultivation. Other areas were depressed below sea-level by the sheer mass of the glacier, resulting in pockets of marine sediments which distinguish MLRAs 142, 144A, 144B, and 145.

Precipitation is sufficient to support productive forestland throughout the Northeastern region. Still, a latitudinal temperature gradient from mesic to frigid soil temperatures results in a general transition from central hardwoods and pine in the southern MLRAs to northern hardwoods and spruce-fir forests farther north (no true boreal forests exist in the region). Elevations are generally low throughout the Northeastern region, with the exception of MLRA 143 which has many high mountain ecosystems with cryic temperature regimes and alpine vegetation above the tree line.

Ecological site concept

This site is found on high mountains and ridges, just below the tree line where balsam fir trees exhibit a stunted growth form. Elevations range from 3500 to 4500 feet. Soils may be organic or mineral, shallow to deep, with or without rock fragments in the profile. All soils are well- to excessively-drained, and pH is typically between 3.4 and 6.0. These broad ranges in soil properties may be explained by balsam fir krummholz being controlled by topographic and climatic factors more than soil properties. Further study is needed to clarify the soil concept of this site.

This site is dominated by stunted balsam fir trees. On the harshest sites, the branches of fir trees may only grow on the leeward side of the tree trunk in relation to the direction of prevailing winds.

This site is resistant to disturbance and exhibits limited community dynamics.

Associated sites

F143XY901ME	<p>Alpine Ridge Meadow The Alpine Ridge Meadow site occurs upslope of the Alpine Ridge Krummholz site--that is, it occurs above tree line rather than below tree-line.</p>
-------------	---

Similar sites

F143XY901ME	<p>Alpine Ridge Meadow The Alpine Ridge Meadow site is dominated by herbs and shrubs rather than stunted balsam fire trees. However, at its lower extent it may include small patches of stunted balsam fir within a matrix of its characteristic shrub and herb species.</p>
-------------	--

Table 1. Dominant plant species

Tree	(1) <i>abies balsamea</i>
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site is found on high mountains and ridges, just below the tree line where balsam fir trees exhibit a stunted growth form. Elevations range from 3500 to 4500 feet. Despite high rainfall, these landscape positions shed water

rapidly, are exposed to high winds and heavy snows, and represent harsh growing conditions for vegetation.

Table 2. Representative physiographic features

Landforms	(1) Mountain (2) Ridge (3) Mountain slope
Flooding frequency	None
Ponding frequency	None
Elevation	1,067–1,372 m
Slope	5–100%
Aspect	Aspect is not a significant factor

Climatic features

The climate of this site is much cooler than the typical climate of MLRA 143, with very cold snowy winters, high winds, cool rainy summers, and a very short growing season. Precipitation is fairly constant from month to month and averages about 83 inches annually. Growing degree days ranges from 88-124 days from June to September. Soil temperature regime is cryic on this site.

Table 3. Representative climatic features

Frost-free period (average)	88 days
Freeze-free period (average)	124 days
Precipitation total (average)	2,108 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly average minimum and maximum temperature




Figure 3. Annual precipitation pattern

Climate stations used

- (1) MT MANSFIELD [USC00435416], Underhill, VT
- (2) PINKHAM NOTCH [USC00276818], Sargents Purchase, NH
- (3) MT WASHINGTON [USW00014755], Sargents Purchase, NH

Influencing water features

Due to its landscape position, this site is not typically influenced by streams or wetlands.

Soil features

This site occurs on a broad range soils. Soils may be organic or mineral, shallow to deep, with or without rock fragments in the profile. All soils are well- to excessively-drained, and pH is typically between 3.4 and 6.0. These broad ranges in soil properties may be explained by balsam fir krummholz being controlled by topographic and climatic factors more than soil properties. Further study is needed to clarify the soil concept of this site.

Table 4. Representative soil features

Parent material	(1) Lodgment till–mica schist (2) Organic material–granite (3) Colluvium–phyllite
Surface texture	(1) Fine sandy loam (2) Silt loam (3) Loamy sand
Family particle size	(1) Loamy
Drainage class	Well drained to excessively drained
Soil depth	25–152 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0–35%
Available water capacity (0-101.6cm)	2.54–38.1 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	3.4–6

Subsurface fragment volume <=3" (Depth not specified)	0–30%
Subsurface fragment volume >3" (Depth not specified)	0–22%

Ecological dynamics

This site is dominated by stunted balsam fir trees. On the harshest sites, the branches of fir trees may only grow on the leeward side of the tree trunk in relation to the direction of prevailing winds.

This site is resistant to disturbance and exhibits limited community dynamics.

State and transition model

F143XY902ME – Alpine Ridge Krummholz

1. Reference State/Current Potential

1.1 Stunted balsam fir dominates

Figure 5. STM

Other references

Barton, A. M., A. S. White, and C. V. Cogbill. 2012. *The Changing Nature of the Maine Woods*. University Press of New England, Lebanon, NH.

Gawler, S. and A. Cutko. 2010. *Natural Landscapes of Maine: A Guide to Natural Communities and Ecosystems*. Maine Natural Areas Program, Maine Department of Conservation, Augusta, Maine.

Johanson, J. K., Butler, N. R. and C. Bickford. 2016. *Classifying Northern New England Landscapes for Improved Conservation*. *Rangelands* 38:6.

Sperduto, D. and B. Kimball. 2011. *The Nature of New Hampshire: Natural Communities of the Granite State*. The Nature Conservancy and The New Hampshire Heritage Bureau. University Press of New England, Lebanon, NH.

Thompson, E. H. and E. R. Sorenson. 2000. Wetland, Woodland, Wildland: A Guide to the Natural Communities of Vermont. The Nature Conservancy and the Vermont Department of Fish and Wildlife. University Press of New England, Hanover, NH.

USDA Natural Resources Conservation Service. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Contributors

Jamin Johanson

Approval

Greg Schmidt, 10/07/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	12/04/2024
Approved by	Greg Schmidt
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

-
7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
