

## Ecological site F149BY004NY Wet Lake Plain

Last updated: 5/23/2020  
Accessed: 05/03/2024

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 149B—Long Island-Cape Cod Coastal Lowland

149B—Long Island-Cape Cod Coastal Lowland

MLRA 149B is in the Embayed Section of the Coastal Plain Province of the Atlantic Plain. It is part of the partially submerged coastal plain of New England. It is mostly an area of nearly level to rolling plains, but it has some steeper hills (glacial moraines). Ridges border the lower plains. Elevation generally ranges from sea level to 80 feet (0 to 25 meters), but it is as much as 330 feet (100 meters) in a few areas. Local relief is mainly 3 to 30 feet (1 to 10 meters), but it is as much as 65 feet (20 meters) in the more hilly areas. The Peconic and Carmans Rivers are on the eastern end of Long Island. The parts of this area in Massachusetts and Rhode Island have no major rivers.

This entire area is made up of deep, unconsolidated glacial outwash deposits of sand and gravel. A thin mantle of glacial till covers most of the surface. Some moraines form ridges and higher hills in this area of generally low relief. Sand dunes and tidal marshes are extensive along the coastline.

### Classification relationships

USDA-NRCS (USDA, 2006):

Land Resource Region (LRR): S—Northern Atlantic Slope Diversified Farming Region

Major Land Resource Area (MLRA): 149B—Long Island-Cape Cod Coastal Lowland

USDA-FS (Cleland et al., 2007):

Province: 221 Eastern Broadleaf Forest Province

Section: 221A Lower New England

Subsection: 221Ab Cape Cod Coastal Lowland and Islands

Subsection: 221An Long Island Coastal Lowland and Moraine

### Ecological site concept

This site consists of very deep, poorly drained soils that formed in silty estuarine or glacio-lacustrine deposits on glacial lake plains. Representative soil is Raynham (coastal).

The reference community in this site is “red maple-swamp white oak” (Edinger et al. 2014). The dominant tree is swamp white oak with red maple. Other trees include pin oak, black gum, and green ash. Characteristic shrubs include winterberry holly, northern arrowwood. Groundcover is commonly ferns: cinnamon fern, royal fern, marsh fern; and sedges: Gray’s sedge, fringed sedge, hop sedge.

### Associated sites

F149BY003MA	Well Drained Lake Plain
-------------	-------------------------

**Table 1. Dominant plant species**

Tree	(1) <i>Acer rubrum</i> (2) <i>Quercus bicolor</i>
Shrub	(1) <i>Vaccinium corymbosum</i>
Herbaceous	Not specified

## Physiographic features

The site occurs on nearly level to gently sloping soils in low-lying positions on glaciolacustrine and marine terraces.

**Table 2. Representative physiographic features**


Landforms	(1) Lake plain (2) Marine terrace (3) Gap
Flooding frequency	None
Elevation	0–152 m
Slope	0–3%
Water table depth	10–30 cm
Aspect	Aspect is not a significant factor

## Climatic features

Mean annual precipitation is 50 inches and is usually uniformly distributed throughout the year. Frost free and freeze free days average 177 and 207, respectively.

**Table 3. Representative climatic features**

Frost-free period (average)	177 days
Freeze-free period (average)	207 days
Precipitation total (average)	1,270 mm



**Figure 1. Monthly precipitation range**




Figure 2. Monthly average minimum and maximum temperature




Figure 3. Annual precipitation pattern

### Climate stations used

- (1) BRIDGEHAMPTON [USC00300889], Sag Harbor, NY
- (2) GREENPORT PWR HOUSE [USC00303464], Greenport, NY
- (3) MINEOLA [USC00305377], Mineola, NY
- (4) RIVERHEAD RSCH FM [USC00307134], Riverhead, NY

### Influencing water features

These sites may be considered depressional or "perched" with seeps, hence maybe considered more minerally-enriched.

### Soil features

This site consists of very deep, poorly drained soils that formed in silty estuarine or glacio-lacustrine deposits on glacial lake plains. Representative soil is Raynham soils mapped within MLRA 149B.

Table 4. Representative soil features

Surface texture	(1) Sand (2) Silt loam (3) Very fine sandy loam
Drainage class	Poorly drained
Permeability class	Very slow
Soil depth	183 cm
Available water capacity (0-101.6cm)	20.32 cm
Soil reaction (1:1 water) (0-101.6cm)	4.5–7.8

## Ecological dynamics

[Caveat: The vegetation information contained in this section and is only provisional, based on concepts, not yet validated with field work.\*]


The vegetation groupings described in this section are based on the terrestrial ecological system classification and vegetation associations developed by NatureServe (Comer 2003). Terrestrial ecological systems are specifically defined as a group of plant community types (associations) that tend to co-occur within landscapes with similar ecological processes, substrates, and/or environmental gradients. They are intended to provide a classification unit that is readily mappable, often from terrain and remote imagery, and readily identifiable by conservation and resource managers in the field. A given system will typically manifest itself in a landscape at intermediate geographic scales of tens-to-thousands of hectares and will persist for 50 or more years. A vegetation association is a plant community that is much more specific to a given soil, geology, landform, climate, hydrology, and disturbance history. It is the basic unit for vegetation classification and recognized by the US National Vegetation Classification (US FDGC 2008; USNVC 2017). Each association will be named by the diagnostic and often dominant species that occupy the different height strata (tree, shrub, and herb). Within the NatureServe Explorer database, ecological systems are numbered by a community Ecological System Code (CES) and individual vegetation associations are assigned an identification number called a Community Element Global Code (CEGL).

[\*Caveat] The information presented is representative of very complex vegetation communities. Key indicator plants and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and geography. The reference plant community is not necessarily the management goal. The drafts of species lists are merely representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

The Wet Lake Plains ecological site occurs on poorly-drained flats and depressions in protected areas along the coast and is characterized mainly by coastal forests, and variants such as wet “sunken” forests. These wet coastal forests are highly varied and range from deciduous swamps dominated by red maple to sunken forests dominated by pin oak and swamp white oak to wet evergreen swamps dominated by Atlantic white cedar (*Chamaecyparis thyoides*). These wet lake plains in coastal settings are not well-described but may in part, include vegetation within the Northern Atlantic Coastal Plain Basin Swamp and Wet Hardwood Forest system (CES203.520) and in part by the Northern Atlantic Coastal Plain Basin Peat Swamp system (CES203.522) (NatureServe 2017).

## State and transition model

## 149BY004 –Wet lake Plains



Transition	Drivers/practices
T1-2	disturbance, invasive plant establishment
T1-3, T2-3	cutting, land clearing, plant establishment
R2-1, R3-1	herbaceous weed treatment, plant removal, plant establishment, successional management
T3-2	abandonment, disturbance, invasive plant establishment
CP1.1-1.2	disturbance
CP1.2-1.1	succession
CP2.1-2.2	Invasive plant establishment
CP2.1-2.1	Invasive plant removal

### State 1

#### Reference State (Wet Lake Plains)

Predominately coastal forests • Northeastern Pin Oak - Swamp White Oak Wet Forest, (Pin Oak - (Swamp White Oak) - Red Maple / Highbush Blueberry / Cinnamon Fern Wet Forest), [*Quercus palustris* - (*Quercus bicolor*) - *Acer rubrum* / *Vaccinium corymbosum* / *Osmunda cinnamomea* Wet Forest], - CEGL006240 • Coastal Plain Atlantic White-cedar Swamp Forest, (Atlantic White-cedar / Inkberry - Swamp Azalea Swamp Forest). [*Chamaecyparis thyoides* / *Ilex glabra* - *Rhododendron viscosum* Swamp Forest], - CEGL006188 Other plant communities may include: • Lower New England Red Maple Swamp Forest, (Red Maple / Swamp Azalea - Coastal Sweet-peppercorn Swamp Forest), [*Acer rubrum* / *Rhododendron viscosum* - *Clethra alnifolia* Swamp Forest], - CEGL006156 • Red Maple / Upright Sedge Wet Woodland, (Red Maple / Upright Sedge - Sensitive Fern Wet Woodland), [*Acer rubrum* / *Carex stricta* - *Onoclea sensibilis* Wet Woodland], - CEGL006119 (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]).

#### Community 1.1

##### Pin Oak - (Swamp White Oak) - Red Maple / Highbush Blueberry / Cinnamon Fern Wet Forest

Northeastern Pin Oak - Swamp White Oak Wet Forest, (Pin Oak - (Swamp White Oak) - Red Maple / Highbush Blueberry / Cinnamon Fern Wet Forest), [*Quercus palustris* - (*Quercus bicolor*) - *Acer rubrum* / *Vaccinium corymbosum* / *Osmunda cinnamomea* Wet Forest], - CEGL006240 This association is a woody, deciduous, seasonally flooded forest with a closed to partially open canopy, often located on sandy loams or clayey soils of

glacial lakeplains in basin areas that are seasonally wet (winter and early spring) with a shallow, perched water table, that may become drier in late summer and early fall creating a hummock-and-hollow microtopography. There is generally some layer that impedes drainage. The canopy is codominated by pin oak (*Quercus palustris*) and/or swamp white oak (*Quercus bicolor*) and red maple (*Acer rubrum*). Common associates include blackgum (*Nyssa sylvatica*) and occasionally bitternut hickory (*Carya cordiformis*) and sometimes eastern hemlock (*Tsuga canadensis*). The shrub layer may be sparse or dense and contains winterberry (*Ilex verticillata*), highbush blueberry (*Vaccinium corymbosum*), smooth arrowwood (*Viburnum dentatum*), buttonbush (*Cephalanthus occidentalis*), and sheep laurel (*Kalmia angustifolia*). The sparse herb layer may include cinnamon fern (*Osmunda cinnamomea*), marshfern (*Thelypteris palustris*), Massachusetts fern (*Thelypteris simulata*), sensitive fern (*Onoclea sensibilis*), royal fern (*Osmunda regalis*), common woosedge (*Scirpus cyperinus*), Frank's sedge (*Carex frankii*), fowl mannagrass (*Glyceria striata*), and fringed sedge (*Carex crinita*). (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]). Cross-referenced plant community concepts (typically by political state): Black Gum - Pin Oak - Swamp White Oak Perched Swamp (Swain 2016) [MA] Red maple - swamp white oak swamp (Edinger et al. 2014) [NY] Mixed Hardwood Swamp Forest (Greller 1977) [NY]

## **Community 1.2**

### **Atlantic White-cedar / Inkberry - Swamp Azalea Swamp Forest**

Coastal Plain Atlantic White-cedar Swamp Forest, (Atlantic White-cedar / Inkberry - Swamp Azalea Swamp Forest). [*Chamaecyparis thyoides* / *Ilex glabra* - *Rhododendron viscosum* Swamp Forest], - C EGL006188 This Atlantic white-cedar swamp occurs in poorly-drained depressions or basins generally overlaying sand and gravel deposits. The canopy is dominated by Atlantic white cedar (*Chamaecyparis thyoides*) or codominated with red maple (*Acer rubrum*). Less frequent canopy associates include pitch pine (*Pinus rigida*), blackgum (*Nyssa sylvatica*), and sometimes eastern white pine (*Pinus strobus*) and occasionally eastern hemlock (*Tsuga canadensis*). The shrub layer is very dense and diverse with sweet pepperbush (*Clethra alnifolia*), inkberry (*Ilex glabra*), northern bayberry (*Morella pensylvanica*), blue huckleberry (*Gaylussacia frondosa*), swamp doghobble (*Leucothoe racemosa*), swamp azalea (*Rhododendron viscosum*), smooth winterberry (*Ilex laevigata*), winterberry (*Ilex verticillata*), black chockberry (*Aronia melanocarpa*), and highbush blueberry (*Vaccinium corymbosum*). The herbaceous layer tends to be sparse or patchy and limited to sunny openings with cinnamon fern (*Osmunda cinnamomea*), marshfern (*Thelypteris palustris*), Virginia chainfern (*Woodwardia virginica*), netted chainfern (*Woodwardia areolata*), Massachusetts fern (*Thelypteris simulata*), eastern teaberry (*Gaultheria procumbens*), sundew (*Drosera* spp.), purple pitcherplant (*Sarracenia purpurea*), rose pogonia (*Pogonia ophioglossoides*), partridgeberry (*Mitchella repens*), Walter's sedge (*Carex striata*), Collins' sedge (*Carex collinsii*), prickly bogsedge (*Carex atlantica*), and poison ivy (*Toxicodendron radicans*). The nonvascular layer can several species of peatmoss (*Sphagnum*), commonly *Sphagnum fallax*, *Sphagnum flavicomans*, *Sphagnum magellanicum*, *Sphagnum pulchrum*, *Sphagnum recurvum*, and *Sphagnum palustre*. (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]). Coastal Atlantic White Cedar Swamp (Swain 2016) [MA] Coastal plain Atlantic white cedar swamp (Edinger et al. 2014) [NY] Coastal Plain Atlantic White Cedar Swamp (Sneddon et al. 2010) [Cape Cod National Seashore]

## **Pathway CP1.1-2.1**

### **Community 1.1 to 1.2**

Disturbance

## **Pathway CP1.2-1.1**

### **Community 1.2 to 1.1**

Abandonment, Sucession

## **State 2**

### **Semi-natural State**

Vegetation on lands somewhat conditioned by land use, e.g., managed native plant communities or invasive plant communities.

## **Community 2.1**

### **Managed forest/woodlands/meadows**

## Community 2.2 Invasive Plants

Black locust (*Robinia pseudoacacia*), Norway Maple (*Acer platanoides*), tree-of-heaven (*Ailanthus altissima*), winged burningbush (*Euonymus alatus*), Japanese knotweed (*Polygonum cuspidatum*), and oriental bittersweet (*Celastrus orbiculatus*), multiflora rose (*Rosa multiflora*), wineberry (*Rubus phoenicolasius*), garlic mustard (*Alliaria petiolata*), major celindine (*Chelidonium majus*), ground ivy (*Glechoma hederacea*), and European lily-of-the-valley (*Convallaria majalis*), can characterize the herb layer, which may also include native plants.

## Pathway CP2.1-2.2 Community 2.1 to 2.2

Disturbance, Invasive species establishment

## Pathway CP2.2-2.1 Community 2.2 to 2.1

Invasive spp. Control, Forest mgmt.

## State 3 Cultural State

Landscapes heavily conditioned by land use, e.g., Plantations / Gardens / Cultivated / Pasture.

## Community 3.1 Plantations / Gardens / Cultivated / Pasture

## Transition T1-2 State 1 to 2

altered by human- induced Disturbance or Management

### Conservation practices

Tree/Shrub Establishment
Forest Land Management
Forest stand improvement for habitat and soil quality

## Transition T1-3 State 1 to 3

Disturbance, clearing, cutting

### Conservation practices

Brush Management
Land Clearing
Herbaceous Weed Control

## Restoration pathway R2-1 State 2 to 1

Plant removals, plantings, Invasive plant control, successional mgmt., forestry practices Restoration & Mgmt, Forest Stand Improvement, Early Successional Habitat Development, Upland Wildlife Mgmt, Invasive spp. Control, Plant establishment

### Conservation practices

Brush Management
Tree/Shrub Establishment
Early Successional Habitat Development/Management
Forest Stand Improvement
Restoration and Management of Natural Ecosystems
Native Plant Community Restoration and Management
Forest Land Management
Invasive Plant Species Control

### Transition T2-3

#### State 2 to 3

Land clearing, cutting

### Conservation practices

Brush Management
Land Clearing
Herbaceous Weed Control

### Restoration pathway R3-1

#### State 3 to 1

Plant removals, plantings, Invasive plant control, successional mgmt., forestry practices Restoration & Mgmt, Forest Stand Improvement, Early Successional Habitat Development, Upland Wildlife Mgmt, Invasive spp. Control, Plant establishment

### Conservation practices

Restoration and Management of Natural Ecosystems
Native Plant Community Restoration and Management

### Transition T3-2

#### State 3 to 2

Abandonment. Plant establishment, Forest mgmt.

### Conservation practices

Tree/Shrub Establishment
Forest Stand Improvement
Forest Land Management

### Additional community tables

#### Inventory data references

Site Development and Testing Plan

Future work is needed, as described in a project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation



ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

## References

- Comer, P., D. Faber-Langendoen, R. Evans, S. Grawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schultz, K. Snow, and J. Teague. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia..
- FGDC [Federal Geographic Data Committee]. 2008. National Vegetation Classification Standard, Version 2. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC..
- Greller, A. 1977. A Classification of Mature Forests on Long Island, New York . Bulletin of the Torrey Botanical Club 104:376–382.
- Parshall, T., D.R. Foster, E. Faison, D. MacDonald, and Hansen. 2003. Long-term history of vegetation and fire in pitch pine–oak forests on cape cod, Massachusetts.. Ecology 84:736–748.
- Swain, P.C. 2016. Classification of the natural communities of Massachusetts, Natural Heritage & Endangered Species Program, Massachusetts Division of Fisheries and Wildlife,.
- USNVC [United States National Vegetation Classification]. 2017 (Date accessed). United States National Vegetation Classification Database V2.01. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC.

## Contributors

Nels Barrett, Ph.D.  
Michael Margo

## Approval

Nels Barrett, 5/23/2020

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/21/2020
Approved by	Nels Barrett
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**  

---
2. **Presence of water flow patterns:**  

---
3. **Number and height of erosional pedestals or terracettes:**  

---
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**  

---
5. **Number of gullies and erosion associated with gullies:**  

---
6. **Extent of wind scoured, blowouts and/or depositional areas:**  

---
7. **Amount of litter movement (describe size and distance expected to travel):**  

---
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**  

---
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**  

---
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**  

---
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**  

---
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---