

# Ecological site F149BY008MA

## Very Wet Outwash

Last updated: 5/23/2020  
Accessed: 04/24/2024

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 149B—Long Island-Cape Cod Coastal Lowland

149B—Long Island-Cape Cod Coastal Lowland

This area is in the Embayed Section of the Coastal Plain Province of the Atlantic Plain. It is part of the partially submerged coastal plain of New England. It is mostly an area of nearly level to rolling plains, but it has some steeper hills (glacial moraines). Ridges border the lower plains. Elevation generally ranges from sea level to 80 feet (0 to 25 meters), but it is as much as 330 feet (100 meters) in a few areas. Local relief is mainly 3 to 30 feet (1 to 10 meters), but it is as much as 65 feet (20 meters) in the more hilly areas. The Peconic and Carmans Rivers are on the eastern end of Long Island. The parts of this area in Massachusetts and Rhode Island have no major rivers.

This entire area is made up of deep, unconsolidated glacial outwash deposits of sand and gravel. A thin mantle of glacial till covers most of the surface. Some moraines form ridges and higher hills in this area of generally low relief. Sand dunes and tidal marshes are extensive along the coastline.

### Classification relationships

USDA-NRCS (USDA, 2006):

Land Resource Region (LRR): S—Northern Atlantic Slope Diversified Farming Region

Major Land Resource Area (MLRA): 149B—Long Island-Cape Cod Coastal Lowland

USDA-FS (Cleland et al., 2007):

Province: 221 Eastern Broadleaf Forest Province

Section: 221A Lower New England

Subsection: 221Ab Cape Cod Coastal Lowland and Islands

Subsection: 221An Long Island Coastal Lowland and Moraine

### Ecological site concept

This site consists of very deep, very poorly drained soils that formed in sandy and gravelly glaciofluvial deposits. They are on toeslopes in wet depressions on outwash plains and deltas, along drainage ways, and adjacent to swamps and bogs. Representative soils are Rainberry.

Many of the plant communities are considered maritime if influenced by salt spray from coastal storms or coastal if only influenced by the coastal climate. The representative plant communities are dominated by more open plant communities e.g., “shrub swamps” (Swain and Kearsley 2001); and “shallow/deep emergent marshes” (Swain and Kearsley 2001). Eutrophication can convert these site to common reed marshes.

### Associated sites

F149BY007NY	<b>Moist Outwash</b> Moist Outwash
-------------	---------------------------------------

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	(1) <i>Vaccinium corymbosum</i> (2) <i>Cephalanthus occidentalis</i>
Herbaceous	Not specified

## Physiographic features

The site occurs in wet depressions, drainageways, and are adjacent to swamps and bogs in toeslope positions on glaciofluvial and glaciolacustrine landforms. Slopes range from 0 to 3 percent.

**Table 2. Representative physiographic features**


Landforms	(1) Outwash plain (2) Depression (3) Drainageway
Ponding frequency	Occasional
Slope	0–3%

## Climatic features

Mean annual precipitation is 49 inches and is usually uniformly distributed throughout the year. Frost free and freeze free days average 180 and 201, respectively.

**Table 3. Representative climatic features**

Frost-free period (average)	180 days
Freeze-free period (average)	201 days
Precipitation total (average)	48 in



**Figure 1. Monthly precipitation range**




Figure 2. Monthly average minimum and maximum temperature




Figure 3. Annual precipitation pattern

### Climate stations used

- (1) HYANNIS [USC00193821], Hyannis, MA
- (2) EAST WAREHAM [USC00192451], East Wareham, MA

### Influencing water features

wet depressions

### Wetland description

Palustrine scrubshrub deciduous seasonally saturated (PSSIE) (Cowardin

### Soil features

The site consists of very deep, very poorly drained soils that formed in sandy and gravelly glaciofluvial deposits.

Table 4. Representative soil features

Parent material	(1) Glaciofluvial deposits–granite
Drainage class	Very poorly drained
Permeability class	Moderate
Soil depth	72 in
Soil reaction (1:1 water) (0-40in)	3.5–6

### Ecological dynamics

[Caveat: The vegetation information contained in this section and is only provisional, based on concepts, not yet validated with field work.\*]


The vegetation groupings described in this section are based on the terrestrial ecological system classification and vegetation associations developed by NatureServe (Comer 2003). Terrestrial ecological systems are specifically defined as a group of plant community types (associations) that tend to co-occur within landscapes with similar ecological processes, substrates, and/or environmental gradients. They are intended to provide a classification unit that is readily mappable, often from terrain and remote imagery, and readily identifiable by conservation and resource managers in the field. A given system will typically manifest itself in a landscape at intermediate geographic scales of tens-to-thousands of hectares and will persist for 50 or more years. A vegetation association is a plant community that is much more specific to a given soil, geology, landform, climate, hydrology, and disturbance history. It is the basic unit for vegetation classification and recognized by the US National Vegetation Classification (US FDGC 2008; USNVC 2017). Each association will be named by the diagnostic and often dominant species that occupy the different height strata (tree, shrub, and herb). Within the NatureServe Explorer database, ecological systems are numbered by a community Ecological System Code (CES) and individual vegetation associations are assigned an identification number called a Community Element Global Code (CEGL).

[\*Caveat] The information presented is representative of very complex vegetation communities. Key indicator plants and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and geography. The reference plant community is not necessarily the management goal. The drafts of species lists are merely representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

The Very Wet Outwash ecological site, set in wet basins with saturated hydrologic conditions, is characterized by wetland plant communities with coastal affinities from Long Island, New York, north to Cape Cod, Massachusetts. These plant communities are mostly within the Northern Atlantic Coastal Plain Basin Peat Swamp system (CES203.522). The prevailing ecological processes are related to wetness, but also coastal influences, such as a coastal climate and storms, and if within close proximity to the coast, maritime effects of wind exposure, salt spray, and sand movement. These wetlands are quite varied and may be wet thickets dominated by highbush blueberry (*Vaccinium corymbosum*), swamp azalea (*Rhododendron viscosum*), and sweet pepperbush (*Clethra alnifolia*) or buttonbush (*Cephalanthus occidentalis*) and swamp loosestrife (*Decodon verticillatus*) or marshes of cattails (*Typha* spp.). Other swamp dominants may be red maple (*Acer rubrum*) and/or Atlantic white cedar (*Chamaecyparis thyoides*). Threats include drainage and conversion to developed land, invasives such as common reed (*Phragmites australis*). (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]).

## **State and transition model**

## 149BY008 – Very Wet Outwash



Transition	Drivers/practices
T1-2	disturbance, invasive plant establishment
T1-3, T2-3	cutting, land clearing, plant establishment
R2-1, R3-1	herbaceous weed treatment, plant removal, plant establishment, successional management
T3-2	abandonment, disturbance, invasive plant establishment
CP1.1-1.2	disturbance
CP1.2-1.1	succession
CP2.1-2.2	Invasive plant establishment
CP2.1-2.1	Invasive plant removal

### State 1

#### Reference State (Very Wet Outwash)

The predominant plant communities of the Very Wet Outwash ecological site's Reference State (minimally-managed) include: • Blueberry Wetland Thicket, (Highbush Blueberry - Swamp Azalea - Coastal Sweet-pepperbush Acidic Peatland), [*Vaccinium corymbosum* - *Rhododendron viscosum* - *Clethra alnifolia* Acidic Peatland], - CEGLO06371 • Northeastern Buttonbush Shrub Swamp, (Common Buttonbush - Swamp-loosestrife Shrub Swamp), [*Cephalanthus occidentalis* - *Decodon verticillatus* Shrub Swamp], - CEGLO06069 Other common plant communities in more enriched conditions may include: • Eastern Cattail Marsh, Cattail (Narrowleaf Cattail, Broadleaf Cattail) - (Bulrush species) Eastern Marsh), [*Typha* (*angustifolia*, *latifolia*) - (*Schoenoplectus* spp.) Eastern Marsh], - CEGLO06153 • Lower New England Red Maple Swamp Forest, (Red Maple / Swamp Azalea - Coastal Sweet-pepperbush Swamp Forest), [*Acer rubrum* / *Rhododendron viscosum* - *Clethra alnifolia* Swamp Forest], - CEGLO06156 • Coastal Plain Atlantic White Cedar Swamp Forest, (Coastal Plain Atlantic White Cedar Swamp Forest), [*Chamaecyparis thyoides* / *Ilex glabra* - *Rhododendron viscosum* Swamp Forest], - CEGLO06188 (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]).

### Community 1.1

#### Highbush Blueberry - Swamp Azalea - Coastal Sweet-pepperbush Thicket

Blueberry Wetland Thicket, (Highbush Blueberry - Swamp Azalea - Coastal Sweet-pepperbush Thicket),

[*Vaccinium corymbosum* - *Rhododendron viscosum* - *Clethra alnifolia* Thicket], - CEG006371 Typically, these associations occurs in small open basins, closed sandplain basins, and seasonally flooded zones within larger wetlands. It may also form as a result of beaver activity. This vegetation can occur on the margins of Coastal Plain ponds. This community is influenced by a strongly fluctuating water table with flooded conditions in spring and early summer followed by a drop in the water table below soil surface usually by late summer. Dominant shrubs include highbush blueberry (*Vaccinium corymbosum*), winterberry (*Ilex verticillata*), and swamp azalea (*Rhododendron viscosum*). Scattered red maple (*Acer rubrum*) are occasional. Maleberry (*Lyonia ligustrina*) and buttonbush (*Cephalanthus occidentalis*) are characteristic. Associated shrub species may include sweet pepperbush (*Clethra alnifolia*), rosy meadowsweet (*Spiraea tomentosa*), leatherleaf (*Chamaedaphne calyculata*), inkberry (*Ilex glabra*), swamp doghobble (*Leucothoe racemosa*), swamp loostriffe (*Decodon verticillatus*), sheep laurel (*Kalmia angustifolia*), smooth alder (*Alnus serrulata*), sweetgale (*Myrica gale*), and chokeberry (*Aronia* spp.). Herbaceous composition is variable; some of the more typical species include cinnamon fern (*Osmunda cinnamomea*), royal fern (*Osmunda regalis*), marsh fern (*Thelypteris palustris*), sensitive fern (*Onoclea sensibilis*), water arum (*Calla palustris*), northern water horehound (*Lycopus uniflorus*), Virginia marsh St. John'swort (*Triadenum virginicum*), fowl mannagrass (*Glyceria striata*), rice cutgrass (*Leersia oryzoides*), threeway sedge (*Dulichium arundinaceum*), common sofrush (*Juncus effusus*), and Virginia chainfern (*Woodwardia virginica*). A layer of peatmoss (*Sphagnum* spp. may occur, including: *Sphagnum fimbriatum*, *Sphagnum rubellum*, *Sphagnum magellanicum*, *Sphagnum fallax*, and *Sphagnum viridum*). (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]). Cross-referenced plant community concepts (typically by political state): Shrub Swamp (Swain 2016) [MA] Shrub Swamp (Edinger et al. 2014) [NY] Blueberry wetland thicket (Sneddon et al. 2010) [Cape Cod National Seashore]

## Community 1.2

### Common Buttonbush - Swamp-loosestrife Shrub Swamp

Northeastern Buttonbush Shrub Swamp, (Common Buttonbush - Swamp-loosestrife Shrub Swamp), [*Cephalanthus occidentalis* - *Decodon verticillatus* Shrub Swamp], - CEG006069 These swamps can experience prolonged or semipermanent flooding for much of the growing season, with water tables receding below the soil surface only during drought or very late in the growing season. They occur in a variety of environmental settings. Buttonbush (*Cephalanthus occidentalis*) is dominant and often monotypic. Occasional associates depend on the environmental setting and most often occur in drier areas. They include highbush blueberry (*Vaccinium corymbosum*), swamp azalea (*Rhododendron viscosum*), and possibly red maple (*Acer rubrum*) and dogwoods (*Cornus* spp). closer to upland borders or swamp loostriffe (*Decodon verticillatus*), leatherleaf (*Chamaedaphne calyculata*), and white meadowsweet (*Spiraea alba* var. *latifolia*) in more stagnant basins. Herbaceous species tend to be sparse but can include Canada mannagrass (*Glyceria canadensis*), threeway sedge (*Dulichium arundinaceum*), tussock sedge (*Carex stricta*), common woolsedge (*Scirpus cyperinus*), marshfern (*Thelypteris palustris*), rice cutgrass (*Leersia oryzoides*), European sweetflag (*Acorus calamus*), European waterplantain (*Alisma plantago-aquatica*), smartweeds/knitweeds (*Polygonum* spp.), (bur-reeds (*Sparganium* spp)., and possibly floating or submerged aquatic species such as duckweed (*Lemna minor*), pondweed (*Potamogeton* spp.), and variagated yello pondlily (*Nuphar variegata*). Bryophytes, if present, cling to shrub bases and include warnstofia moss (*Warnstorfia fluitans*), dreplanoclatus moss (*Drepanocladus aduncus*), or peatmoss (*Sphagnum fallax*). In disturbed areas, these wetland may be invaded by purple loosestrife (*Lythrum salicaria*). (Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]). Cross-referenced plant community concepts (typically by political state): Shrub Swamp (Swain 2016) [MA] Shrub Swamp (Edinger et al. 2014) [NY] Northeastern Buttonbush Shrub Swamp (Sneddon et al. 2010) [Cape Cod National Seashore]

## Pathway CP1.1-1.2

### Community 1.1 to 1.2

disturbance, greater fire frequency, coastal proximity

## Pathway CP1.2-1.1

### Community 1.2 to 1.1

succession, lessor fire frequency

## State 2

### Semi-natural State

Vegetation on lands somewhat conditioned by land use, e.g., managed native plant communities or invasive plant communities.

**Community 2.1**  
**Managed wet woodland/thicket**

**Community 2.2**  
**Invasive Plant Community**

purple loosestrife (*Lythrum salicaria*), common reed (*Phragmites australis*)

**Pathway CP2.1-2.1**  
**Community 2.1 to 2.2**

Invasive Plant establishment

**Pathway CP2.2-2.1**  
**Community 2.2 to 2.1**

Invasive Plant Management

**Conservation practices**

Invasive Plant Species Control
--------------------------------

**State 3**  
**Cultural State**

Landscapes heavily conditioned by land use, e.g., Plantations/gardens/cultivation(?)

**Community 3.1**  
**Cultivated/Cranberry Bogs(?)**

cranberry (*Vaccinium macrocarpon*)

**Transition T1-2**  
**State 1 to 2**

disturbance, invasive plant establishment

**Conservation practices**

Forest Land Management
------------------------

**Transition T1-3**  
**State 1 to 3**

cutting, land clearing, plant establishment

**Conservation practices**

Brush Management
Land Clearing

**Restoration pathway R2-1**

## State 2 to 1

herbaceous weed treatment, plant removal, plant establishment, successional management

### Conservation practices

Brush Management
Restoration and Management of Natural Ecosystems
Native Plant Community Restoration and Management
Forest Land Management
Invasive Plant Species Control
Monitoring and Evaluation

## Transition T2-3

### State 2 to 3

cutting, land clearing, plant establishment

### Conservation practices

Land Clearing
Invasive Plant Species Control
Herbaceous Weed Control

## Restoration pathway R3-1

### State 3 to 1

herbaceous weed treatment, plant removal, plant establishment, successional management

### Conservation practices

Brush Management
Restoration and Management of Natural Ecosystems
Native Plant Community Restoration and Management
Invasive Plant Species Control
Monitoring and Evaluation
Herbaceous Weed Control

## Transition T3-2

### State 3 to 2

disturbance, invasive plant establishment

## Additional community tables

### References

Cleland, D.T., J.A. Freeouf, J.E. Keys, G.J. Nowacki, C. Carpenter, and W.H. McNab. 2007. Ecological Subregions: Sections and Subsections of the Coterminous United States. USDA Forest Service, General Technical Report WO-76. Washington, DC. 1–92.

Comer, P., D. Faber-Langendoen, R. Evans, S. Grawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K.

Schultz, K. Snow, and J. Teague. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia..

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States..

NatureServe. 2018 (Date accessed). NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available <http://explorer.natureserve.org>.  
<http://explorer.natureserve.org>.

Swain, P.C. 2016. Classification of the natural communities of Massachusetts, Natural Heritage & Endangered Species Program, Massachusetts Division of Fisheries and Wildlife,.

USNVC [United States National Vegetation Classification]. 2017 (Date accessed). United States National Vegetation Classification Database V2.01. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC.

### Other references

Swain, P.C. and J.B. Kearsley. 2011. Classification of the Natural Communities of Massachusetts. Version 1.4. Natural Heritage & Endangered Species Program, Massachusetts Division of Fisheries and Wildlife. Westborough, MA.

### Contributors

Nels Barrett, Ph.D.  
Michael Margo

### Approval

Nels Barrett, 5/23/2020

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/24/2024
Approved by	Nels Barrett
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

1. **Number and extent of rills:**

- 
2. **Presence of water flow patterns:**
- 
3. **Number and height of erosional pedestals or terracettes:**
- 
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
- 
5. **Number of gullies and erosion associated with gullies:**
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:**
- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or**

decadence):

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---