

Ecological site R149BY002MA

Coastal Dunes

Last updated: 5/21/2020

Accessed: 05/15/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 149B—Long Island-Cape Cod Coastal Lowland

149B—Long Island-Cape Cod Coastal Lowland

This area is in the Embayed Section of the Coastal Plain Province of the Atlantic Plain. It is part of the partially submerged coastal plain of New England. It is mostly an area of nearly level to rolling plains, but it has some steeper hills (glacial moraines). Ridges border the lower plains. Elevation generally ranges from sea level to 80 feet (0 to 25 meters), but it is as much as 330 feet (100 meters) in a few areas. Local relief is mainly 3 to 30 feet (1 to 10 meters), but it is as much as 65 feet (20 meters) in the more hilly areas. The Peconic and Carmans Rivers are on the eastern end of Long Island. The parts of this area in Massachusetts and Rhode Island have no major rivers.

This entire area is made up of deep, unconsolidated glacial outwash deposits of sand and gravel. A thin mantle of glacial till covers most of the surface. Some moraines form ridges and higher hills in this area of generally low relief. Sand dunes and tidal marshes are extensive along the coastline.

Ecological site concept

The site consists of very deep, excessively drained and poorly developed eolian sands derived from marine deposits. The site occurs on both active and stabilized coastal sand dunes. Slopes range from 0 to 35 percent. Representative soil is Hooksan.

Associated sites

| | |
|-------------|---|
| R149BY012MA | Coastal Backbarrier Dune Flats Coastal Backbarrier Dune Flats |
|-------------|---|

Table 1. Dominant plant species

| | |
|------------|------------------------------------|
| Tree | Not specified |
| Shrub | (1) <i>Morella pensylvanica</i> |
| Herbaceous | (1) <i>Ammophila breviligulata</i> |

Physiographic features

The site occurs on both active and stabilized coastal sand dunes influenced by a maritime environment of wind exposure, shifting sands, and salt spray in varying degrees depending on the distance from the coast. Slopes range from 0 to 35 percent.




Figure 1.

Table 2. Representative physiographic features

| | |
|-----------|----------|
| Landforms | (1) Dune |
| Slope | 0–35% |

Climatic features

Mean annual precipitation is 48 inches and is usually uniformly distributed throughout the year. Frost free and freeze free days average 177 and 204, respectively.

Table 3. Representative climatic features

| | |
|-------------------------------|----------|
| Frost-free period (average) | 177 days |
| Freeze-free period (average) | 204 days |
| Precipitation total (average) | 1,219 mm |




Figure 2. Monthly precipitation range




Figure 3. Monthly average minimum and maximum temperature




Figure 4. Annual precipitation pattern

Climate stations used

- (1) HYANNIS [USC00193821], Hyannis, MA
- (2) BRIDGEHAMPTON [USC00300889], Sag Harbor, NY

Influencing water features

Soil features

The site consists of very deep, excessively drained and poorly developed eolian sands derived from marine deposits. The site occurs on both active and stabilized coastal sand dunes. Slopes range from 0 to 35 percent. Representative soil is Hooksan.



Figure 6. Hooksan photo by Pete Fletcher, Horseneck Beach, W

Table 4. Representative soil features

| | |
|--------------------|--|
| Surface texture | (1) Sand (2) Fine sand (3) Coarse sand |
| Drainage class | Excessively drained |
| Permeability class | Very rapid |
| Soil depth | 183 cm |

Ecological dynamics

[Caveat: The vegetation information contained in this section and is only provisional, based on concepts, not yet validated with field work.*]

The vegetation groupings described in this section are based on the terrestrial ecological system classification and vegetation associations developed by NatureServe (Comer et al. 2003). Terrestrial ecological systems are specifically defined as a group of plant community types (associations) that tend to co-occur within landscapes with similar ecological processes, substrates, and/or environmental gradients. They are intended to provide a classification unit that is readily mappable, often from terrain and remote imagery, and readily identifiable by conservation and resource managers in the field. A given system will typically manifest itself in a landscape at intermediate geographic scales of tens-to-thousands of hectares and will persist for 50 or more years. A vegetation association is a plant community that is much more specific to a given soil, geology, landform, climate, hydrology, and disturbance history. It is the basic unit for vegetation classification and recognized by the US National Vegetation Classification (FDGC 2008, USNVC 2017). Each association will be named by the diagnostic and often dominant species that occupy the different height strata (tree, shrub, and herb). Within the NatureServe Explorer database, ecological systems are numbered by a community Ecological System Code (CES) and individual vegetation associations are assigned an identification number called a Community Element Global Code (CEGL) (NatureServe 2019).

[*Caveat] The information presented is representative of very complex vegetation communities. Key indicator plants and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and geography. The reference plant community is not necessarily the management goal. The drafts of species lists are merely representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.


This ecological site is characterized by a mix of xeric plant communities coinciding Northern Atlantic Coastal Plain Dune and Swale system (CES203.264). The prevailing ecological processes are due to maritime environment, including frequent salt spray, wind exposure, overwash, and sand movement. Other disturbances considered as threats include: development and fragmentation, barriers to connectivity to the open ocean, off-road vehicles, trampling, and invasive plants: Japanese black pine (*Pinus thunbergii*), Japanese sedge (*Carex kobomugi*), and common reed (*Phragmites australis*).

Plant community succession on the site coincides with coastal dune succession. Young, active foredunes, are nearest the ocean and are exposed to significant environmental stresses of onshore winds, high light intensity, high temperatures, and salt spray. Vegetation on the young dunes is a grassland community dominated by American beachgrass (*Ammophila breviligulata*). Beachgrass plays a valuable role in jump starting succession and in stabilizing dunes. Beach pea is a common associate and can dominate at times (Sneddon et al. 2010). Other associated species includes American searocket (*Cakile edentula*), and seaside sandmat (*Chamaesyce polygonifolia*), pinweed (*Lechea maritima*), seabeach sedge (*Carex silicea*), coastal jointweed (*Polygonella articulata*), and seaside goldenrod (*Solidago sempervirens*). Northern bayberry (*Morella pensylvanica*), beach plum (*Prunus maritima*), and pitch pine (*Pinus rigida*) can occur sporadically within the grassland. Common exotic and invasive plants are Japanese black pine (*Pinus thunbergii*), beach rose (*Rosa rugosa*) and beach wormwood [dusty miller] (*Artemisia stelleriana*). The community is interspersed with small to large areas of bare sand. The active foredune communities coincides with Maritime Dune Community (Swain and Kearsley 2011) and Northern Beachgrass dune community (Sneddon et al. 2010). Northern bayberry dune shrublands (Sneddon et al. 2010) are a mid-seral community occurring on more stabilized dunes. Other shrubs include beach plum (*Prunus maritima*), Canadian serviceberry (*Amelanchier canadensis*), and occasionally, eastern baccharis (*Baccharis halimifolia*).

American beachgrass (*Ammophila breviligulata*), wavy hairgrass (*Deschampsia flexuosa*) and seaside goldendrod (*Solidago sempervirens*) are interspersed among the shrubs. On older and more stabilized backdunes or secondary dunes, a pitch pine dune woodland (Sneddon et al. 2010) community will occur. This woodland community is dominated by pitch pine (*Pinus rigida*) with scattered black oak (*Quercus velutina*). Other trees include black cherry (*Prunus serotina*), red maple (*Acer rubrum*), and American Beech (*Fagus grandifolia*). The understory is can be very open with northern bayberry (*Morella pensylvanica*) and beach plum (*Prunus maritima*) (Sneddon et al. 2010). Woolly beachheather (*Hudsonia tomentosa*) and kinnikinnik (*Arctostaphylos uva-ursi*) are found in open areas. Herbaceous plants include starflower (*Trientalis borealis*), whorled yellow loosestrife (*Lysimachia quadrifolia*), wild sarsaparilla (*Aralia nudicaulis*), western brackenfern, and wavy hairgrass (*Deschampsia flexuosa*). Common vines include eastern poison ivy (*Toxicodendron radicans*), roundleaf greenbrier (*Smilax rotundifolia*), and Virginia creeper (*Parthenocissus quinquefolia*).

State and transition model

149BY002 – Coastal Dunes



| Transition | Drivers/practices |
|------------|---|
| T1-2 | disturbance, invasive plant establishment |
| T1-3, T2-3 | cutting, land clearing, plant establishment, wind erosion control |
| R2-1, R3-1 | herbaceous weed treatment, plant removal, plant establishment, successional management |
| T3-2 | abandonment, disturbance, invasive plant establishment |
| CP1.1-1.2 | succession |
| CP1.2-1.1 | disturbance, herbaceous weed treatment, plant removal, plant establishment, successional management |

State 1

Reference State - Coastal Dunes

This ecological site is characterized by a mix of xeric plant communities coinciding Northern Atlantic Coastal Plain Dune and Swale system (CES203.264) (NatureServe 2019). The predominant reference plant community is largely herbaceous: • Northern Beachgrass Dune, (American Beachgrass - Beach Pea Grassland, [*Ammophila breviligulata* - *Lathyrus japonicus* Grassland] - CEGLO06274); and under stable dune conditions, it may succeed to shrublands, predominantly: • Northern Bayberry Dune Shrubland (Northern Bayberry - Beach Plum Shrubland, [*Morella pensylvanica* - *Prunus maritima* Shrubland] – CEGLO06295). Other associated plant communities include: • Northern Beach-heather Dune Dwarf-shrubland (Woolly Beach-heather - Bearberry Dwarf-shrubland [*Hudsonia tomentosa* - *Arctostaphylos uva-ursi* Dwarf-shrubland] – CEGLO06143), and • North Atlantic Coastal Plain Dune Vine/shrubland (Cat Greenbrier - Eastern Poison-ivy Vine-Shrubland [*Smilax glauca* - *Toxicodendron radicans* Vine-Shrubland] – CEGLO03886). On older more stabilized backdunes may support: • Pitch Pine Dune Woodland

(Pitch Pine / Woolly Beach-heather Woodland [*Pinus rigida* / *Hudsonia tomentosa* Woodland] – CEG006117). • Long Island Maritime Beech Forest (American Beech / Roundleaf Greenbrier Forest [*Fagus grandifolia* / *Smilax rotundifolia* Forest – CEG006043]) [Source: NatureServe 2019, USNVC 2017]

Community 1.1

Northern Beachgrass Dune

Northern Beachgrass Dune, (American Beachgrass - Beach Pea Grassland, [*Ammophila breviligulata* - *Lathyrus japonicus* Grassland] - CEG006274) is the typical community type. This association is characterized and dominated by American beachgrass (*Ammophila breviligulata*), which is often the only plant present, especially on foredunes or other areas of active and rapid sand deposition. Beach pea (*Lathyrus japonicus*) is the most common associate and sometimes codominant. Other associated species include seaside goldenrod (*Solidago sempervirens*), beach pinweed (*Lechea maritima*), seaside threeawn (*Aristida tuberculosa*), little bluestem (*Schizachyrium scoparium*), beach sedge (*Carex silicea*), coastal jointweed (*Polygonella articulata*), and dusty miller (*Artemisia stelleriana*). Dwarf-shrubs, such as woolly beachheather (*Hudsonia tomentosa*), beach rose (*Rosa rugosa*), northern bayberry (*Morella pensylvanica* [= *Myrica pensylvanica*]), or stunted beach plum (*Prunus maritima*), can occur sporadically and form patches within the grassland [Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]. Cross-referenced plant community concepts (typically by political state): Maritime Dunes (Swain 2016) [MA] Maritime Dunes (Edinger et al. 2014) [NY] Maritime Herbaceous Dune (Enser et al. 2011) [RI] Northern Beachgrass Dune (Sneddon et al 2010) [Cape Cod National Seashore]

Community 1.2

Northern Bayberry Dune Shrubland

Northern Bayberry Dune Shrubland (Northern Bayberry - Beach Plum Shrubland, [*Morella pensylvanica* - *Prunus maritima* Shrubland] – CEG006295) typically occurs on more stabilized dunes. Northern bayberry (*Morella pensylvanica* [= *Myrica pensylvanica*]) and beach plum (*Prunus maritima*) dominate. Another common and often codominant shrub is seaside rose (*Rosa rugosa*), while not native, where naturalized it's restricted to duneland sites. Other common shrubs or stunted trees in less abundance can include: eastern baccharis (*Baccharis halimifolia*) and sporadic red cedar (*Juniperus virginiana*) and black cherry (*Prunus serotina*). The herbaceous layer tends to be sparse where shrub growth is dense, and can include dune grassland species: American beachgrass (*Ammophila breviligulata*), seaside goldenrod (*Solidago sempervirens*), woolly beachheather (*Hudsonia tomentosa*), beach pinweed (*Lechea maritima*), Greene's rush (*Juncus greenii*), seabeach sedge (*Carex silicea*), coastal jointweed (*Polygonella articulata*), annual saltmarsh American-aster (*Symphotrichum subulatum* [= *Aster subulatus*]) or occasionally adjacent upland species such as, common wrinkleleaved goldenrod *Solidago rugosa* ssp. *aspera*, small flowered evening primrose (*Oenothera parviflora*), little bluestem (*Schizachyrium scoparium*), and others. Typical vine associates are poison ivy (*Toxicodendron radicans*), Virginia creeper *Parthenocissus quinquefolia*, and various Greenbriers (*Smilax* spp). Open, bare patches to sparsely vegetated sand are present in some examples [Source: NatureServe 2018 [accessed 2019], USNVC 2017 [accessed 2019]. Cross-referenced plant community concepts (typically by political state): Maritime Shrubland (Swain 2016) [MA] Maritime Shrubland (Edinger et al. 2014) [NY] Maritime Shrub Dune (Enser et al. 2011) [RI] Northern Bayberry Dune Shrubland (Sneddon et al 2010) [Cape Cod National Seashore]

Pathway CP1.1-1.2

Community 1.1 to 1.2

succession

Pathway CP1.2-1.1

Community 1.2 to 1.1

disturbance, herbaceous weed treatment, plant removal, plant establishment, successional management

Conservation practices

| |
|---|
| Restoration and Management of Rare and Declining Habitats |
| Upland Wildlife Habitat Management |

| |
|--|
| Restoration and Management of Natural Ecosystems |
| Herbaceous Weed Control |

State 2

Semi-Natural State

Semi-natural dunescapes may be dominated by occasional non-native plants include Japanese Sedge (*Carex kobomugi*), dusty miller (*Aretemisia stelleriana*), sea poppy (*Glaucium flavum*), beach rose (*Rosa rugosa*), Morrow's honeysuckle (*Lonicera morrowii*), and Japanese pine (*Pinus thunbergii*).

Community 2.1

Invasive plant community

Carex kobomugi (Japanesese sedge) *Ammophila arenaria* (European beechgrass)

State 3

Cultural State

The non-native Japanese Pine (*Pinus thunbergii*) and beach rose (*Rosa rugosa*) may have been be planted.

Community 3.1

Plantation(?) *Pinus thunbergii*

Transition T1-2

State 1 to 2

disturbance, invasive plant establishment

Transition T1-3

State 1 to 3

cutting, land clearing plant establishment, wind erosion control

Conservation practices

| |
|--------------------------------|
| Brush Management |
| Land Clearing |
| Invasive Plant Species Control |

Restoration pathway R2-1

State 2 to 1

herbaceous weed treatment, plant removal, plant establishment, successional management American beachgrass (*Ammophila breviligulata*) plugs generally planted on a grid (0.5 m on center?) [Avoid planting the non-native European beachgrass (*Ammophila arenaria*)]

Conservation practices

| |
|---|
| Brush Management |
| Critical Area Planting |
| Restoration and Management of Rare and Declining Habitats |
| Upland Wildlife Habitat Management |
| Early Successional Habitat Development/Management |

| |
|--|
| Record Keeping |
| Restoration and Management of Natural Ecosystems |
| Invasive Plant Species Control |
| Herbaceous Weed Control |

Transition T2-3

State 2 to 3

cutting, land clearing, plant establishment, wind erosion control

Conservation practices

| |
|--------------------------------|
| Land Clearing |
| Invasive Plant Species Control |

Restoration pathway R3-1

State 3 to 1

herbaceous weed treatment, plant removal, plant establishment, successional management

Conservation practices

| |
|---|
| Brush Management |
| Critical Area Planting |
| Restoration and Management of Rare and Declining Habitats |
| Upland Wildlife Habitat Management |
| Early Successional Habitat Development/Management |
| Restoration and Management of Natural Ecosystems |
| Herbaceous Weed Control |

Transition T3-2

State 3 to 2

abandonment, disturbance, invasive plant establishment

Additional community tables

Animal community

Wildlife

Terns, gulls, shorebirds and some songbirds nest in dune especially Piping Plover and Least Tern which prefer sparse vegetation. Dunes may also provide food and cover for some migrating birds. Diamondback Terrapins nest in dunes. Moths that inhabit stable dunes with patches of low shrubs include the Chain-dotted Geometer and Dune Sympistis.

Other information

Conservation Strategies

Maintain dynamic coastal process allowing dunes to develop; maintain or enhance connectivity across the beach-dune complex and avoid development that contributes to fragmentation; prevent recreational overuse, e.g., driving and trampling, and encourage the public to use marked trails; effect dune restoration where excessive disturbance warrants, e.g., American beachgrass planting for stabilization, removal of invasive plants for ecological integrity.

Inventory data references

Site Development and Testing Plan

Future work is needed, as described in a project plan, to validate the information presented in this provisional ecological site description. Future work includes field sampling, data collection and analysis by qualified vegetation ecologists and soil scientists. As warranted, annual reviews of the project plan can be conducted by the Ecological Site Technical Team. A final field review, peer review, quality control, and quality assurance reviews of the ESD are necessary to approve a final document.

References

- Comer, P., D. Faber-Langendoen, R. Evans, S. Grawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schultz, K. Snow, and J. Teague. 2003. Ecological Systems of the United States: A Working Classification of U.S. Terrestrial Systems. NatureServe, Arlington, Virginia..
- Edinger, G.J., D.J. Evans, S. Gebauer, T.J. Howard, D. Hunt, and A. Olivero. 2014. Ecological Communities of New York State, Second Edition: A revised and expanded edition of Carol Reschke's Ecological Communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation, Albany, NY..
- Enser, R., D. Gregg, C. Sparks, P. August, P. Jordan, J. Coit, C. Raithel, B. Tefft, B. Payton, C. Brown, and C. LaBash. 2011.. Rhode Island ecological communities classification. Rhode Island Natural History Survey, Kingston, RI..
- FGDC [Federal Geographic Data Committee]. 2008. National Vegetation Classification Standard, Version 2. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC..
- NatureServe. 2018 (Date accessed). NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available <http://explorer.natureserve.org>.
<http://explorer.natureserve.org>.
- Smith, S.M. and K.T. Killingbeck. 2008. Development of vegetation in dune slack wetlands of Cape Cod National Seashore (Massachusetts, USA). Plant Ecology 194:243–256.
- Swain, P.C. 2016. Classification of the natural communities of Massachusetts, Natural Heritage & Endangered Species Program, Massachusetts Division of Fisheries and Wildlife,.
- USNVC [United States National Vegetation Classification]. 2017 (Date accessed). United States National Vegetation Classification Database V2.01. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC.

Contributors

Nels Barrett, Ph.D.
Michael Margo

Approval

Nels Barrett, 5/21/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

| | |
|---|-------------------|
| Author(s)/participant(s) | |
| Contact for lead author | |
| Date | 05/15/2024 |
| Approved by | Nels Barrett |
| Approval date | |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
