

# Ecological site F152BY007TX Poorly Drained Loamy Upland

Last updated: 9/22/2023  
Accessed: 05/06/2024

## General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

## MLRA notes

Major Land Resource Area (MLRA): 152B–Western Gulf Coast Flatwoods

Major Land Resource Area (MLRA) 152B, Western Gulf Coast Flatwoods, is in eastern Texas and western Louisiana. Locally termed the Flatwoods, the area is dominated by coniferous forest covering 5,681 square miles (14,714 square kilometers). The region is a hugely diverse transition zone between the northern/eastern mixed forests and southern/western coastal prairies and grasslands.

## Classification relationships

Major Land Resource Area (MLRA) (USDA-Natural Resources Conservation Service, 2006)

## Ecological site concept

The Poorly Drained Loamy Upland ecological site has very deep, poorly drained soils. The poor drainage pattern causes water to persist for long periods of time after precipitation events. The resulting hydrophytic plant communities are able to withstand the presence of water.

## Associated sites

F152BY004TX	<b>Clayey Flat</b> Soils are comprised of shrink-swell clays.
F152BY005TX	<b>Seasonally Wet Loamy Upland</b> Soils have seasonally high water table and are not as poorly drained.
F152BY006TX	<b>Well Drained Loamy Upland</b> Soils are well drained.
F152BY008TX	<b>Acid Baygall</b> Soils have a spodic horizon.
F152BY010TX	<b>Terrace</b> Soils are on terraces.
F152BY001TX	<b>Depressional</b> Soils are on a lower landform and stay ponded longer.
F152BY002TX	<b>Sodic Flats</b> Soils have high salt concentrations and bioturbation.
F152BY013TX	<b>Poorly Drained Loamy Bottomland</b> Soils are on a lower landform and flooded for extended periods.
F152BY014TX	<b>Poorly Drained Clayey Bottomland</b> Soils are clayey and on floodplains.

### Similar sites

F152BY001TX	<b>Depressional</b> Soils are on a lower landform and stay ponded longer.
F152BY004TX	<b>Clayey Flat</b> Soils are comprised of shrink-swell clays.
F152BY005TX	<b>Seasonally Wet Loamy Upland</b> Soils have seasonally high water table and are not as poorly drained.
F152BY002TX	<b>Sodic Flats</b> Soils are influenced by salt and bioturbation.

**Table 1. Dominant plant species**

Tree	(1) <i>Pinus taeda</i> (2) <i>Pinus palustris</i>
Shrub	(1) <i>Sabal minor</i>
Herbaceous	(1) <i>Eleocharis</i> (2) <i>Saururus cernuus</i>

### Physiographic features

The ecological site occurs on relict bars on terrace risers. Slope ranges from 0 to 3 percent, but are most commonly 0 to 1 percent. Elevation ranges from 10 to 151 feet. Water table depth fluctuates throughout the year. From October to May, the depth to the top of the water table will be 0 to 16 inches. The water table will deepen during the warmer months of the year.

**Table 2. Representative physiographic features**

Landforms	(1) Coastal plain > Flat (2) Coastal plain > Depression
Runoff class	Negligible to high
Flooding frequency	None
Ponding duration	Long (7 to 30 days)

Ponding frequency	None to frequent
Elevation	3–46 m
Slope	0–3%
Ponding depth	0–46 cm
Water table depth	0–41 cm
Aspect	Aspect is not a significant factor

## Climatic features

The Western Gulf Coast Flatwoods (MLRA 152B) is within the humid subtropical climate zone. The region boasts one of the highest rainfall averages in the southern United States, over 60 inches (152 centimeters) annually. This is due to the gulf currents that carry humid air to the region, where it condenses and precipitates. Rainfall averages are fairly consistent month by month, ranging from the lowest of 3.5 inches (8.9 centimeters) in March and the highest of 5.6 inches (14.3 centimeters) in June.

The area is prone to severe thunderstorms and tornadoes when the proper conditions exist, generally in the springtime. Sometimes excessive rainfall occurs, leading to flooding. Hurricanes also strike the region, generally in late summer or early fall. These extreme weather events can be quite destructive, toppling trees, and serves to naturally reset the vegetation to primary succession. The higher humidity of the region amplifies the feeling of heat during the summer. Prolonged droughts and snowfall events are rare.

**Table 3. Representative climatic features**

Frost-free period (average)	249 days
Freeze-free period (average)	289 days
Precipitation total (average)	1,600 mm

## Climate stations used

- (1) CLEVELAND [USC00411810], Cleveland, TX
- (2) LIBERTY [USC00415196], Liberty, TX
- (3) LUMBERTON [USC00415435], Silsbee, TX
- (4) TOWN BLUFF DAM [USC00419101], Jasper, TX
- (5) DE RIDDER [USC00162367], Deridder, LA
- (6) DE QUINCY [USC00162361], Dequincy, LA
- (7) ELIZABETH [USC00162800], Oakdale, LA
- (8) WILDWOOD [USC00419754], Kountze, TX
- (9) OBERLIN FIRE TWR [USC00166938], Oberlin, LA
- (10) ORANGE 9 N [USC00416680], Orange, TX

## Influencing water features

Water affects these soils due to their poor drainage. Water can be ponded up to 18 inches for extended periods of time. This influences the plants and many are classified as hydrophytic.

## Wetland description

The soils are hydric and are potentially wetlands. Onsite field determinations using water, soil, and plant indicators should be used to verify.

## Soil features

The site consists of very deep, poorly to very poorly drained soils formed in loamy fluviomarine deposits. The representative series are Bissonnet, Camptown, Evadale, Jasco, Kinder, Olive, Sorter, Tyden, and Waller. Since so

many soils belong to this ecological site, their taxonomies vary, but all belong to the orders of alfisols and ultisols. The soils extend throughout the whole region and are widely prolific.

**Table 4. Representative soil features**

Parent material	(1) Fluviomarine deposits–igneous, metamorphic and sedimentary rock
Surface texture	(1) Loam (2) Silt loam (3) Fine sandy loam
Family particle size	(1) Fine-silty (2) Coarse-silty
Drainage class	Poorly drained to very poorly drained
Permeability class	Slow to very slow
Soil depth	203 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-152.4cm)	22.86–33.02 cm
Calcium carbonate equivalent (0-152.4cm)	0%
Electrical conductivity (0-152.4cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-152.4cm)	0–6
Soil reaction (1:1 water) (0-152.4cm)	3.5–6
Subsurface fragment volume <=3" (61-152.4cm)	0–2%
Subsurface fragment volume >3" (Depth not specified)	0%

## Ecological dynamics

The information in this ecological site description (ESD), including the state-and-transition model (STM), was developed using archeological and historical data, professional experience, and scientific studies. The information is representative of a complex set of plant communities. Not all scenarios or plants are included. Key indicator plants, animals, and ecological processes are described to inform land management decisions.

**Introduction** – In southeastern Texas and southwestern Louisiana the transition from coastal grasslands to the large expanse of coniferous forest has been deemed the “Flatwoods”. As the name suggests, the region is relatively flat and, with many transitional areas, highly diverse in flora and fauna. Historically, the area was covered by pines with mixed hardwoods, sparse shrubs, and a diverse understory of grasses and forbs. Fire and drainage patterns play a significant role in shaping the plant communities and their development. Fire suppression, drainage alterations, and land conversion have reduced the amount of historical communities in existence today.

**Background** – Prior to settlement by the Europeans, the reference state for the Poorly Drained Loamy Uplands were Loblolly Pine/Longleaf Pine Woodlands. Remnants of this presumed historic plant community still exist where natural conditions are intact. Evidence of the reference state is found in accounts of early historic explorers to the area, historic forest and biological survey teams, as well as recent ecological studies in the last 30 years. The age of this woodland community varies, and has a diverse flora.

**Settlement Management** – As human settlement increased throughout the area, so did the increase in logging and grazing by domestic livestock. The logging became so extensive that by the 1930’s most of the region had been cut-over. Replanting trees to historic communities was not common and early foresters began planting loblolly pine

(*Pinus taeda*) for its quick growth. As more people colonized they began suppressing fire, which allowed dense thickets of shrubs to replace the herbaceous understory.


**Current Management and State** – Today much of the historic forest is gone, replaced by pine plantations, crops, and pastures. The areas that were not converted have been fire-suppressed so long that loblolly pine and fire intolerant hardwoods populate the overstory structure. Currently, federally-managed properties are the best place to view the remnant sites (National Park Service, U.S. Fish and Wildlife Service, etc.). Some private individuals have begun restoring communities through selective tree planting and retention of communities that remain. Other restoration efforts include mimicking natural-disturbance regimes through gap-phase regeneration on plantation sites.

**Fire Regimes** – Fire was a natural and important disturbance throughout the region. Fire occurred naturally from lightning strikes, by Native Americans for game movement, and eventually early European settlers. Fires throughout the Flatwoods occurred at two different times. Early in the year, they would occur during late winter and early spring, removing senescent vegetation, recycling nutrients and minerals, and spurring new plant growth. Late summer and early fall fires occurred as well, but with a different community effect. Summer fires burned hotter and with more intensity, greatly suppressing the shrub canopy layer. The summer fires also shifted the ecological site transitional state by decreasing grass densities and increasing forb densities. The topography, fuel loads, and other conditions caused patchy burns throughout the region resulting in mosaic patterns of plant communities and a heterogeneous landscape.

**Disturbance Regimes** – Extreme weather events occur occasionally throughout the region. Tornadoes uproot trees and open canopies in the spring months. In the late summer and early fall, hurricanes or tropical depressions can make landfall, dumping excessive amounts of rain and toppling trees with high winds. Another cause of large canopy openings is the effects of the southern pine beetle (*Dendroctonus frontalis*). Starting in the late 1950's, beetle outbreaks have occurred every 6 to 9 years (although a major attack has not occurred in some time); usually when the trees are stressed due to multiple environmental factors.

## State and transition model

### Ecosystem states



**T1A** - Absence of disturbance, coupled with natural regeneration over time

**T1B** - Merchantable timber is harvested by clearcut and site is planted to a monoculture of pine trees

**T1C** - Removal of native vegetation and introduction of improved forage species or annual crops

**R2A** - Reduction of overstory canopy using fire and selective thinning

**T2A** - Merchantable timber is harvested by clearcut and site is planted to a monoculture of pine trees

**T2B** - Merchantable timber harvested by clearcut, followed by planting of improved forage species or annual crops

**R3A** - Selective harvest combined with reintroduction of natural disturbances and native species

**T3B** - Lack of natural/anthropogenic disturbance and natural regeneration over time

**T3A** - Timber harvest by clearcut, followed by planting improved forage species or annual crops

**T4A** - Lack of natural/anthropogenic disturbance and natural regeneration over time

**T4B** - Site is planted to a monoculture of pine trees

#### State 1 submodel, plant communities

1.1. Loblolly  
Pine/Longleaf Pine  
Woodland

#### State 2 submodel, plant communities

2.1. Pine/Hardwood  
Forest

#### State 3 submodel, plant communities

3.1. Plantation

#### State 4 submodel, plant communities

4.1. Planted Pasture  
and Row Crop

### State 1 Reference

The Poorly Drained Loamy Upland ecological site is a Loblolly Pine/Longleaf Pine Woodland. The deep loamy soils provide an excellent growing medium for plants, but the poor drainage restricts what plants will grow. Because of ponding and extended soil saturation, many of the plants are classified as facultative wetland (FACW) and obligate (OBL) by the U.S. Army Corps of Engineers Wetland Delineation Manual. Despite the influence of water, the ecological site has a moderately frequent burning regime of 5 to 10 years. The site ranges from 60 to 80 percent canopy cover with basal areas from 75 to 105 square feet per acre.

#### Dominant plant species

- loblolly pine (*Pinus taeda*), tree
- longleaf pine (*Pinus palustris*), tree

### Community 1.1 Loblolly Pine/Longleaf Pine Woodland



The overstory canopy is dominated by loblolly pine, with longleaf pine as a co-dominant. Other species such as sweetgum (*Liquidambar straciflua*), sweetbay (*Magnolia virginiana*), and blackgum (*Nyssa sylvatica*) may be present, but not in large numbers. The midstory and understory vary significantly, especially with time since last fire. An herbaceous understory will be found directly after a burn and more shrub species will be present as the time since fire lengthens. Common midstory shrubs and trees include wax myrtle (*Morella cerifera*), dwarf palmetto (*Sabal minor*), green ash (*Fraxinus pennsylvanica*), and redbay (*Persea borbonia*). Understory species include a variety of sedges (*Carex* sp.), flatsedges (*Cyperus* sp.), rushes (*Juncus* sp.), spikerushes (*Eleocharis* sp.), and fimbry (*Fimbristylis* sp.). The presence of spikerushes is oftentimes an indicator of the site. Lizards tail (*Saururus cernuus*) is the most prolific forb species.

## State 2 Encroached

A long-term lack of fire and management has caused the community to cross a threshold, resulting in an Encroached State (2). The crossing of this threshold represents a closure in the overstory canopy, which limits the productivity of the ground layer. The limited ground layer does not provide enough fuel to harbor a burn with the intensity found in State 1. Fire-intolerant hardwoods have become part of the overstory. The overstory trees are overstocked and limit the growth of neighboring species. The overstocking reduces tree growth and causes stress, making them vulnerable to attacks from insects and/or diseases. Longleaf recruitment may be nonexistent due to lack of light and bare ground. Loblolly pine will especially take advantage of the current conditions. The plant communities will stay in this constant state and continue to age without disturbance or intervention.

## Community 2.1 Pine/Hardwood Forest



The understory plant layer only contains remnants of the reference community and possibly no reference community indicator species. Shade-tolerant grasses, such as longleaf woodoats (*Chasmanthium sessiliflorum*), forbs, and greenbriers (*Smilax* sp.) may be the only ground-layer species. Because the site lacks the diversity of the reference state, the wildlife diversity will be limited to generalist species, species requiring a closed canopy, and

those seeking refuge.

### **State 3 Plantation**

The Plantation State is a result of conversion activities. The landowner has maximized silviculture production by planting a monoculture of pine species, usually loblolly pine, but sometimes slash pine (*Pinus elliotti*) is planted.

#### **Community 3.1 Plantation**

In the immediate years following the initial plantation tree planting, the understory community will resemble the reference state (State 1). During this early growth period, the landowner will typically remove unwanted hardwoods and herbaceous plants to reduce competition with the planted pine trees. As the overstory canopy closes, less understory management is required due to sunlight restrictions to the ground layer.

### **State 4 Pasture and Crop**

The Pasture and Crop state is a result of conversion activities. The landowner has maximized agriculture production by planting a monoculture of introduced grass species or agricultural row crops.

#### **Community 4.1 Planted Pasture and Row Crop**

Typical introduced pasture grass species include bahiagrass (*Paspalum notatum*) and different varieties of bermudagrass (*Cynodon dactylon*). The grasses are grown for livestock production through direct grazing or baling hay for later use. Agricultural row crops are grown for food and fiber production. Many farmers use herbicides to reduce unwanted plant competition which yields a plant community unrepresentative of the reference (State 1) or subsequent vegetative states.

#### **Transition T1A State 1 to 2**

The transition from State 1 to State 2 is a result of time and long periods, greater than 15 years, of no fire and/or forest management practices. Without fire to suppress tree seedlings, biomass and diversity is lost from the grass and forb layers of the system.

#### **Transition T1B State 1 to 3**

The transition is due to the land manager maximizing agricultural production. Merchantable timber is harvested by clearcut. Then, the site is prepared and planted to either an improved grass or row crops.

#### **Transition T1C State 1 to 4**

The transition is due to the land manager maximizing agricultural production. Merchantable timber is harvested by clearcut, the site prepared and planted to either an improved grass or row crops.

#### **Restoration pathway R2A State 2 to 1**

Restoration of this community to the reference state begins with a selective timber harvest. Removing unwanted trees opens up the canopy, allowing sunlight penetration to the ground. Years of overstory growth have limited the fuel necessary to have an effective fire. Time will be needed to encourage understory growth. Once the herbaceous layer has established, more frequent than natural burns (3 to 5 years) may be required to suppress the woody



vegetation.

### **Transition T2A**

#### **State 2 to 3**

The transition is due to the land manager maximizing silviculture potential. Merchantable timber is harvested by clearcut. Then, the site is prepared and planted to a monoculture of pine trees.

### **Transition T2B**

#### **State 2 to 4**

The transition is due to the land manager maximizing agricultural potential. Merchantable timber is harvested by clearcut. Then, the site is prepared and planted to either an improved grass or row crops.

### **Restoration pathway R3A**

#### **State 3 to 1**

When restoring a plantation, the land manager can either clearcut the timber, the site prepared, and trees planted. Otherwise, gap-phase regeneration is possible through selective timber harvests. This involves replanting the desired overstory species in small openings within the current structure of the forest. The benefit is a slow progression of restoration instead of starting from primary succession.

### **Transition T3B**

#### **State 3 to 2**

This community transition is caused by neglecting the plantation understory. Without fire, mowing, or herbicides, unwanted understory saplings can begin to grow into the overstory.

### **Transition T3A**

#### **State 3 to 4**

The transition is due to the land manager maximizing agricultural production. Merchantable timber is harvested by clearcut. Then, the site is prepared and planted to either an improved grass or row crops.

### **Transition T4A**

#### **State 4 to 2**

This community transition is caused by neglecting crop or pasture. Without continuation of agricultural management, first-successional herbaceous plants will occupy the ground layer, followed by shrubs, and eventually shade-loving, fire-intolerant overstory species.

### **Transition T4B**

#### **State 4 to 3**

The transition is due to the land manager maximizing silviculture production. The site prepared and planted to a monoculture of pine trees.

## **Additional community tables**

Table 5. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (M)	Canopy Cover (%)	Diameter (Cm)	Basal Area (Square M/Hectare)
<b>Tree</b>							
sweetgum	LIST2	<i>Liquidambar styraciflua</i>	Native	–	–	–	–
blackgum	NYSY	<i>Nyssa sylvatica</i>	Native	–	–	–	–
loblolly pine	PITA	<i>Pinus taeda</i>	Native	–	–	–	–
longleaf pine	PIPA2	<i>Pinus palustris</i>	Native	–	–	–	–
sweetbay	MAV12	<i>Magnolia virginiana</i>	Native	–	–	–	–

Table 6. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (M)	Canopy Cover (%)
<b>Grass/grass-like (Graminoids)</b>					
flatsedge	CYPER	<i>Cyperus</i>	Native	–	–
needleleaf rosette grass	DIAC	<i>Dichanthelium aciculare</i>	Native	–	–
Ravenel's rosette grass	DIRA	<i>Dichanthelium ravenelii</i>	Native	–	–
spikerush	ELEOC	<i>Eleocharis</i>	Native	–	–
fimbry	FIMBR	<i>Fimbristylis</i>	Native	–	–
leathery rush	JUCO4	<i>Juncus coriaceus</i>	Native	–	–
sedge	CAREX	<i>Carex</i>	Native	–	–
Indian woodoats	CHLA5	<i>Chasmanthium latifolium</i>	Native	–	–
longleaf woodoats	CHSE2	<i>Chasmanthium sessiliflorum</i>	Native	–	–
little bluestem	SCSC	<i>Schizachyrium scoparium</i>	Native	–	–
yelloweyed grass	XYRIS	<i>Xyris</i>	Native	–	–
<b>Forb/Herb</b>					
lizard's tail	SACE	<i>Saururus cernuus</i>	Native	–	–
sidebeak pencilflower	STBI2	<i>Stylosanthes biflora</i>	Native	–	–
Texas ironweed	VETE3	<i>Vernonia texana</i>	Native	–	–
fourvalve mimosa	MIQU2	<i>Mimosa quadrivalvis</i>	Native	–	–
eastern poison ivy	TORA2	<i>Toxicodendron radicans</i>	Native	–	–
<b>Fern/fern ally</b>					
western brackenfern	PTAQ	<i>Pteridium aquilinum</i>	Native	–	–
<b>Shrub/Subshrub</b>					
American beautyberry	CAAM2	<i>Callicarpa americana</i>	Native	–	–
yaupon	ILVO	<i>Ilex vomitoria</i>	Native	–	–
wax myrtle	MOCE2	<i>Morella cerifera</i>	Native	–	–
dwarf palmetto	SAMI8	<i>Sabal minor</i>	Native	–	–
<b>Tree</b>					
American holly	ILOP	<i>Ilex opaca</i>	Native	–	–
redbay	PEBO	<i>Persea borbonia</i>	Native	–	–
green ash	FRPE	<i>Fraxinus pennsylvanica</i>	Native	–	–
<b>Vine/Liana</b>					
greenbrier	SMILA2	<i>Smilax</i>	Native	–	–
southern dewberry	RUTR	<i>Rubus trivialis</i>	Native	–	–

## Wood products

These soils occur in the Woodland Suitability Group 2w9. They have a high potential for woodland management, both pine and hardwood. The 50-year site index for loblolly pine averages 90 feet (approximately 60 feet on a 25-year curve). For oaks, the site index ranges from 75 to 85 feet. The yield from an unmanaged, natural stand of loblolly pine, over a 50-year period, is approximately 330 board feet (Doyle Rule), 2.64 tons, or 90 cubic feet per acre per year. Management can substantially increase this yield. Access and equipment operability on these soils is poor during wet periods due to saturation of the soil. Harvesting and other operations may need to be suspended during such periods. Wetness also makes these soils poorly suited for log landings and roads. Wetness and low strength will lead to severe rutting problems and make them moderately suited for road construction. Raising and crowning the road bed will be necessary and care must be taken to avoid interrupting the natural drainage. Site preparation operations should be limited to the dry months and planting should be planned for the drier part of the planting season. Use of herbicides for site preparation must also take into consideration the poor drainage on these soils. Applications should not be made during wet periods. Wetness may cause a moderate loss in pine seedling survival. Bedding may be needed.

## Type locality

Location 1: Hardin County, TX	
UTM zone	N
UTM northing	30.5453611
UTM easting	-94.417833
General legal description	Big Thicket National Forest – Sundew Trail

## Other references

### References

- Ajilvsgi, G. 2003. *Wildflowers of Texas*. Revised edition. Shearer Publishing, Fredericksburg, TX.
- Ajilvsgi, G. 1979. *Wildflowers of the Big Thicket*. Texas A&M University Press, College Station, TX.
- Allen, J. A., B. D. Keeland, J. A. Stanturf, and A. F. Kennedy Jr. 2001. *A guide to bottomland hardwood restoration*. Technical report, USGS/BRD/ITR-2000-0011.
- Bray, W. L. 1904. *Forest resources of Texas*. Bureau of Forestry Bulletin 47, Government Printing Office, Washington D.C.
- Diggs, G. M., B. L. Lipscomb, M. D. Reed, and R. J. O'Kennon. 2006. *Illustrated flora of East Texas*. Second edition. Botanical Research Institute of Texas & Austin College, Fort Worth, TX.
- Jones, S. D., J. K. Wipff, and P. M. Montgomery. 1997. *Vascular plants of Texas: a comprehensive checklist including synonymy, bibliography, and index*. University of Texas Press, Austin.
- Liu, C., P. A. Harcombe, and I. S. Elsik. 1990. *Fire study report, including Roy E. Larsen Preserve species list*. Summer 1990. Department of Ecology and Evolutionary Biology, Rice University, Houston, TX.
- Marks, P. L., and P. A. Harcombe. 1981. *Forest Vegetation of the Big Thicket, southeast Texas*. *Ecological Monographs* 51:287-305.
- Matos, J. A. 1985. *Roy E. Larsen Sandylands Sanctuary vascular plant species list*. Master thesis, Stephen F. Austin University, Nacogdoches, TX.
- NatureServe. 2002. *International classification of ecological communities: Terrestrial vegetation of the United States*. National forests in Texas final report. NatureServe, Arlington, VA.
- Nixon, E. S. 2000. *Trees, shrubs & woody vines of East Texas*. Second edition. Bruce Lyndon Cunningham Productions, Nacogdoches, TX.
- Randall, J. M., and J. Marinelli. 1996. *Invasive plants: weeds of the global garden*. Volume 149. Brooklyn Botanic Garden, Brooklyn, NY.
- Stanturf, J. A., S. H. Schoenholtz, C. J. Schweitzer, and J. P. Shepard. 2001. *Achieving restoration success: Myths in bottomland hardwood forests*. *Restoration Ecology*, 9:189-200.
- Stringham, T. K., W. C. Krueger, and P. L. Shaver. 2003. *State and transition modeling: An ecological process approach*. *Journal of Range Management* 56:106-113.
- Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. *Soil Survey Geographic (SSURGO) Database*.
- Truett, J. C. 1984. *Land of bears and honey: A natural history of East Texas*. The University of Texas Press, Austin,

TX.

Van Kley, J. E., R. L. Turner, L. S. Smith, and R. E. Evans. 2007. Ecological classification system for the national forests and adjacent areas of the West Gulf Coastal Plain. Second approximation. Stephen F. Austin University and The Nature Conservancy, Nacogdoches, TX.

USDA-NRCS Ag Handbook 296 (2006).

Vines, R. A. 1960. Trees, shrubs, and woody vines of the Southwest. University of Texas Press, Austin, TX.

Watson, G. E. 2006. Big Thicket Plant Ecology. Third Edition. University of North Texas Press, Denton, TX.

## Contributors

Tyson Hart

## Approval

Bryan Christensen, 9/22/2023

## Acknowledgments

Thanks to all involved during the preparation, sampling, and reviewing of the Flatwoods project. Thanks to Josh Berry, Dennis Brezina, Kenny Hall, Jason Hohlt, Stacey Kloesel, Ricky Lambert, Cody Langston, Mark Moseley, Ramiro Molina, Mike Oliver, Alan Peer, Sara Russell, Don Sabo, Mary Webb-Marek, and Jon Wiedenfeld for all their help.

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	09/21/2021
Approved by	Bryan Christensen
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native).** List species which **BOTH** characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-