

Ecological site R222XY356AK Alpine Dwarf Scrub Dry Gravelly Slopes

Last updated: 6/20/2019
Accessed: 05/20/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 222X–Southern Alaska Coastal Mountains

The Southern Alaska Coastal Mountains (MLRA 222) encompasses the Pacific Border Ranges and Coast Mountains physiographic provinces (Wahrhaftig 1965). Spanning approximately 26,355 square miles, the elevation ranges from sea level at the base of glaciers and ice fields to 18,008 feet at Mt. St. Elias (Figure 1). The MLRA was covered by glacial ice during the Pleistocene epoch, a time period spanning from 2.6 million to 11,700 years ago. During interglacial periods glacial extent was reduced, leaving behind various glacial deposits. Over time these deposits have been eroded or buried by colluvium and slope alluvium, which now covers more than 90 percent of the unglaciated landscape. Paleozoic, Mesozoic, and Lower Tertiary stratified sedimentary rocks, and occasionally Paleozoic intrusive rocks, underlie much of the area and are exposed on steep mountain slopes and ridges (USDA-NRCS 2006).

This landscape lies in the true alpine zone where glacial ice is, and has been, the dominant ground cover. Glacial ice encompassed all the MLRA during the Late Wisconsinan glaciation, 25,000 – 21,000 years ago (Kauffman et al. 2011). Changes in climatic conditions following this period resulted in the recession of some glaciers and slowly exposed new surfaces for inhabitation by terrestrial ecosystems. Pioneer plant communities began to establish on the new substrate within the first 30 years and consisted mostly of evergreen, herbaceous dwarf shrubs with some pockets of low shrubs. These communities quickly transitioned to tall shrubs within 100 years of deglaciation. By approximately 13,000 years ago, four stable plant communities emerged across the non-glaciated landscape –

ericaceous dwarf shrub, low shrub, tall shrub, and herbaceous communities – and form the present-day ecosystems (Boggs et al. 2010).

Classification relationships

USFS Ecoregion Province: Marine Mountains (M240), Forest-Meadow High (M242b) (Bailey 2007)

U.S. EPA Level III Ecoregion: Pacific Coastal Mountains (119) (Gallant et al. 2010)

National Vegetation Classification – Ecological Systems: Alaskan Pacific Maritime Alpine Sparse Shrub and Fell-Field (CES204.318) (NatureServe 2015)

Biophysical Settings: Alaskan Pacific Maritime Herbaceous Dwarf Shrubland (BpS 7816430) (LANDFIRE 2009)

Alaska Natural Heritage Program Landcover Class: Dwarf Shrub (Boggs et al. 2016)

Alaskan Vegetation Classification: Crowberry Dwarf Shrub Tundra (Viereck et al. 1992)

Ecological site concept

Alpine Dwarf Scrub Gravelly Slopes, Bedrock ecological sites occur on high-elevation mountain slopes that are exposed to harsh climatic conditions. The soils are gravelly Inceptisols that are well-drained and shallow, formed in colluvium over weathered residuum.

The reference vegetation on this ecological site is defined by a sparse cover of dwarf shrubs interspersed with large patches of bedrock and surface rock. Bog blueberry (*Vaccinium uliginosum* (Hook.) Coville) and black crowberry (*Empetrum nigrum* L.) are the dominant dwarf shrubs, and star reindeer lichen (*Cladina stellaris* (Opiz) Brodo) is the dominant non-vascular ground cover. Continuous exposure to cold temperatures, wind, and a short growing season maintain this plant community (LANDFIRE 2009).

Associated sites

R222XY352AK	Alpine Dwarf Scrub Dry Organic Slopes Organic soils that are shallow to bedrock
R222XY357AK	Alpine Dwarf Scrub Moist Gravelly Slopes Mineral soils that are shallow to bedrock, occurring on concave slope shapes

Similar sites

R222XY357AK	Alpine Dwarf Scrub Moist Gravelly Slopes Supports a similar dwarf shrub community but bedrock cover is less than 10 percent
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Vaccinium uliginosum</i> (2) <i>Empetrum nigrum</i>
Herbaceous	(1) <i>Cladina</i> (2) <i>Rock, bedrock</i>

Physiographic features

Alpine Dwarf Scrub Gravelly Slopes, Bedrock ecological sites occur on alpine mountain slopes. They are situated on elevations ranging from approximately 2500 to 8000 feet. This ecological site does not experience flooding, but rather generates runoff to adjacent, downslope ecological sites.




Figure 2. Representative block diagram of Alpine Dwarf Scrub Gravelly Slopes, Bedrock and associated ecological sites.

Table 2. Representative physiographic features

Geomorphic position, mountains	(1) Upper third of mountainflank
Landforms	(1) Mountains
Elevation	762–2,438 m
Slope	30–60%
Water table depth	203 cm
Aspect	Aspect is not a significant factor

Climatic features

Climate data and analyses are derived from 30-year averages gathered from National Oceanic and Atmospheric Administration (NOAA) weather stations contained within the range of an ecological site. However, no weather stations are available for this ecological site. The following information is a general climate description of the MLRA.

The Southern Alaska Coastal Mountains falls into two Köppen-Geiger climate classifications (Peel et al. 2007): tundra climate (ET) dominates the majority of the MLRA with small portions falling into the subarctic with cool summers and year around rainfall climate (Dfc). In the tundra climate, average temperatures are below 50°F for all months of the year, while the subarctic climate can experience highs above 50°F. Precipitation does not differ significantly across the seasons, but due to the high latitude environment solar radiation extremes occur with seasonal variability. The soil temperature regime of MLRA 222 is classified as cryic, where the mean annual soil temperature is between 32°F and 46°F (USDA-NRCS 2006).

Temperature and precipitation are affected by latitude, elevation, and proximity to maritime or continental zones. The average annual temperature and length of freeze-free period are not known. At the higher elevations, freezing temperatures are likely to occur during any month of the year. Most of the precipitation occurs as snowfall with rainfall increasing in importance in the southeast. Average annual precipitation is 120 to 200 inches but can be 250 inches or more at the highest elevations. Average annual snowfall ranges from about 200 to 800 inches. The snowfall greatly exceeds the annual snowmelt in many places, as evidenced by the abundance and extent of glaciers and ice fields (USDA-NRCS 2006).

Influencing water features

Alpine Dwarf Scrub Gravelly Slopes, Bedrock ecological sites are not influenced by wetland or riparian water features. Precipitation is the main source of water for this ecological site. Infiltration is very slow (Hydrologic Group D), and surface runoff is high. Surface runoff contributes some water to downslope ecological sites.




Figure 3. Hydrologic cycling in Alpine Dwarf Scrub Gravelly Slopes, Bedrock ecological site.

Soil features

Soils of this ecological site are in the Inceptisols order, further classified as Lithic Humicrypts with very slow infiltration and high runoff potential. The soil series, D22-Alpine Herbaceous Gravelly Diorite Slopes, is the most dominant soil series associated with this site. Parent material is colluvium over weathered residuum. The soils are well-drained and shallow, with a soil pH class ranging from extremely acid to strongly acid. A shallow depth to bedrock is noted as a rooting restriction for the soils of this ecological site.



Figure 4. Profile of soil series associated with Alpine Dwarf Scrub Gravelly Slopes, Bedrock.

Table 3. Representative soil features

Parent material	(1) Colluvium
Surface texture	(1) Gravelly loam
Drainage class	Well drained
Depth to restrictive layer	33 cm
Soil depth	33 cm
Surface fragment cover <=3"	0–2%
Surface fragment cover >3"	0–15%
Available water capacity (Depth not specified)	5.08 cm
Electrical conductivity (Depth not specified)	0–2 mmhos/cm

Soil reaction (1:1 water) (Depth not specified)	3.5–5.3
Subsurface fragment volume <=3" (Depth not specified)	0–20%
Subsurface fragment volume >3" (Depth not specified)	0–30%

Ecological dynamics

The information in this Ecological Dynamics section, including the state-and-transition model (STM), was developed based on historical data, current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.


The MLRA lies within the true alpine zone where glaciers are the dominant land cover. The non-glaciated areas are inhabited by a vegetative matrix resulting from a complex interaction among elevation, varying microclimates resulting from landscape topography, and natural disturbance regimes. The result is a heterogeneous landscape of ericaceous dwarf shrubs, low shrubs, and tall shrubs. Alpine Dwarf Scrub Gravelly Slopes, Bedrock ecological sites form an aspect of this vegetative continuum. This ecological site occurs on mountain backslopes on well-drained soils. Species characteristic of this ecological site consist of a sparse mix of dwarf ericaceous shrubs and herbaceous vegetation.

Located in the alpine life zone, the Alpine Dwarf Scrub Gravelly Slopes, Bedrock ecological site is exposed to a variety of harsh environmental conditions that drive and maintain the plant community. It is typically under snow for most of the year leaving a short season for plants to grow and reproduce. When the site is snow-free, cold temperatures and high winds in exposed positions further contribute to inhibited plant growth and performance.

The state-and-transition model that follows provides a detailed description of each state, community phase, pathway, and transition. This model is based on available experimental research, field observations, literature reviews, professional consensus, and interpretations.

State and transition model

Legend
No known state transitions or phase shifts



State 1

STATE 1 - REFERENCE STATE

The reference plant community is categorized as a sparse dwarf scrub-herbaceous vegetative community interspersed with bedrock and surface rock. The one community phase within the reference state is maintained by exposure to cold temperatures, wind, and a short growing season.

Community 1.1

Bog blueberry-Black crowberry/Star reindeer lichen-bedrock



Figure 5. Alpine Dwarf Scrub Gravelly Slopes, Bedrock ecological site at Skagway-Klondike Gold Rush National Historical Park.

This community phase is characterized by dwarf scrub and herbaceous vegetation interspersed with exposed bedrock and surface rock fragments. Bedrock cover is 10 to 70 percent, and surface rock fragment cover is as much as 50 percent. The plant community is dominantly dwarf scrub species such as bog blueberry, black crowberry, yellow mountainheath (*Phyllodoce glanduliflora* (Hook.) Coville), western moss heather (*Cassiope mertensiana* (Bong.) G. Don), alpine azalea (*Loiseleuria procumbens* (L.) Desv.), lingonberry (*Vaccinium vitis-idaea* L.), partridgefoot (*Luetkea pectinata* (Pursh) Kuntze), and arctic willow (*Salix arctica* Pall.). Dwarf shrub cover is generally 35 to 45 percent. Lichen cover is 5 to 60 percent. Common lichens include star reindeer lichen, greygreen reindeer lichen (*Cladina rangiferina* (L.) Nyl.), reindeer lichen (*Cladina arbuscular* (Wallr.) Hale & W.L. Culb.), cup lichen (*Cladonia squamosa* (Scop.) Hoffm. and *Cladonia gracilis* (L.) Willd.), snow lichen (*Stereocaulon* Hoffm.), and felt lichen (*Peltigera* Willd.). Forb cover is generally less than 10 percent with species such as redstem saxifrage (*Saxifraga lyallii* Engl.), dwarf fireweed (*Chamerion latifolium* (L.) Holub), russethair saxifrage (*Saxifraga ferruginea* Graham), Eschscholtz's buttercup (*Ranunculus eschscholtzii* Schldl.), groundcedar (*Lycopodium complanatum* L.), stiff clubmoss (*Lycopodium annotinum* L.), western oak fern (*Gymnocarpium Dryopteris* (L.) Newman), arctic starflower (*Trientalis europaea* L.), and arctic sweet coltsfoot (*Petasites frigidus* (L.) Fr.). Moss cover is minimal, but it may be as much as 30 percent in areas where lichen cover is minimal. Moss species include sphagnum (*Sphagnum capillifolium* (Ehrh.) Hedw.), splendid feather moss (*Hylocomium splendens* (Hedw.) Schimp.), Schreber's big red stem moss (*Pleurozium schreberi* (Brid.) Mitt.), dicranum moss (*Dicranum scoparium* Hedw.), polytrichum moss (*Polytrichum commune* Hedw.), and goose neck moss (*Rhytidiadelphus* (Lindb. ex Limpr.) Warnst.).

Additional community tables

Animal community

Vegetation in this ecological site does not seem to be commonly browsed and grazed as few of the ericaceous dwarf shrubs are palatable. However, while the site may not be an important foraging ground for wildlife, game trails may be abundant. Large mammals, such as Sitka deer (*Odocoileus hemionus sitkensis*), mountain goats (*Oreamnos americanus*), bears (*Ursus* sp.), wolves (*Canis lupus*), and wolverines (*Gulo gulo*), are known to utilize the associated mountain ridges of this site for migration and hunting. A limited number of bird species – e.g., rock ptarmigan (*Lagopus mutus*) and American pipit (*Anthus rubescens*) – can successfully nest in this harsh environment (Carsten 2007).

Other references

- Bailey, R.G. 1995. Ecoregions of North America. U.S. Department of Agriculture, Forest Service, Washington, DC, map scale 1: 15,000,000. Available at <https://www.fs.fed.us/rm/ecoregions/products/map-ecoregions-north-america/>.
- Boggs, K., S.C. Klein, J. Grunblatt, T. Boucher, B. Koltun, M. Sturdy, and G.P. Streveler. 2010. Alpine and subalpine vegetation chronosequences following deglaciation in coastal Alaska. *Arctic, Antarctic, and Alpine Research* 42: 385-395.
- Boggs, K., L. Flagstad, T. Boucher, T. Kuo, M. Aisu, J. Tande, and J. Michaelson. 2016. Vegetation Map and Classification: Southern Alaska and Aleutian Islands. Alaska Natural Heritage Program, Alaska Center for Conservation Science, University of Alaska Anchorage. 90 pps.
- Carsten, R. 2007. Chapter 5.2 Terrestrial habitats of Southeast Alaska. In: Schoen, J.W. and E. Dovichin (eds). *The Coastal Forests and Mountains Ecoregion of Southeastern Alaska and the Tongass National Forest: A Conservation Assessment and Resource Synthesis*. Audubon Alaska and The Nature Conservancy, Anchorage, AK.
- Gallant, A.L., E.F. Binnian, J.M. Omernick, and M.B. Shasby. 2010. Level III Ecoregions of Alaska. Corvallis, OR, U.S. EPA, National Health and Environmental Effects Research Laboratory, map scale 1: 5,000,000. Available at <http://http://www.epa.gov/eco-research//ecoregion-download-files-state-region-10>. (Accessed 11 September 2018).
- Kauffman, D.S., N.E. Young, J.P. Briner, and W.F. Manley. 2011. Alaska Palaeo-Glacier Atlas (Version 2), pps. 427-445. In: Ehlers, J., P.L. Gibbard, and P.D. Hughes (eds.). *Developments in Quaternary Science, Volume 15*. Amsterdam, The Netherlands.
- LANDFIRE. 2009. Biophysical Setting 7816430 Alaskan Pacific Maritime Alpine Herbaceous Dwarf Shrubland. In: *LANDFIRE National Vegetation Dynamics Models*. USDA Forest Service and US Department of Interior. Washington, DC.
- NatureServe. 2018. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1 NatureServe, Arlington, VA. Available at <http://explorer.natureserve.org>. (Accessed 10 September 2018).
- Peel, M.C., B.L. Finlayson, and T.A. McMahon. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11: 1633-1644.
- United States Department of Agriculture – Natural Resources Conservation Service (USDA-NRCS). 2006. *Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin*. U.S. Department of Agriculture Handbook 296. 682 pps.
- Viereck, L.A., C.T. Dyrness, A.R. Batten, and K.J. Wenzlick. 1992. *The Alaska Vegetation Classification*. General Technical Report PNW-GTR-286. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 278 pps.
- Wahrhaftig, C. 1965. *Physiographic Divisions of Alaska*. Geological Survey Professional paper 482. U.S. Department of the Interior, Geological Survey, U.S. Government Printing Office, Washington, DC. 52 pps.

Approval

Michael Margo, 6/20/2019

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community

cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-