

Ecological site R224XY500AK Loamy Wet Flood Plains Boreal-riparian scrub loamy wet flood plains

Accessed: 05/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

Table 2. Representative physiographic features

Landforms	(1) Flood plain
Flooding duration	Long (7 to 30 days)
Flooding frequency	Rare to occasional
Ponding duration	Brief (2 to 7 days) to long (7 to 30 days)
Ponding frequency	Frequent
Elevation	61–487 m
Slope	0–2%
Water table depth	0–25 cm

Climatic features

Influencing water features

Soil features

Boreal-Riparian Scrub Loamy Wet Flood Plains component is on channels on flood plains. The parent material consists of sandy and silty alluvium over sandy and gravelly alluvium. The depth to restrictive layer is 13 to 31 inches to strongly contrasting textural stratification. It is very poorly drained. The slowest permeability of the soil material is moderately slow. Available water capacity is low and shrink swell potential is low. This soil is occasionally flooded and is frequently ponded. The top of the seasonal high water table is at 0 inches. There are no saline horizons within 30 inches of the soil surface.

Table 3. Representative soil features

Surface texture	(1) Fine sand (2) Silt
Family particle size	(1) Sandy
Drainage class	Poorly drained
Permeability class	Moderately slow
Soil depth	33–104 cm
Surface fragment cover <=3"	0–5%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	10.16 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0–1 mmhos/cm
Soil reaction (1:1 water) (0-101.6cm)	6.2–6.9
Subsurface fragment volume <=3" (Depth not specified)	20–65%
Subsurface fragment volume >3" (Depth not specified)	0–25%

Ecological dynamics

State and transition model

Contributors

Dennis Moore Michelle Schuman

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be

known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators		
1.	Number and extent of rills:	
2.	Presence of water flow patterns:	
3.	Number and height of erosional pedestals or terracettes:	
4.	Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):	
5.	Number of gullies and erosion associated with gullies:	
6.	Extent of wind scoured, blowouts and/or depositional areas:	
7.	Amount of litter movement (describe size and distance expected to travel):	
8.	Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):	
9.	Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):	
10.	Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:	

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be

	mistaken for compaction on this site):
12.	Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):
	Dominant:
	Sub-dominant:
	Other:
	Additional:
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
14.	Average percent litter cover (%) and depth (in):
15.	Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):
16.	Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
17.	Perennial plant reproductive capability: