

Ecological site R225XY363AK

Southern Alaska Scrub Loamy Hill Backslope

Last updated: 6/14/2025
Accessed: 03/10/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 225X–Southern Alaska Peninsula Mountains

Major Land Resource Area 225X (Southern Alaska Peninsula Mountains), herein called area, is in the Southern Alaska Land Resource Region (LRR). MLRA 225X covers approximately 23,935 square miles of the southern parts of the Kodiak Archipelago and the slopes of the southern Aleutian Mountains on the Alaska Peninsula. The landscape is comprised of rugged mountains separated by narrow valleys, with outwash plains and low hills towards the coast. Glaciers and ice fields cover the highest mountains. Flood plains are predominantly restricted to broad river valleys. Elevations range from sea level to 9,372 feet at the summit of Shishaldin Volcano. The soils and vegetation in this MLRA have been influenced by ash deposits from nearby Mount Katmai and surrounding volcanoes. This MLRA is primarily comprised of wilderness, with towns and villages primarily located along rivers, lakes, and the coast. Small villages are scattered along the coastline and include King Cove, Old Harbor, Karluk, and Larsen Bay among others. There is no road access to MLRA 225X from Anchorage, AK, and access is primarily via plane or boat.

MLRA 225X, excluding high peaks and steep upper backslopes, was glaciated during the Late Pleistocene. Glacial deposits were eroded or covered by colluvium or alluvium during the Holocene, which comprises 60 percent of the current landscape. Modified glacial moraines and outwash landforms are prevalent. Volcanic activity continues through the present day, and ash deposits are represented in many of the soils of this MLRA.

The dominant soil orders are Andisols, Histosols, and Inceptisols. Soils have a cryic temperature regime or subgelic soil temperature class, an aquic or udic soil moisture regime, and primarily amorphous mineralogy. Permafrost is sporadic in the Southern Alaska

LRR. The primary soils in this MLRA developed from volcanic ash over colluvium or from thick organic material. Miscellaneous (non-soil) areas comprise greater than 50 percent of all acreage in this MLRA and includes “rock outcrops, rubble land, glaciers, riverwash, and beaches” (USDA-NRCS, 2022).

The climate in this MLRA is shaped by maritime influences from Bristol Bay to the west and the Pacific Ocean to the south. Mountains effect local patterns in temperature and precipitation. Temperatures are typically cool throughout the year. The mean annual temperature at sea level is 37 to 43 degrees Fahrenheit. Precipitation ranges from 30 inches along the coast to over 100 inches at high elevations. Snowfall ranges from 50 to 200 inches and glaciers and icefields are present at higher elevations.

This MLRA is dominated by tall alder and willow shrubs at lower elevations. Vegetation shifts to low and then dwarf shrubs at increased elevations. Herbaceous communities are on exposed plains and hills, and sedges dominate wet depressions. Balsam poplar forests are restricted to flood plains and warm, low mountain slopes in the northern parts of the MLRA (USDA-NRCS, 2022).

LRU notes

MLRA 225X supports three life zones delineated by the physiological limits of plant communities along longitudinal and elevational gradients: Aleutian, South Alaska maritime, and South Alaska alpine. The Aleutian climate covers hills of the southern Kodiak archipelago and on the extended Alaska Peninsula. These regions are low-lying, exposed and are scoured by winter winds. Vegetation is primarily low shrublands, heathlands and often diverse herbaceous meadows. The South Alaska maritime climate is common in mountainous areas where local site conditions delineate alpine and lowland areas. Certain vascular plant species are common in the lowlands and much less common in the alpine (i.e. *Populus balsamifera*, *Alnus* spp., *Salix pulchra*, *Betula nana*, *Ledum palustre* ssp. *decumbens*, and *Calamagrostis canadensis*). The alpine generally occurs at elevations above 1,500 feet, though may extend higher on warm, north-facing slopes, and lower on cooler slopes. Vascular plants are restricted in height and often exclude common lowland species. The transition between South Alaska maritime and alpine vegetation can occur within a range of elevations, and is highly dependent on latitude, slope, aspect, and shading from adjacent mountains.

Classification relationships

Alaska Vegetation Classification:

Open low scrub (II.C.2 – level III) / Willow (II.C.2.g - level IV)

(Viereck et al., 1992)

Circumboreal Vegetation Map – Alaska-Yukon Region: Aleutian Heath and Meadow (Jorgensen and Meidinger, 2015)

LANDFIRE BioPhysical Settings: 7616400 – Aleutian Mesic-Wet Willow Shrubland (Landfire, 2009)

Kodiak Archipelago Land Cover Classification:

Best fit: Class 25 – heath hummocks with forbs (Fleming and Spencer, 2007)

Ecological site concept

Ecological Site characteristics:

- Supports an open willow shrubland in the reference plant community
- Occurs on hill slopes
- Soils are moderately deep (33 to 50 inches) to a root-limiting layer
- No ponding or flooding, and no water table during the growing season

Associated sites

F225XY315AK	Southern Alaska Forest Loamy Terraces Forested alluvial terraces and footslopes
F225XY360AK	Southern Alaska Forest Loamy Hill Backslopes Forested mountain backslopes downslope of this site
F225XY310AK	Southern Alaska Riparian Complex Flood Plains Forested flood plain complex on river valley bottoms

Similar sites

R225XY333AK	Southern Alaska Scrub Steep Backslopes Birch-ericaceous scrubland on steeper backslopes
R225XY334AK	Southern Alaska Scrub Shallow Ashy Backslopes Ericaceous heath dwarf scrubland on shallower soils on hill slopes

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Salix</i> (2) <i>Vaccinium uliginosum</i>
Herbaceous	(1) <i>Lupinus nootkatensis</i> (2) <i>Chamerion angustifolium</i>

Physiographic features

This ecological site occurs on hill backslopes across plain landscapes. Slopes are linear, with gentle to strongly slopes gradients. Flooding and ponding do not occur and there is no water table in the soil profile. Aspect is not a factor.

Table 2. Representative physiographic features

Slope shape across	(1) Linear
Slope shape up-down	(1) Linear
Hillslope profile	(1) Backslope
Geomorphic position, hills	(1) Side Slope
Landforms	(1) Plains > Hill
Runoff class	Low to medium
Flooding frequency	None
Ponding frequency	None
Elevation	30–305 m
Slope	5–20%
Water table depth	152 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Negligible to high
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	3–457 m
Slope	0–30%
Water table depth	Not specified

Climatic features

The climate in this MLRA is shaped by the maritime influences of Bristol Bay to the west and the Pacific Ocean to the south. Cloudy days are the norm. Temperature and precipitation are patterned around mountainous effects. Temperatures are typically cool throughout the year. The mean annual temperature at sea level is between 37 and 43 degrees Fahrenheit and generally decreases as elevation increases. Precipitation ranges from 30 inches along the coast to over 100 inches at high elevations. Snowfall ranges from 50 to 200 inches and supports glaciers and icefields at higher elevations (USDA-NRCS, 2022). Frost-free and freeze-free periods in the city of Kodiak are presented in the table below (WRCC, 2024). These periods shorten as elevation increases, culminating in the shortest frost-free and freeze-free periods in the alpine.

Table 4. Representative climatic features

Frost-free period (characteristic range)	124-156 days
Freeze-free period (characteristic range)	164-195 days
Precipitation total (characteristic range)	762-1,930 mm
Frost-free period (actual range)	121-167 days
Freeze-free period (actual range)	153-199 days
Precipitation total (actual range)	356-2,540 mm
Frost-free period (average)	140 days
Freeze-free period (average)	173 days
Precipitation total (average)	1,346 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Influencing water features

This ecological site is not associated with or influenced by streams or wetlands. Precipitation and throughflow are the main source of water for this ecological site. Surface runoff and throughflow contribute water to downslope ecological sites.

Wetland description

This ecological site is not associated with wetlands.

Soil features

Soils are cold, moderately deep Andisols (Soil Survey Staff, 2013). Parent material is ash over till over paralithic residuum. Permafrost is absent. Surface rock fragment cover is low to absent. There is little to no organic soil horizon. Soil pH is very strongly to strongly acidic. Soils are dry during the growing season.

Correlated soil components in MLRA 225X: E25-Maritime scrub/herbaceous-ashy slopes

Table 5. Representative soil features

Parent material	(1) Residuum
Surface texture	(1) Medial silt loam
Drainage class	Well drained
Permeability class	Moderate
Depth to restrictive layer	84–127 cm
Soil depth	84–127 cm
Surface fragment cover ≤3"	0–6%
Surface fragment cover >3"	0–6%

Available water capacity (0-25.4cm)	5.33–6.86 cm
Soil reaction (1:1 water) (0-25.4cm)	4.5–5.5
Subsurface fragment volume ≤3" (Depth not specified)	17–28%
Subsurface fragment volume >3" (Depth not specified)	3–7%

Ecological dynamics

The open willow shrubland reference plant community occurs on moderately deep soils that are root restricted by paralithic bedrock between 33 and 50 inches. Larger shrubs and trees are generally excluded at low elevations. This ecological site grades into an alpine scrubland at higher elevations, and a tall shrubland or boreal forest at lower elevations.

The reference plant community (1.1) is unlikely to shift to a new plant community due to natural disturbance (Viereck et al., 1992; Landfire, 2009). However, vegetation can be patchy and shrubs and herbaceous species may form a mosaic across the slope. Fuel load is low and cool temperatures generally inhibit fire. Root systems stabilize the slope against mass erosion. Some striped sorting of surface fragments and vegetation may be evident (Landfire, 2009).

The information in this Ecological Dynamics section, including the state-and-transition model (STM), was developed based on current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

State and transition model

Ecosystem states

1. Reference State

State 1 submodel, plant communities

1.1. Lingonberry -
black crowberry /
sedge / lupine -
fireweed

State 1 Reference State



Figure 7. Open willow scrubland hosts a diversity of herbaceous species in unshaded areas.

The reference state supports one community phase. The reference plant community is a stable open willow scrubland community. This community supports a mix of low and dwarf shrubs. Willow is common, but total canopy cover is site specific and can range from 10 – 75 percent. Areas with low willow cover commonly support a mix of shrubs such as bog blueberry, black crowberry, and dwarf Labrador tea. Herbaceous cover varies. All community phases in this report are characterized using the Alaska vegetation classification system (Vioreck et al., 1992).

Dominant plant species

- willow (*Salix*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- sedge (*Carex*), grass
- bluejoint (*Calamagrostis canadensis*), grass
- Nootka lupine (*Lupinus nootkatensis*), other herbaceous
- fireweed (*Chamerion angustifolium*), other herbaceous

Community 1.1

Lingonberry - black crowberry / sedge / lupine - fireweed



Figure 8. Open willow scrubland hosts a diversity of herbaceous species in unshaded areas.

The reference plant community is an open to closed willow scrubland. The major plant groups are low shrubs, dwarf shrubs, medium graminoids, and medium forbs. Willow is common, but total canopy cover is site specific and can range from 10 – 75 percent. Non-willow areas support a mix of shrubs, including lingonberry, black crowberry, bog blueberry, alpine azalea, and dwarf Labrador tea. Herbaceous cover varies from area to area. Graminoids are usually present as sedges. Ground cover is comprised of mosses, lichens, and herbaceous litter. This community may present as a mosaic in ecotonal areas where it abuts with alder stands and alpine shrubland communities.

Dominant plant species

- willow (*Salix*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- sedge (*Carex*), grass
- bluejoint (*Calamagrostis canadensis*), grass
- Nootka lupine (*Lupinus nootkatensis*), other herbaceous
- fireweed (*Chamerion angustifolium*), other herbaceous

Additional community tables

Inventory data references

Vegetative communities and transitions are described using existing models and expert knowledge. There are no vegetation inventory data points in NASIS associated with this

ecological site.

External data sources:

The Alaska Vegetation Classification (Vioreck et al., 1992)

The Alaska-Yukon Region of the Circumboreal Vegetation Map (CBVM) (Jorgensen and Meidinger, 2015)

LANDFIRE Biophysical Settings Models (Landfire, 2009)

References

Vioreck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Other references

Fleming, M.D., and Spencer, P. 2007. Kodiak Archipelago Land Cover Classification User Guide. Available at https://akevt.gi.alaska.edu/data/Kodiak_UsersGuide_v1.1.pdf. Accessed April 24, 2024.

Jorgensen, T., and D. Meidinger. 2015. The Alaska Yukon Region of the Circumboreal Vegetation Map (CBVM). CAFF Strategies Series Report. Conservation of Arctic Flora and Fauna, Akureyri, Iceland. ISBN: 978-9935-431-48-6.

Landfire. 2009. Biophysical Setting 7616400 – Aleutian Mesic-Wet Willow Shrubland. In: LANDFIRE Biophysical Setting Model: Map zone 76, [Online]. In: Vegetation Dynamics Models. In: LANDFIRE. Washington, DC: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory; U.S. Geological Survey; Arlington, VA: The Nature Conservancy (Producers). Available: https://www.landfire.gov/national_veg_models_op2.php. Accessed Mar 8, 2024

PRISM Climate Group. 2018. “Alaska – average monthly and annual precipitation and minimum, maximum, and mean temperature for the period 1981-2010.” Oregon State University, Corvallis, Oregon. <https://prism.oregonstate.edu/projects/alaska.php>. Accessed Sept 17, 2024.

Scenarios network for Alaska and arctic planning (SNAP). 2014. “Historical Monthly Temperature – 1km, 1901-2009”. <http://ckan.snap.uaf.edu/dataset/>. Accessed Sept 17, 2024.

Scenarios network for Alaska and arctic planning (SNAP). 2014. “Historical monthly and derived precipitation products downscaled from CRU TS data via the delta methods – 2km, 1901-2009”. <http://ckan.snap.uaf.edu/dataset/>. Accessed Sept 17, 2024.

Soil Survey Staff, 2013. Simplified Guide to Soil Taxonomy. USDA-Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

United States Department of Agriculture, Natural Resources Conservation Service. 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture, Agriculture Handbook 296.

Western Regional Climate Center. 2024. "Kodiak WSO Airport, Alaska 'Freeze Free' Season Probabilities." <https://wrcc.dri.edu/cgi-bin/cliTFrezD.pl?akkodi>. Accessed Sept 19, 2024.

Contributors

Phil Barber
Blaine Spellman
Marji Patz

Acknowledgments

This soil – ecological site correlation was reviewed by a workshop team during a February 2024 PES workshop in Wasilla, AK.

This ecological site description (ESD) fulfills the requirements of the Provisional Ecological Site (PES) national initiative. This ESD is published to fit current site-soil correlations as they are currently mapped and understood. Further data collection may provide the information to update this ESD from the provisional level to the approved level.

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	03/10/2026
Approved by	Blaine Spellman

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. Perennial plant reproductive capability:
