

# Ecological site R226XY032AK Crowberry (Upland) (Umnak Island)

Accessed: 05/02/2024

## General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

## Physiographic features

Occurs on shoulders and back-slopes of ridges associated with volcanic cones and on very steep side-slopes of ridges associated with volcanic cones.


**Table 2. Representative physiographic features**

Landforms	(1) Hillside
Elevation	120–300 ft
Slope	0–120%

## Climatic features

**Table 3. Representative climatic features**

Frost-free period (average)	120 days
Freeze-free period (average)	100 days
Precipitation total (average)	24 in



**Figure 1. Monthly precipitation range**




Figure 2. Monthly average minimum and maximum temperature

## Influencing water features

### Soil features

Soils are moderately deep and well drained. Surface textures are gravelly silt loam and overlie gravel-sized scoria. Soil pH is moderately acid. Runoff is very low and permeability is rapid to very rapid.

Table 4. Representative soil features

Surface texture	(1) Gravelly silt loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Rapid to very rapid
Soil depth	20–40 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	2.5–2.7 in
Calcium carbonate equivalent (0-40in)	0%
Electrical conductivity (0-40in)	0 mmhos/cm
Sodium adsorption ratio (0-40in)	0
Soil reaction (1:1 water) (0-40in)	5.6–6
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

## Ecological dynamics

### State and transition model

## Ecosystem states

1. Crowberry
--------------

## State 1 submodel, plant communities

1.1. Crowberry
----------------

## State 1 Crowberry

## Community 1.1 Crowberry

Shrubs make up about 75% of the composition, sedges and grasses 20% and forbs about 5% of the composition. Total annual vascular herbage production is 840 pounds/acre. Total live lichen biomass is 3,000 pounds/acre. Moss biomass is high and suppresses production of this site.

## Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
<b>Shrub/Vine</b>					
1				600–650	
	black crowberry	EMNI	<i>Empetrum nigrum</i>	600–650	–
<b>Grass/Grasslike</b>					
1				150–170	
	longawn sedge	CAMA11	<i>Carex macrochaeta</i>	150–170	–
	Bering hairgrass	DEBR2	<i>Deschampsia brevifolia</i>	5–15	–
	bluegrass	POA	<i>Poa</i>	10–15	–
	wideleaf polargrass	ARLA2	<i>Arctagrostis latifolia</i>	0	–
<b>Forb</b>					
1				20–30	
	Nootka lupine	LUNO	<i>Lupinus nootkatensis</i>	20–25	–
	cloudberry	RUCH	<i>Rubus chamaemorus</i>	0–5	–
	saxifrage	SAXIF	<i>Saxifraga</i>	0	–
	seacoast angelica	ANLU	<i>Angelica lucida</i>	0	–
	mountain harebell	CALA7	<i>Campanula lasiocarpa</i>	0	–
<b>Lichen</b>					
1				20–30	
	whiteworm lichen	THAMN3	<i>Thamnolia</i>	25–30	–
	cup lichen	CLAM60	<i>Cladonia amaurocraea</i>	0–5	–
	reindeer lichen	CLAR60	<i>Cladina arbuscula</i>	0–5	–
	greengreen reindeer lichen	CLRA60	<i>Cladina rangiferina</i>	0–5	–

## Animal community

Because of the high subsistence value and use of black crowberry berries, this site should only be lightly grazed. Reindeer have a tendency to concentrate on this site because they are attracted to the lichens during winter.

## Recreational uses

Traditional Mossberry picking, hiking

## Contributors

Swanson

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	

Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 

14. **Average percent litter cover (%) and depth ( in):**
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 

17. **Perennial plant reproductive capability:**
-