

## Ecological site R226XY053AK Wet Meadow Complex (AK653-St Paul Island)

Accessed: 07/17/2024

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

### Physiographic features

This site occurs in nearly flats adjacent to the coast in the proximity of laggons and salt marshes. This site is subject to periodic tidal inundation.


**Table 2. Representative physiographic features**

Landforms	(1) Tidal flat
Flooding duration	Very brief (4 to 48 hours) to brief (2 to 7 days)
Flooding frequency	Rare to occasional
Elevation	0–2 m
Slope	0–3%


### Climatic features

**Table 3. Representative climatic features**

Frost-free period (average)	120 days
Freeze-free period (average)	100 days
Precipitation total (average)	610 mm



**Figure 1. Monthly precipitation range**



**Figure 2. Monthly average minimum and maximum temperature**

## Influencing water features

### Soil features

Soils are very deep and poorly or very poorly drained. Soils can have a thick organic layer or sand at the surface. Soil pH is neutral. Runoff is very low to negligible and permeability is rapid.

**Table 4. Representative soil features**

Surface texture	(1) Mucky sand
Family particle size	(1) Sandy
Drainage class	Poorly drained
Permeability class	Rapid
Soil depth	152–165 cm
Surface fragment cover ≤3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	15.75–16.26 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	6.6–7.3

Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

## Ecological dynamics

### State and transition model

#### Ecosystem states

1. Carex spp./Arctagrostis latifolia
--

#### State 1 submodel, plant communities

1.1. Carex spp./Arctagrostis latifolia
--

### State 1

#### Carex spp./Arctagrostis latifolia

#### Community 1.1

#### Carex spp./Arctagrostis latifolia

Sedges and grasses make up 95% and forbs 5% of the composition. Total annual vascular herbage production is 1970 pounds/acre.

### Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1				2046–2186	
	sedge	CAREX	<i>Carex</i>	1569–1793	–
	wideleaf polargrass	ARLA2	<i>Arctagrostis latifolia</i>	213–235	–
	American dunegrass	LEMOM2	<i>Leymus mollis ssp. mollis</i>	196–207	–
<b>Forb</b>					
1				101–106	
	starwort	STELL	<i>Stellaria</i>	90–95	–
	Aleutian violet	VILA6	<i>Viola langsdorffii</i>	0–6	–
	Pacific hemlockparsley	COGM	<i>Conioselinum gmelinii</i>	0–6	–

### Animal community

Sedges provide high value reindeer forage during winter when covered by snow. Wide leaf polargrass maintains its forage value well into the winter and provides excellent early spring forage. Lyme grass is seldom selected by

reindeer during spring and summer and is of no value during winter.

## Contributors

Swanson

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---