

Ecological site R226XY061AK Lake Margin (AK653 St Paul Island)

Accessed: 05/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occupies fringe areas around freshwater lakes.

Table 2. Representative physiographic features

Landforms	(1) Lakeshore
Elevation	3–9 m
Slope	0–3%

Climatic features

Table 3. Representative climatic features

Frost-free period (average)	120 days
Freeze-free period (average)	100 days
Precipitation total (average)	610 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly average minimum and maximum temperature

Influencing water features

Soil features

Soils are very deep and poorly tovery poorly drained. Textures are coarse and can be high in organic matter content; soil pH is moderately acid. Runoff is negligible and permeability is moderately slow to rapid.

Table 4. Representative soil features

Surface texture	(1) Mucky sand
Family particle size	(1) Sandy
Drainage class	Poorly drained
Permeability class	Moderately slow to rapid
Soil depth	152–165 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	13.46–13.97 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	5.6–6
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

State and transition model

Ecosystem states

Deschampsia beringensis/Juncus arcticus

State 1 submodel, plant communities

1.1. Deschampsia beringensis/Juncus arcticus

State 1

Deschampsia beringensis/Juncus arcticus

Community 1.1

Deschampsia beringensis/Juncus arcticus

Sedges and grasses make up about 98% of the composition. Forbs make up 2%. Total annual vascular herbage production is 550 pounds/acre.

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass	/Grasslike				
1				588–644	
	Bering's tufted hairgrass	DEBE2	Deschampsia beringensis	219–230	_
	arctic rush	JUAR2	Juncus arcticus	213–224	_
	sedge	CAREX	Carex	106–118	_
	bluegrass	POA	Poa	50–62	_
Forb	Forb				
1				0–6	
	boreal yarrow	ACMIB	Achillea millefolium var. borealis	0–1	_
	seacoast angelica	ANLU	Angelica lucida	0–1	_
	Tilesius' wormwood	ARTI	Artemisia tilesii	0–1	_
	Nootka lupine	LUNO	Lupinus nootkatensis	0–1	_

Contributors

Swanson

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author	r(s)/participant(s)		
Conta	ct for lead author		1
Date			1
Appro	ved by		1
Appro	val date		1
Comp	osition (Indicators 10 and 12) based on	Annual Production	
ndica	ators		
1. Nu	mber and extent of rills:		
2. Pr	esence of water flow patterns:		
3. Nu	mber and height of erosional pedesta	als or terracettes:	
	re ground from Ecological Site Descr re ground):	iption or other stud	dies (rock, litter, lichen, moss, plant canopy are not
 5. Nu	mber of gullies and erosion associate	ed with gullies:	
6. Ex	tent of wind scoured, blowouts and/o	r depositional areas	as:
7. A n	nount of litter movement (describe siz	ze and distance exp	pected to travel):
	il surface (top few mm) resistance to lues):	erosion (stability va	values are averages - most sites will show a range of
9. So	il surface structure and SOM content	(include type of str	tructure and A-horizon color and thickness):
	fect of community phase composition stribution on infiltration and runoff:	า (relative proportion	on of different functional groups) and spatial
	esence and thickness of compaction staken for compaction on this site):	layer (usually none;	e; describe soil profile features which may be

12.	Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):
	Dominant:
	Sub-dominant:
	Other:
	Additional:
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
14.	Average percent litter cover (%) and depth (in):
15.	Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):
16.	Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
17.	Perennial plant reproductive capability: