

Ecological site R228XY704AK

Dryas/dwarf ericaceous scrub gravelly slopes

Last updated: 6/12/2025

Accessed: 01/22/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 228X—Interior Alaska Mountains

Physiography

The Interior Alaska Mountains Major Land Resources Area (MLRA) (228X) includes portions of the high mountain slopes, hills, and plains of the Alaska Range, Talkeetna Mountains, Chugach Mountains, Wrangell Mountains and the northern Aleutian Range. This MLRA comprises 54,205 square miles and consists of rugged, high mountains and low, rounded hills and extended footslopes along the base of the mountains. Most of the area is undeveloped wildland and includes true alpine and subalpine life zones. Geology consists of sedimentary, metamorphic, and igneous bedrock. Climate is considered continental subarctic.

MLRA boundaries

MLRA 228X is expansive and therefore shares a boundary with many MLRAs. Boundaries with other mountainous MLRAs such as 222X (Southern Alaska Coastal mountains), 223X (Cook Inlet Mountains), and 225X (Southern Alaska Peninsula Mountains) result from orogenic differences (225X,223X) or variation in climate (222X). Other MLRAs such as 236X (Bristol-Bay-Northern Alaska Peninsula Lowlands), 227X (Copper River Basin), 229X (Interior Alaska Lowlands), and 230X (Yukon-Kuskokwim Highlands) have distinct physiographic boundaries where steep mountains meet lowlands, basins, and floodplains.

Waterways

Encompassed within the Pacific Mountain system, the mountains of MLRA 228X are dissected by high-gradient valleys with braided floodplains in the valley bottoms. Glaciers, snowfields, and ice fields make up 15 percent of the area and elevations range from about 1,500 feet in the Copper River Basin to 20,320 feet at the summit of Denali. The major

rivers of this MLRA include the Tanana, Kuskokwim, and Copper, and drain into the Bering Sea (Tanana, Kuskokwim), and the Gulf of Alaska (Copper). The headwaters of the Susitna River are part of this MLRA and drain into Cook Inlet through the Cook Inlet Mountains, and Cook inlet Lowlands (MLRA 223X and 224X, respectively). This MLRA is in the zone of discontinuous permafrost, where permafrost mostly occurs in fine-textured soils on gently sloping landforms and/or on northerly aspects.

Geology

Except for the highest peaks and upper ridges, all of this area was glaciated during the late Pleistocene. Glacial deposits have mostly eroded or have been buried by colluvium and alluvium throughout the Holocene, yet some highly modified glacial deposits remain at lower elevations on low mountain slopes and valleys. Loess also occurs at lower elevations, and most valley bottoms have been buried by recent alluvial deposits. Bedrock geology is comprised of sedimentary, metamorphic, igneous, and volcanic rock, and gold mining does take place in this MLRA.

Soils

The dominant soil orders in MLRA 228X are Gelisols, Inceptisols, Spodosols, and Entisols. The Gelisols are shallow or moderately deep to permafrost, occur on finer textured sediments, and are poorly drained or very poorly drained. Common Gelisol suborders are Histels, Orthels, and Turbels. The Histels have thick accumulations of surface organic material and occur in depressions and peat plateau. The Orthels and Turbels have comparably thinner surface organic material and occur on drainageways, stream terraces, and outwash plains. The Inceptisols, Spodosols, and Entisols lack permafrost in the soil profile. Spodosols are formed from weathering processes that strip organic matter combined with aluminum from upper horizons and deposit them into lower horizons. Entisols and Inceptisols are characteristically undeveloped, with Inceptisols exhibiting only moderate weathering and development while Entisols exhibit little to no evidence of development at all. Soils have a subgelic or cryic temperature regime with aquic or udic moisture regime and mixed mineralogy. Miscellaneous areas make up 58 percent of this MLRA and are classified as rock outcrop, rubble land, and glaciers.

Climate

The climate of this area is characterized by brief, cool summer, and long, cold winters, but extreme variation in elevation results in a wide range of climatic conditions. Average annual precipitation ranges from 15 to 20 inches at lower elevations to 100 inches at high elevations. Rainfall is generally highest in July, August, and early September. The average annual snowfall ranges from 70 to 400 inches, and the average annual temperature at Denali Park headquarters is 27 degrees F. Freeze-free period ranges from 50 to 80 days, but at higher elevations, freezing temperatures can occur at any point throughout the year.

Vegetation

The Interior Mountains MLRA is defined by subalpine and alpine life zones; therefore, true forested communities do not occur and are restricted to surrounding lowland MLRAs. Black and white spruce trees do occur in the subalpine zone, but are often sporadic, and

exhibit Krummholz (stunted and/or crooked) growth forms and do not produce viable seed. Subalpine vegetation is characterized by birch-willow scrublands or spruce-scrub woodlands on loamy, stable mountain slopes. Unstable, colluvial slopes are typically dominated by alder scrub communities which can, on occasion, include scattered black and white spruce. Willow typically dominates drainages, while wet, poorly to very poorly drained swales are comprised of tussock sedge-scrub species. Low birch-ericaceous scrub communities climb up mountain slopes until they are replaced by dryas-ericaceous dwarf scrub communities in the true alpine zone. Lichen also plays an important role in skeletal and bedrock-controlled high elevation mountain slopes, ridges, and summits. There is generally little to no plant growth at elevations above 7,500 feet (USDA, 2022).

LRU notes

This area supports two life zones defined by the physiological limits of plant communities along an elevational gradient: subalpine, and alpine. In this area, the boreal life zone occurs below 2500 feet elevation on average, and is relegated to surrounding, lowland MLRAs. The subalpine zone is typically a narrow transitional band between the boreal and the alpine life zones, and is characterized by sparse, stunted trees. In the subalpine, certain types of birch and willow shrub species grow at over one meter in height (commonly *Betula glandulosa* and *Salix pulchra*). In the alpine, trees no longer occur, and all shrubs are dwarf or lay prostrate on the ground. The transition between boreal and alpine vegetation can occur within a range of elevations, and is highly dependent on slope, aspect, and shading from adjacent mountains.

Within each life zone, there are plant assemblages that are typically associated with cold slopes and warm slopes. Cold slopes and warm slopes are created by the combination of the steepness of the slope, the aspect, and shading from surrounding ridges and mountains. Warm slope positions typically occur on southeast to west facing slopes that are moderate to very steep (over 10 percent slope) and are not shaded by the surrounding landscape. Cold slopes typically occur on northwest to east facing slopes, occur in shaded slope positions, or occur in low-lying areas that are cold air sinks. Examples of shaded positions include head slopes, low relief backslopes of hills, and the base of hills and mountains shaded by adjacent mountain peaks. These warm and cold slopes can shift the elevation at which life zones occur. Warm slopes can allow communities to persist at higher elevations, while cold slopes can restrict these same communities to lower elevations.

Classification relationships

Alaska Vegetation Classification
Dryas dwarf scrub tundra (II.D.1.a – Level IV)
(Viereck et al. 1992)

Circumboreal Vegetation Map – Alaska-Yukon Region:
Central-Northern Alaska-Yukon Alpine Dwarf Scrub and Meadows

(Jorgensen and Meidinger, 2015)

LANDFIRE Biophysical Settings

7416310 – Western North American Boreal Alpine Dwarf-Shrub Summit

(LANDFIRE biophysical settings, 2009)

Ecological site concept

- Ecological site R228XY704AK is an alpine dryas/dwarf ericaceous scrub community on mountain slopes, hillslopes, and glacial landforms
- Soils are well-drained and formed in organic material and silty loess over colluvium
- Elevations range from 2460 to 5250 feet
- Plant community is influenced by harsh weather conditions, short growing season, and persistent snowpack
- The representative plant community (1.1) is characterized by a dryas dwarf scrub tundra community

Associated sites

R228XY713AK	Alpine tussock tundra hummocks The alpine scrub earth hummocks ecological site occur adjacent to the dryas/ericaceous scrub gravelly slopes ecological on cold slopes where enough water is present for organic matter and ice lenses to form dome-like hummocks.
-------------	---

Similar sites

R228XY705AK	White spruce/alder scrub unstable slopes The white spruce/willow-birch ericaceous scrub loamy slopes ecological site occurs on similar landforms as the dryas/ericaceous scrub gravelly slopes, but occurs in the subalpine zone. As elevation increases, R228XY705AK will shift into an ecotonal community that resembles R228XY704AK as birch begins to lose dominance while ericaceous dwarf scrubs and dryas increase in cover.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Dryas octopetala</i> ssp. <i>octopetala</i> (2) <i>Vaccinium uliginosum</i>
Herbaceous	(1) <i>Cladina</i> (2) <i>Thuidium recognitum</i>

Physiographic features

Ecological site R228XY704AK occurs on mountain slopes and hillslopes as well as various

glacial landforms. Elevation generally ranges from 2460 to 5250 feet above sea level on slopes ranging from 2 to 50 percent on all aspects. Soils are considered well drained, and no water table has been detected on this site.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope (2) Mountains > Moraine (3) Mountains > Hillslope
Runoff class	Very low to medium
Flooding frequency	None
Ponding frequency	None
Elevation	750–1,600 m
Slope	2–50%
Water table depth	152 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	546–2,164 m
Slope	0–90%
Water table depth	Not specified

Climatic features

The climate of this high-elevation area is characterized by short growing season, cool summers, and long winters. Mean annual precipitation is around 15 inches at lower elevations but can reach much higher totals at higher elevations. June, July, and August are the wettest months of the year, while February, March, and April are the driest. On average, there are 17 frost free days per year, but at high elevations, freezing temperatures can occur any month of the year. The mean maximum temperature is 67 degrees Fahrenheit in July, while the mean low temperature is -9 degrees Fahrenheit in January. At higher elevations, this temperature range will be greatly skewed towards colder temperatures.

Table 4. Representative climatic features

Frost-free period (characteristic range)	5-30 days
--	-----------

Freeze-free period (characteristic range)	63-77 days
Precipitation total (characteristic range)	305-432 mm
Frost-free period (actual range)	1-41 days
Freeze-free period (actual range)	50-84 days
Precipitation total (actual range)	305-457 mm
Frost-free period (average)	17 days
Freeze-free period (average)	69 days
Precipitation total (average)	381 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) TONSINA [USC00509385], Copper Center, AK
- (2) NABESNA [USC00506147], Gakona, AK
- (3) PAXSON [USC00507097], Copper Center, AK
- (4) MCKINLEY PARK [USC00505778], Healy, AK
- (5) CANTWELL 2 E [USC00501243], Cantwell, AK
- (6) FAREWELL LAKE [USC00503009], Mc Grath, AK

Influencing water features

Due to topographic position, no wetlands or water features are associated with this site.

Soil features

The soils of ecological site R228XY704AK formed in organic material and loess over gravelly colluvium. Surface fragments are common, but do not exceed 25 percent, and surface textures are silt loams. The mineral soil is considered loamy-skeletal and forms in windblown silty eolian deposits and gravelly colluvium. Strongly contrasting textural stratification is a common restrictive layer at very shallow depth. Soil depth is controlled by lithic contact which typically occurs at deep and very deep depths. Soil pH ranges from acidic to basic depending on mineral composition of the parent material. Soils of the ecological site are well-drained.

Table 5. Representative soil features

Parent material	(1) Organic material (2) Loess (3) Colluvium
Surface texture	(1) Silt loam
Family particle size	(1) Loamy-skeletal

Drainage class	Well drained
Permeability class	Moderately rapid to rapid
Depth to restrictive layer	3–20 cm
Soil depth	150–178 cm
Surface fragment cover <=3"	0–25%
Surface fragment cover >3"	0–10%
Available water capacity (0-25.4cm)	1.27–7.37 cm
Calcium carbonate equivalent (0-101.6cm)	0–2%
Clay content (0-50.8cm)	5–15%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–3
Soil reaction (1:1 water) (0-101.6cm)	3.8–8.3
Subsurface fragment volume <=3" (0-101.6cm)	0–75%
Subsurface fragment volume >3" (0-101.6cm)	0–43%

Table 6. Representative soil features (actual values)

Drainage class	Somewhat poorly drained to excessively drained
Permeability class	Not specified
Depth to restrictive layer	3–178 cm
Soil depth	25–178 cm
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-25.4cm)	0.51–8.64 cm
Calcium carbonate equivalent (0-101.6cm)	Not specified
Clay content (0-50.8cm)	Not specified

Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-101.6cm)	Not specified
Subsurface fragment volume <=3" (0-101.6cm)	Not specified
Subsurface fragment volume >3" (0-101.6cm)	Not specified

Ecological dynamics

Growing conditions

Located in alpine life zone, ecological site R228XY704AK is exposed to a variety of harsh conditions including high winds, persistent snowpack, and extremely cold temperatures. Persistent snowpack and cold temperatures reduce the growing season in the alpine, when compared to lower elevations. As a result of these harsh growing conditions, alpine vegetation exhibits diminished growth forms in the low and dwarf shrubs that occupy this site.

Disturbance

Although fire plays an important role in shaping plant communities across Alaska, fire frequency in high elevation communities is largely unstudied, when compared to interior forest stands. Most wildfires in Alaska are caused by lightning strikes which tend to occur near tree line, decreasing in frequency into the subalpine and alpine zones (Dewilde et al, 2006). Despite the propensity of fires to move from boreal stands upslope into higher life zones, it is likely a general lack of fuel in scrub communities that accounts for diminished fire frequency (Kasischke et al. 2002, Dewild et al. 2006). The disturbance regime in this community is likely driven by avalanche, rockslides, and other mass movement events associated with eroding, unstable mountain slopes coupled with substantial snowfall.

Community 1.1

Field data suggest a single plant community exists on this ecological site. The reference plant community (1.1) is characterized by a dryas dwarf scrub tundra community (Viereck et al., 1992). Notable species include eightpetal mountain avens (*Dryas octopetala*), bog blueberry (*Vaccinium uliginosum*), White arctic mountain heather (*Cassiope tetragona*), black crowberry (*Empetrum nigrum*), several willow (*Salix* spp.) species, and various lichen species. Community composition in this alpine community is variable across plots, with some plots exhibiting greater cover in one dwarf shrub species or the other. This is likely due to several factors including retention of snowpack, recent disturbance or browsing pressure, as well as seed source. In addition to these factors, many of the

species above express prostrate ground-covering growth forms, taking any opportunity to spread out. This could lead to higher or lower cover values of key species and could lead one to assign a Viereck classification of both Dryas dwarf scrub (II.D.1 – level III) or Ericaceous dwarf scrub (II.D.2 – level III) (Viereck et al., 1992).

State and transition model

Ecosystem states

- 1. Reference state

State 1 submodel, plant communities

- 1.1. Eight petal mountain avens - bog blueberry - netleaf willow / reindeer lichen

State 1 Reference state

The reference state is a dryas dwarf scrub tundra community (Viereck et al., 1992). Common shrubs include eightpetal mountain avens (*Dryas octopetala*), bog blueberry (*Vaccinium uliginosum*), white arctic mountain heather (*Cassiope tetragona*), black crowberry (*Empetrum nigrum*), and several willow (*Salix* spp.) species. Lichen and moss cover are also considerable on this site.

Dominant plant species

- bog blueberry (*Vaccinium uliginosum*), shrub
- eightpetal mountain-avens (*Dryas octopetala* ssp. *octopetala*), shrub
- netleaf willow (*Salix reticulata*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- alpine bearberry (*Arctostaphylos alpina*), shrub
- reindeer lichen (*Cladina*), other herbaceous
- cup lichen (*Cladonia*), other herbaceous
- thuidium moss (*Thuidium recognitum*), other herbaceous

Community 1.1

Eight petal mountain avens - bog blueberry - netleaf willow / reindeer

lichen

The reference state is a dryas dwarf scrub tundra community (Viereck et al., 1992). The major plant groups are low and dwarf shrubs (less than 8 inches tall), medium graminoids (less than 24 inches tall), mosses, and lichen. Common shrubs include eightpetal mountain avens (*Dryas octopetala*), bog blueberry (*Vaccinium uliginosum*), white arctic mountain heather (*Cassiope tetragona*), black crowberry (*Empetrum nigrum*), and several willow (*Salix* spp.) species. Various lichen and moss species are common on this site, including reindeer lichen and thuidium moss.

Dominant plant species

- bog blueberry (*Vaccinium uliginosum*), shrub
- eightpetal mountain-avens (*Dryas octopetala* ssp. *octopetala*), shrub
- netleaf willow (*Salix reticulata*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- alpine bearberry (*Arctostaphylos alpina*), shrub
- polar willow (*Salix polaris*), shrub
- reindeer lichen (*Cladina*), other herbaceous
- cup lichen (*Cladonia*), other herbaceous
- thuidium moss (*Thuidium recognitum*), other herbaceous

Additional community tables

Inventory data references

The vegetation modeled for this site has limited data and is considered provisional. The associated model was largely developed from NRCS staff with working knowledge of the area and literature review.

Plant community composition is largely based on ecological sites from AK638: Soil Survey of Cantwell Area, Alaska.

References

Jorgensen, T. and D. Meidinger. 2015. The Alaska Yukon Region of the Circumboreal Vegetation map (CBVM). CAFF Strategies Series Report. Conservation of Arctic Flora and Fauna, Akureyri, Iceland..

Other references

Bernhardt, E.L., T.N. Hollingsworth. 2011. Fire severity mediates climate-driven shifts in understory community composition of black spruce stands of interior Alaska. *Journal of Vegetation Science* 22:32–44.

Chapin, F.S., L.A. Viereck, P.C. Adams, K.V. Cleve, C.L. Fastie, R.A. Ott, D. Mann, and J.F. Johnstone. 2006. Successional processes in the Alaskan boreal forest. Page 100 in Alaska's changing boreal forest. Oxford University Press.

Hinzman, L.D., L.A. Viereck, P.C. Adams, V.E. Romanovsky, and K. Yoshikawa. 2006. Climate and permafrost dynamics of the Alaskan boreal forest. Alaska's changing boreal forest 39–61.

I. H. Myers-Smith, J. W. Harden, M. Wilmking, C. C. Fuller, A. D. McGuire, and F. S. Chapin III. 2008. Wetland succession in a permafrost collapse: interactions between fire and thermokarst. *Biogeosciences* 5:1273–1286.

Johnstone, J.F., T.N. Hollingsworth, F.S. CHAPIN III, and M.C. Mack. 2010. Changes in fire regime break the legacy lock on successional trajectories in Alaskan boreal forest. *Global change biology* 16:1281–1295.

Johnstone, J.F., F.S. Chapin, T.N. Hollingsworth, M.C. Mack, V. Romanovsky, and M. Turetsky. 2010. Fire, climate change, and forest resilience in interior Alaska. *Canadian Journal of Forest Research* 40:1302–1312.

Johnstone, J.F. 2008. A key for predicting postfire successional trajectories in black spruce stands of interior Alaska. US Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Kelly, R., M.L. Chipman, P.E. Higuera, I. Stefanova, L.B. Brubaker, and F.S. Hu. 2013. Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National Academy of Sciences* 110:13055–13060.

LANDFIRE. 2009. Western North American Boreal Black Spruce Dwarf-tree Peatland – Boreal Complex. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.

Schoeneberger, P.J. and D.A. Wysocki. 2012. Geomorphic Description System. Natural Resources Conservation Service, 4.2 edition. National Soil Survey Center, Lincoln, NE.

Smith, R.D., A.P. Ammann, C.C. Bartoldus, and M.M. Brinson. 1995. An approach for assessing wetland functions using hydrogeomorphic classification, reference wetlands, and functional indices.

United States Department of Agriculture. 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-

Contributors

Tyler Annetts
Matthew J Mayer

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	01/22/2026
Approved by	Blaine Spellman
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. Amount of plant mortality and decadence (include which functional groups are

expected to show mortality or decadence):

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
