

Ecological site F229XY033AK

Boreal Forest Loamy Slopes

Last updated: 6/12/2025

Accessed: 03/24/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 229X–Interior Alaska Lowlands

The Interior Alaska Lowlands MLRA includes the flood plains and terraces along the upper reaches of the Tanana and Kuskokwim Rivers and the middle reaches of the Yukon River. This area makes up 39,065 square miles. The northern portion of this area that contains the cities of Fairbanks and North Pole are the second most densely populated lands in Alaska. Other towns along the road system include Nenana, Delta Junction, and Tok; and parts of Fort Wainwright and Fort Greely, the two largest military reservations in Alaska. Elsewhere, the area is mostly undeveloped wild land and is sparsely populated. In the western part of the area, the communities of Tanana, Galena, and McGrath are accessible only by air or by river. Parts of the Denali National Park and Preserve and Tetlin National Wildlife Refuge are in this area. The Trans-Alaska Pipeline parallels the Alaska Highway from Delta Junction to Fairbanks.

This area is on broad, nearly level, braided to meandering flood plains, stream terraces, and outwash plains. In many places, shallow basins and undulating stream terraces are dotted with hundreds of small and medium size lakes and interconnecting wetlands. Sloughs, oxbow lakes, and escarpments along river channels are features associated with the flood plains, terraces, and basins. Isolated bedrock-controlled hills and low- to moderate-relief mountains are in scattered places throughout the area. Extended foot slopes are common at the base of hills and mountains and along the boundaries with adjoining mountainous MLRAs. Elevation ranges from about 100 feet in the southwestern part of the area, along the lower Yukon River, to about 1,900 feet in the upper Tanana Valley.

Geology and Soils

Although never glaciated, this area is filled with a deep layer of Pleistocene glaciofluvial deposits. Additional fluvial sediments from the Alaska Range and the northern Aleutian Range accumulated along the Tanana and Kuskokwim Rivers during the Holocene Epoch. The Koyukuk and lower Yukon River drainages have undergone several periods of deposition followed by erosion. In some places old terraces are 33 to 250 feet above the flood plain. Quaternary glaciofluvial and fluvial sediments are estimated to be as much as 330 to 660 feet thick throughout the area. Much of the MLRA, particularly along the Tanana and Kuskokwim Rivers, is mantled with a layer of silty micaceous loess originating from the unvegetated flood plains and outwash plains along the Alaska Range. Thick eolian deposits, including loess dunes, sand dunes, and sand sheets, make up about 12 percent of the area. Inclusions of glacial moraines and drift are near the mountains. Unconsolidated sediments bury the bedrock geology, except for structural hills in some places.

This area is in the zone of discontinuous permafrost. Permafrost is close to the surface in lands with finer textured silty sediments on stream terraces, plains, and the more gently sloping foot slopes and hills throughout the area. Isolated masses of ground ice occur on terraces and the lower side slopes of hills. Permafrost does not generally occur on floodplains, soils with sandy and gravelly parent material, and in areas near lakes and other water bodies.

The dominant soil orders in this area are Gelisols, Entisols, Inceptisols, and Spodosols. The Gelisols are shallow or moderately deep to permafrost, occur on finer textured sediments, and are poorly drained or very poorly drained. Common Gelisol suborders are Histels, Orthels, and Turbels. The Histels have thick accumulations of surface organic material and occur in depressions and peat plateau. The Orthels and Turbels have comparably thinner surface organic material and occur on drainageways, stream terraces, and outwash plains. The Inceptisols, Spodosols, and Entisols lack permafrost in the soil profile. Two important factors that prevent permafrost aggradation are groundwater connectivity and thick bands of sandy and/or gravelly soil horizons. Entisols most commonly occur on the floodplain with the most common suborder being Fluvents. Inceptisols and Spodosols occurs on streams terraces and slopes of hills and outwash plains. Dry soils on these landforms support soil suborders related to Cryods and Cryepts, while wet soils support Aquepts. Miscellaneous (non-soil) areas make up about 19 percent of this MLRA. The most common are riverwash and water.

For Gelisols, wildfires disturb the insulating organic material at the soil surface and can change the presence and/or depth of permafrost in the soil profile. These fire related changes to permafrost can also change the depth and presence of perched water tables. Gelisols that burn in this area can change soil taxonomic classification. For instance, depending on fire-severity, Histels may change to Orthels and Orthels may change to Inceptisols. Depending on the frequency and intensity of fires, landform position, and soil texture, the soils may or may not revert back to their original taxonomic classification.

Climate

Short, warm summers and long, very cold winters characterize the continental subarctic climate of the area. The average annual precipitation for this area ranges from 11 to 17 inches. The maximum precipitation occurs in late summer, mainly during thunderstorms. The average annual snowfall ranges from 30 to 80 inches. The average annual temperature for this area ranges from about 23 to 30 degrees Fahrenheit. The freeze-free period averages about 90 to 110 days. The temperature usually remains above freezing from June through early-September.

Vegetation

Much of the soil in this area supports forested communities in some stage of post-fire recovery. Mesic to dry soils in the uplands support mixed forests of black spruce, white spruce, paper birch, and quaking aspen. White spruce and white spruce-balsam poplar forests occur on occasionally to rarely flooded portions of the floodplain-step. Black spruce woodlands occur on stream terraces and other places with wet soils and/or shallow permafrost. On permafrost-affected alluvial flats, tamarack occurs in association with black spruce. Lightning-caused wildfires are common. Many thousands of acres are often burned during a single fire. Following wildfires, willow, shrub birch, and ericaceous shrub scrub invade most previously forested ecological sites until they eventually are replaced by forest vegetation. After fire, resin birch is common on wet and mesic soils and quaking aspen on dry soils. On all forest and woodland ecological sites, post-fire succession leads to a relatively rapid accumulation of organic matter and mosses on the surface. This accumulation results in decreases in soil temperature, biologic activity, and nutrient availability and a gradual decrease in site productivity.

Frequently flooded and continuously ponded soils lack forested communities. Non-forested soils include shrub birch, ericaceous shrub scrub, and tussock tundra in peat areas and in drainageways. Wet sedge meadows, sedge-moss bog meadows, and sedge-grass meadows are along the margins of lakes and on continuously ponded sites such as fens. Low to tall willow and alder scrub are on low flood plains.

Classification relationships

Landfire BPS – 16030 – Western North American Boreal White Spruce-Hardwood Forest (Landfire 2009)

Landfire BPS – 16050 – Western North American Boreal Mesic Birch-Aspen Forest (Landfire 2009)

Ecological site concept

- Associated with well drained soils on dunes, sand sheets, stream terraces, and slopes of hills and outwash plains.
- Soils formed in silty parent material such as loess over sandy or gravelly material such

as sandy eolian deposits and alluvium. Mineral soils are capped with two to five inches of organic material.

- Soils do not pond or flood. A water table occurs at deep depths during the growing season.

- Soils are considered very deep and lack restrictions like permafrost.

- The reference plant community is open forest (Viereck et al. 1992) with black spruce and/or white spruce the dominant trees. Commonly observed understory species include russet buffaloberry, bog Labrador tea, lingonberry, false toadflax, splendid feathermoss, and Schreber's big redstem moss. Five plant communities have been identified within the reference state related to fire.

Associated sites

F229XY021AK	Boreal Peat Terraces and Slopes Complex Occurs on the same terraces and slopes but in lower positions with wetter soils that pond frequently and have permafrost. Associated with black spruce stunted woodlands.
F229XY031AK	Boreal Forest Loamy Flats and Slopes Moist Occurs on the same slopes but in lower positions with moister soils. Associated with black spruce forests.
R229XY010AK	Boreal Scrub Loamy Frozen Drainageways Occurs in adjacent drainageways with closed low scrub vegetation and abundant obligate wetland species.
F229XY020AK	Boreal Woodland Loamy Frozen Slopes Occurs on the same slopes but in lower positions with wetter soils that have permafrost. Associated with black spruce woodlands.
F229XY032AK	Boreal Woodland Sandy Slopes Occurs on the same slopes but on drier soils. Associated with black spruce woodlands.

Similar sites

F229XY031AK	Boreal Forest Loamy Flats and Slopes Moist Ecological site 31 occurs on the same slopes but in lower positions with moister soils. Ecological site 31 has less productive trees and more facultative wet plant species.
F231XY186AK	Boreal Forest Silty Slopes Ecological site 186 occurs on hills with similar soils and vegetation but occurs to the North in the Interior Alaska Highlands MLRA.
XA232X02Y210	Boreal Forest Loamy Frozen Plains Warm Ecological site 210 occurs on hills and plains with similar soils and vegetation but occurs to the North in the Yukon Flats Lowlands MLRA.

Table 1. Dominant plant species

Tree	(1) <i>Picea glauca</i> (2) <i>Picea mariana</i>
Shrub	(1) <i>Shepherdia canadensis</i> (2) <i>Vaccinium vitis-idaea</i>
Herbaceous	(1) <i>Hylocomium splendens</i> (2) <i>Pleurozium schreberi</i>

Physiographic features

- Typically occurs on stream terraces, hills, outwash plains, dunes, and sand sheets. Rarely occurs on moraines. Hillslope profile is summits, shoulders, and backslopes.
- Elevation occurs between 400 and 1200 feet.
- Occurs on warmer south and west facing backslopes. Terraces and summits of hills are nearly level, while dunes and backslopes are gently to strongly sloping.
- This ecological site does not typically flood or pond. On occasion, soils pond but before the growing season and during the months of April and/or May. A water table occurs in the soil profile at deep depths (40 and 60 inches).
- Associated with negligible to low amounts of runoff to adjacent, downslope ecological sites.

Table 2. Representative physiographic features

Landforms	(1) Alluvial plain > Stream terrace (2) Alluvial plain > Dune (3) Lowland > Hill (4) Lowland > Sand sheet (5) Lowland > Alluvial fan (6) Lowland > Hill (7) Lowland > Plain (8) Lowland > Moraine
Runoff class	Negligible to low
Flooding frequency	None
Ponding duration	Not specified
Ponding frequency	None
Elevation	122–366 m
Slope	1–12%
Water table depth	102–152 cm
Aspect	W, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
--------------	---------------

Flooding frequency	Not specified
Ponding duration	Long (7 to 30 days)
Ponding frequency	None to occasional
Elevation	30–579 m
Slope	0–45%
Water table depth	Not specified

Climatic features

Short, warm summers and long, very cold winters characterize the subarctic continental climate associated with this ecological site. The mean annual temperature for Interior Alaska Lowlands area ranges from 23 to 31 degrees Fahrenheit. The warmest months span May through August with mean temperatures ranging from 60 to 71 degrees Fahrenheit. The coldest months span December through February with mean temperatures ranging from -15 to -10 degrees Fahrenheit. The freeze-free period ranges between 96 and 115 days and typically lasts from late May through early-September.

The average annual precipitation across the Interior Alaska Lowlands area ranges between 11 and 17 inches. July through September are the wettest months with approximately 60 percent of the annual precipitation occurring, and thunderstorms are common. The average annual snowfall ranges from 30 to 80 inches (USDA 2022). The ground is consistently covered with snow from November through March.

Table 4. Representative climatic features

Frost-free period (characteristic range)	68-84 days
Freeze-free period (characteristic range)	96-115 days
Precipitation total (characteristic range)	305-356 mm
Frost-free period (actual range)	56-94 days
Freeze-free period (actual range)	88-120 days
Precipitation total (actual range)	279-432 mm
Frost-free period (average)	75 days
Freeze-free period (average)	105 days
Precipitation total (average)	330 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) GALENA [USC00503212], Nulato, AK
- (2) TANANA CALHOUN MEM AP [USW00026529], Tanana, AK
- (3) MCGRATH AP [USW00026510], Mc Grath, AK
- (4) MINCHUMINA [USW00026512], Lake Minchumina, AK

- (5) KOBE HILL [USC00504971], Healy, AK
- (6) NENANA MUNI AP [USW00026435], Clear, AK
- (7) FAIRBANKS INTL AP [USW00026411], Fairbanks, AK
- (8) NORTH POLE [USC00506581], North Pole, AK
- (9) EIELSON FLD [USC00502707], Eielson AFB, AK
- (10) SALCHA [USC00508140], Salcha, AK
- (11) BIG DELTA AP [USW00026415], Delta Junction, AK

Influencing water features

Due to its landscape position, this ecological site is neither associated with or influenced by streams or wetlands. Precipitation and throughflow are the main source of water for this ecological site. Surface runoff and throughflow contribute some water to downslope ecological sites.

Depth to the water table may decrease following summer storm events or spring snowmelt and increase during extended dry periods.

Wetland description

Not a wetland.

Soil features

- Soils formed in silty parent material over sandy or gravelly parent material. The silty material is loess or glaciofluvial deposits, while the sandy or gravelly material is sandy eolian deposits, glaciofluvial deposits, and/or alluvium.
- Rock fragments do not occur on the soil surface.
- Mineral soils are capped with two to five inches of organic material. The surface mineral horizon is typically a silt loam but on occasion is a fine sandy loam or very fine sandy loam.
- Soils are typically very deep and lack restrictions. On rare occasion, soils contact bedrock at moderate to deep depths (30 to 60 inches) or permafrost at shallow depth (11 to 16 inches).
- Soils typically lack rock fragments in the soil profile. In soils with gravelly material at depth, rock fragments can range up to 65 percent of the soil profile by volume.
- The pH of the soil profile ranges from strongly acidic to slightly acidic.
- Soils are considered well drained.

Table 5. Representative soil features

Parent material	(1) Loess (2) Eolian deposits (3) Glaciofluvial deposits (4) Alluvium
-----------------	--

Surface texture	(1) Silt loam (2) Fine sandy loam (3) Very fine sandy loam
Family particle size	(1) Coarse-loamy (2) Coarse-loamy over sandy or sandy-skeletal (3) Coarse-silty (4) Coarse-silty over sandy or sandy-skeletal
Drainage class	Well drained
Permeability class	Moderately rapid
Depth to restrictive layer	Not specified
Soil depth	152 cm
Surface fragment cover $\leq 3''$	0%
Surface fragment cover $> 3''$	0%
Available water capacity (0-101.6cm)	10.16–17.53 cm
Calcium carbonate equivalent (25.4-101.6cm)	0%
Clay content (0-50.8cm)	5–9%
Electrical conductivity (25.4-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (25.4-101.6cm)	0
Soil reaction (1:1 water) (25.4-101.6cm)	5.1–6.5
Subsurface fragment volume $\leq 3''$ (0-152.4cm)	0–2%
Subsurface fragment volume $> 3''$ (0-152.4cm)	0%

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Permeability class	Moderately slow to rapid
Depth to restrictive layer	28–152 cm
Soil depth	127–152 cm
Surface fragment cover $\leq 3''$	Not specified
Surface fragment cover $> 3''$	Not specified

Available water capacity (0-101.6cm)	Not specified
Calcium carbonate equivalent (25.4-101.6cm)	0–10%
Clay content (0-50.8cm)	Not specified
Electrical conductivity (25.4-101.6cm)	Not specified
Sodium adsorption ratio (25.4-101.6cm)	0–2
Soil reaction (1:1 water) (25.4-101.6cm)	4–8.4
Subsurface fragment volume ≤3" (0-152.4cm)	0–50%
Subsurface fragment volume >3" (0-152.4cm)	0–15%

Ecological dynamics

Fire

In the Interior Alaska Lowlands MLRA, fire is a common and natural event that has a significant control on the vegetation dynamics across the landscape. A typical fire event in the lands associated with this ecological site will reset plant succession and alter dynamic soil properties (e.g., thickness of the organic material). For this ecological site to progress from the earliest stages of post-fire succession dominated by grasses and forbs to the oldest stages of succession dominated by white spruce and black spruce forests, data suggest that 130 to 150 years or more must elapse without another fire event (Foot 1982; Chapin et al. 2006; Landfire 2009).

Within this area, wildfire is considered a natural and common event that in many places goes unmanaged. Fire suppression is limited and occurs adjacent to the municipalities spread throughout the area or on allotments with known structures, all of which have a relatively limited acre footprint. Most fires are caused by lightning strikes. From 2000 to 2020, 513 known fire events occurred in this area and the burn perimeter of the fires totaled approximately 12.4 million acres (AICC 2022). Fire-related disturbances are highly patchy and can leave undisturbed areas within the burn perimeter. During this time frame, 73 percent of the fire events were smaller than 20,000 acres but 34 fire events were greater than 100,000 acres in size (AICC 2022). Over this period of 20 years, these burn perimeters cover approximately 50 percent of this area.

The fire regime within Interior Alaska follows two general scenarios—low-severity burns and high-severity burns. It should be noted, however, that the fire regime in Interior Alaska

can be considered more complex (Johnstone et al. 2008). Burn severity refers to the proportion of the vegetative canopy and organic material consumed in a fire event (Chapin et al. 2006). Fires in cool and moist habitat tend to result in low-severity burns, while fires in warm and dry habitat tend to result in high-severity burns. Because the soils have a thin organic cap and are well drained, the typical fire scenario for this ecological site is considered to result in a high severity burn.

Large portions of the organic mat are consumed during a high-severity fire event, commonly exposing pockets of mineral soil. The loss of this organic mat, which insulates the mineral soil, and the decrease in site albedo tends to cause overall soil temperatures to increase (Hinzman et al. 2006). These alterations to soil temperature may result in increased depths of seasonal frost in the soil profile. High-severity fire events also destroy a majority of the vascular and nonvascular biomass above ground.

Field data suggest that each of the forested communities burn and that fire events will cause a transition to the pioneering stage of fire succession. This stage (community 1.5) is a mix of species that either regenerate in place (e.g., subterranean root crowns for willow and rhizomes for graminoids) and/or from wind-dispersed seed or spores that colonize exposed mineral soil (e.g., quaking aspen [*Populus tremuloides*] and *Ceratodon* moss [*Ceratodon purpureus*]). The pioneering stage of fire succession is primarily composed of tree seedlings, forbs, grasses, and weedy bryophytes. This stage of succession is thought to persist for up to 4 years post-fire (Landfire 2009). Willow (*Salix* spp.) and quick growing deciduous tree seedlings continue to colonize and grow in stature on recently burned sites until they become dominant in the overstory, which marks the transition to the early stage of fire succession (community 1.4). This early stage of fire succession is thought to persist 30 to 50 years post-fire (Landfire 2009). In the absence of fire, tree species continue to become more dominant in the stand and eventually develop into forests.


The latter stages of succession have an overstory that is dominantly deciduous trees (community 1.3), a mix of broadleaf and needleleaf trees (community 1.2), or needleleaf trees (community 1.1). The recruitment of trees species during the pioneering and early stages of post-fire succession largely controls the composition of the stand of trees in the later stages of post-fire succession (Johnstone et al. 2010a). During these later stages of succession, the slower growing spruce seedlings mature and eventually replace the shade-intolerant broadleaf tree species. The typical fire return interval for white spruce stands in Interior Alaska is 150 years (Landfire 2009; Abrahamson 2014).

Lands in Interior Alaska are burning more frequently than in the past, which may result in alternative states of succession. The historic fire return interval for white spruce stands in Interior Alaska occurs approximately once every 150 years (Landfire 2009; Abrahamson 2014). Due to global climate change, stands of spruce in certain portions of the Alaskan boreal forest are burning more frequently than these historic averages (Kelly et al. 2013). Increases to burn frequency favors forested stands dominated by quick growing deciduous trees (community 1.3). A major reason being that increased fire frequency decreases the presence and abundance of mature, cone-bearing trees. Less mature trees


result in less spruce seedlings post-fire and an overall decreased abundance of spruce in the developing forest canopy. Increased fire frequency paired with decreased spruce propagules lowers the chances for the site to progress to post-fire successional communities 1.2 and 1.1. Increased burn frequency in the boreal forest may result in alternative states of post-fire succession with stands of deciduous trees persisting for longer than normal durations of time (Johnstone et al. 2010b).

State and transition model


Ecosystem states



State 1 submodel, plant communities



Communities 1, 5 and 2 (additional pathways)



1.1a - A high-severity fire sweeps through and incinerates much of the above ground vegetation

1.2b - 130 to 150 years without wildfire

1.2a - A high-severity fire sweeps through and incinerates much of the above ground vegetation

1.3b - 80 to 100 years without fire.

1.3a - A high-severity fire sweeps through and incinerates much of the above ground vegetation

1.4b - 30 to 50 years without fire

1.4a - A high-severity fire sweeps through and incinerates much of the above ground vegetation

1.5a - 4 to 6 years without wildfire

State 1 Reference State



Figure 7. Typical vegetation associated with the reference state. Photo was taken in the Totchaket area soil survey.

The reference plant community is open needleleaf forest (Vioreck et al. 1992) with the dominant trees being white spruce and black spruce. There are five plant communities within the reference state related to fire. The vegetation modeled for this ecological site has limited data and is considered provisional.

Dominant plant species

- white spruce (*Picea glauca*), tree
- black spruce (*Picea mariana*), tree
- russet buffaloberry (*Shepherdia canadensis*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous

Community 1.1

black spruce - white spruce / russet buffaloberry - lingonberry / splendid feathermoss - Schreber's big redstem moss



Figure 8. Typical plant community associated with community 1.1. Photo is from the Totchaket area soil survey.

The reference plant community is characterized as open needleleaf forest (Vioreck et al. 1992) with white spruce and/or black spruce the dominant trees. Spruce cover primarily occurs in the tall tree strata (greater than 40 feet). Live deciduous trees, primarily quaking aspen, occasionally occur in the tree canopy but with limited cover. Common understory species include russet buffaloberry, lingonberry, bog Labrador tea, bluejoint, false toadflax, splendid feathermoss, and Schreber's big redstem moss. The soil surface is primarily covered with herbaceous litter and moss.

Dominant plant species

- black spruce (*Picea mariana*), tree
- white spruce (*Picea glauca*), tree

- lingonberry (*Vaccinium vitis-idaea*), shrub
- russet buffaloberry (*Shepherdia canadensis*), shrub
- bog Labrador tea (*Ledum groenlandicum*), shrub
- Bebb willow (*Salix bebbiana*), shrub
- twinflower (*Linnaea borealis*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- false toadflax (*Geocaulon lividum*), other herbaceous
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous
- bunchberry dogwood (*Cornus canadensis*), other herbaceous
- groundcedar (*Lycopodium complanatum*), other herbaceous
- felt lichen (*Peltigera*), other herbaceous

Community 1.2

quaking aspen - spruce / russet buffaloberry - lingonberry / splendid feathermoss - Schreber's big redstem moss



Figure 9. Typical plant community associated with community 1.2. Photo is from the Totchaket area soil survey.

Community 1.2 is in the late stage of fire-induced secondary succession for this ecological site. It is characterized as open mixed forest (Viereck et al. 1992) with mature quaking aspen and a mixture of immature and mature spruce as the dominant trees. Tree cover is split between the tall tree (greater than 40 feet) and medium tree strata (between 15 and 40 feet). Common understory species include russet buffaloberry, lingonberry, bog Labrador tea, bluejoint, false toadflax, splendid feathermoss, and Schreber's big redstem moss. The soil surface is primarily covered with herbaceous litter and moss.

Dominant plant species

- quaking aspen (*Populus tremuloides*), tree
- white spruce (*Picea glauca*), tree

- black spruce (*Picea mariana*), tree
- lingonberry (*Vaccinium vitis-idaea*), shrub
- bog Labrador tea (*Ledum groenlandicum*), shrub
- prickly rose (*Rosa acicularis*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- false toadflax (*Geocaulon lividum*), other herbaceous
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous
- bunchberry dogwood (*Cornus canadensis*), other herbaceous

Community 1.3

quaking aspen / bog Labrador tea - lingonberry / bluejoint



Figure 10. Typical plant community associated with community 1.3. Photo is from the Totchaket area soil survey.

Community 1.3 is in the middle stage of fire-induced secondary succession for this ecological site. It is characterized as open deciduous forest (Viereck et al. 1992) with quaking aspen the dominant tree. Seedlings and saplings of spruce are common but have comparatively limited cover. Tree cover is split between the tall tree (greater than 40 feet) and medium tree strata (between 15 and 40 feet). Common understory species include bog Labrador tea, lingonberry, russet buffaloberry, Bebb's willow, bluejoint, and bunchberry dogwood. The soil surface is primarily covered with herbaceous litter.

Dominant plant species

- quaking aspen (*Populus tremuloides*), tree
- white spruce (*Picea glauca*), tree
- black spruce (*Picea mariana*), tree
- bog Labrador tea (*Ledum groenlandicum*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- Bebb willow (*Salix bebbiana*), shrub

- russet buffaloberry (*Shepherdia canadensis*), shrub
- twinflower (*Linnaea borealis*), shrub
- prickly rose (*Rosa acicularis*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- bunchberry dogwood (*Cornus canadensis*), other herbaceous
- ceratodon moss (*Ceratodon purpureus*), other herbaceous
- juniper polytrichum moss (*Polytrichum juniperinum*), other herbaceous
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- alpine sweetvetch (*Hedysarum alpinum*), other herbaceous

Community 1.4

Bebb willow - lingonberry / bluejoint / bunchberry dogwood



Figure 11. Typical plant community associated with community 1.4. Photo is from the Totchaket area soil survey.

Community 1.4 is in the early stage of fire-induced secondary succession for this ecological site. It is best characterized as open tall scrub (Vioreck et al. 1992) with saplings of quaking aspen and Bebb willow the dominant overstory vegetation. Other common species include lingonberry, bog Labrador tea, russet buffaloberry, bluejoint, bunchberry dogwood, juniper polytrichum moss, and Ceratodon moss. The soil surface is primarily covered with woody litter, herbaceous litter, and weedy mosses.

Dominant plant species

- quaking aspen (*Populus tremuloides*), tree
- lingonberry (*Vaccinium vitis-idaea*), shrub
- bog Labrador tea (*Ledum groenlandicum*), shrub
- Bebb willow (*Salix bebbiana*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- Altai fescue (*Festuca altaica*), grass
- ceratodon moss (*Ceratodon purpureus*), other herbaceous

- juniper polytrichum moss (*Polytrichum juniperinum*), other herbaceous
- bunchberry dogwood (*Cornus canadensis*), other herbaceous

Community 1.5

bluejoint / fireweed / juniper polytrichum moss

Community 1.5 is in the pioneering stage of fire-induced secondary succession for this ecological site. It is characterized as a mesic forb or mesic graminoid herbaceous community (Viereck et al. 1992). Tree seedlings of primarily quaking aspen and spruce are common throughout the community but have limited cover. Commonly observed species include bluejoint, fireweed, juniper polytrichum moss, and Ceratodon moss. The soil surface is primarily covered with a mixture of weedy bryophyte species, woody debris, and herbaceous litter.

Dominant plant species

- Bebb willow (*Salix bebbiana*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- Altai fescue (*Festuca altaica*), grass
- juniper polytrichum moss (*Polytrichum juniperinum*), other herbaceous
- ceratodon moss (*Ceratodon purpureus*), other herbaceous
- fireweed (*Chamerion angustifolium*), other herbaceous
- pohlia moss (*Pohlia nutans*), other herbaceous
- (*Marchantia polymorpha*), other herbaceous

Pathway 1.1a

Community 1.1 to 1.5

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

Pathway 1.2b

Community 1.2 to 1.1



quaking aspen - spruce /
russet buffaloberry -
lingonberry / splendid
feathermoss - Schreber's big
redstem moss



black spruce - white spruce /
russet buffaloberry -
lingonberry / splendid
feathermoss - Schreber's big
redstem moss

Community pathway 1.2b occurs 130 to 150 years after wildfire disturbance (Foot 1982; Landfire 2009). Spruce replace quaking aspen in the tree canopy and the community turns into a needleleaf forest community.

Pathway 1.2a **Community 1.2 to 1.5**

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

Pathway 1.3b **Community 1.3 to 1.2**



quaking aspen / bog Labrador tea - lingonberry / bluejoint



quaking aspen - spruce / russet buffaloberry - lingonberry / splendid feathermoss - Schreber's big redstem moss

Community pathway 1.3b is thought to occur 80 to 100 years after wildfire disturbance. Spruce and feathermoss cover increases and the community turns into a mixed forest community.

Pathway 1.3a **Community 1.3 to 1.5**

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

Pathway 1.4b **Community 1.4 to 1.3**



Bebb willow - lingonberry / bluejoint / bunchberry dogwood



quaking aspen / bog Labrador tea - lingonberry / bluejoint

Community pathway 1.4a is thought to occur 30 to 50 years after wildfire disturbance. Quaking aspen mature are turn into a deciduous forest community.

**Pathway 1.4a
Community 1.4 to 1.5**

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

**Pathway 1.5a
Community 1.5 to 1.4**

Community pathway 1.5a is thought to occur 4 to 6 years after disturbance. Deciduous tree and willow cover increases.

Additional community tables

Animal community

not available

Hydrological functions

not available

Recreational uses

not available

Wood products

not available

Other products

not available

Other information

not available

Inventory data references

The vegetation modeled for this site has limited data coverage across the MLRA and is considered provisional. The associated model was largely developed from NRCS staff with working knowledge, a soil survey of Denali National Park Area (Clark and Duffy 2006), and sample plots from the Totchaket Area Soil Survey (AK625). Tier 2 sampling plots used to develop the reference state are below:

2021AK290008, 2021AK290011, 2021AK290013, 2021AK290102, 2021AK290104, 2021AK290105, 2021AK290106, 2021AK290108, 2021AK290109, 2021AK290112, 2021AK290114, 2021AK290115, 2021AK290117, 2021AK290119, 2021AK290121, 2021AK290122, 2021AK290123, 2021AK290125, 2021AK290126, 2021AK290129, 2021AK290136, 2021AK290140, 2021AK290201, 2021AK290203, 2021AK290208, 2021AK290209, 2021AK290210, 2021AK290212, 2021AK290213, 2021AK290214, 2021AK290217, 2021AK290219, 2021AK290220, 2021AK290302, 2021AK290303, 2021AK290306, 2021AK290308, 2021AK290309, 2021AK290312, 2021AK290314, 2021AK290316, 2021AK290317, 2021AK290319, 2021AK290321, 2021AK290322, 2021AK290323, 2021AK290324, 2021AK290325, 2021AK290327, 2021AK290328, 2021AK290329, 2021AK290330, 2021AK290334, 2021AK290455, 2021AK290459

References

Abrahamson, I.L. 2014. Fire Regimes of Alaskan White Spruce Communities.

Chapin, F.S., L.A. Viereck, P.C. Adams, K.V. Cleve, C.L. Fastie, R.A. Ott, D. Mann, and J.F. Johnstone. 2006. Successional processes in the Alaskan boreal forest. Page 100 in Alaska's changing boreal forest. Oxford University Press.

Foote, M.J. 1983. Classification, description, and dynamics of plant communities after fire in the taiga of interior Alaska. US Department of Agriculture, Forest Service, Pacific Northwest Forest and

Hinzman, L.D., L.A. Viereck, P.C. Adams, V.E. Romanovsky, and K. Yoshikawa. 2006. Climate and permafrost dynamics of the Alaskan boreal forest. Alaska's changing boreal forest 39–61.

Johnstone, J.F., F.S. Chapin, T.N. Hollingsworth, M.C. Mack, V. Romanovsky, and M. Turetsky. 2010. Fire, climate change, and forest resilience in interior Alaska. *Canadian Journal of Forest Research* 40:1302–1312.

Kelly, R., M.L. Chipman, P.E. Higuera, I. Stefanova, L.B. Brubaker, and F.S. Hu. 2013. Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National Academy of Sciences* 110:13055–13060.

Schoeneberger, P.J. and D.A. Wysocki. 2012. *Geomorphic Description System*. Natural Resources Conservation Service, 4.2 edition. National Soil Survey Center, Lincoln, NE.

United States Department of Agriculture, . 2022. *Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin*.

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. *The Alaska vegetation classification*. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Other references

Alaska Interagency Coordination Center (AICC). 2022. <http://fire.ak.blm.gov/>

LANDFIRE. 2009. *Western North American Boreal White Spruce-Hardwood Forest*. In: *LANDFIRE National Vegetation Dynamics Models*. USDA Forest Service and US Department of Interior. Washington, DC.

Contributors

Blaine Spellman

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	03/24/2026
Approved by	Blaine Spellman
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment**

and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability:
