

# Ecological site F230XY611AK

## Boreal Forest Loamy Frozen Slopes

Last updated: 6/11/2025

Accessed: 03/28/2026

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 230X–Yukon-Kuskokwim Highlands

#### MLRA notes

The Yukon-Kuskokwim Highlands (MLRA 230X) include the most western parts of Interior Alaska (Land Resource Region X2) and have a continental climate. MLRA 230X is approximately 42,300 square miles spread across mountain, hills, and valleys. Flood plain systems are common. The watershed drains into the Bering Sea to the west and Bristol Bay to the southwest. Major rivers include the Yukon, Innoko, Kuskokwim, Mulchatna, and Nushagak Rivers. This sparsely populated area is mostly undeveloped wildland. Residents use this remote area primarily for subsistence hunting, fishing, and gathering. Villages are primarily located along rivers along the MLRA 230X boundary and include Greyling, Nulato, and Koyukuk. Federally managed lands in the MLRA include parts of Innoko, Nowitna, and Koyukuk National Wildlife Refuges.

#### Geology and Soils

The Yukon-Kuskokwim Highlands MLRA was mostly unglaciated during the Pleistocene. Glaciers were limited to the Lime Hills in the southeast. Glacial moraines and drift are evident in areas of past glacial activity. Unglaciated upland areas are covered with colluvium and slope alluvium originating from bedrock. Loess deposits cover gentle sloping hills and footslopes of mountains near major rivers. Bedrock material is primarily sedimentary rocks with intrusive volcanic rock (USDA, 2022).

This MLRA is in the zone of discontinuous permafrost. Permafrost is most common in finely textured soils on terraces, gently sloping hills, and cold mountain footslopes. It is typically absent from flood plains and mountain backslopes. Across the MLRA, permafrost

presence decreases as proximity to the Yukon-Kuskokwim delta increases.

The dominant soil orders are Gelisols, Entisols, Inceptisols, and Spodosols. Gelisols support shallow to deep permafrost and often have a perched water table for at least part of the growing season. Inceptisols, Spodosols, and Entisols lack permafrost. Two important factors that prevent permafrost aggradation are groundwater connectivity and thick bands of sandy and/or gravelly soil horizons. Inceptisols have minimal development and are common on alpine scrublands and high flood plains. Entisols are common on mountain backslopes and scoured flood plains. Spodosols support a spodic soil horizon and are common in the acidic soils underlying spruce forests and ericaceous shrublands. Non-soil areas such as rock outcrops, rubble lands and beaches make up approximately ten percent of the MLRA surface.

## Climate

The Yukon-Kuskokwim Highlands MLRA has short, warm summers and cold, long winters. Mean annual precipitation is 10 to 15 inches at low elevations and increases to 20 to 40 inches at higher elevations (USDA, 2022). Annual snowfall is between 80 and 100 inches. Mean annual temperatures ranges from 25 to 32 degrees F (SNAP, 2014a; SNAP, 2014b).

## Vegetation

Vegetation is mainly influenced by site and soil characteristics such as temperature-degree days, exposure, soil depth, and soil hydrology. Dwarf scrublands are prevalent on shallow soils on convex slopes and in the alpine. Mesic, lowland slopes are a mix of forests and shrublands of alder, willow, and ericaceous shrubs. Cold slopes generally support black spruce, while warm slopes support white spruce. Valley bottoms and steep slopes support a deciduous forest. Tussock tundra is associated with wet soils underlain by shallow permafrost and is ubiquitous across the lower footslopes of mountains and the coastal plain (USDA, 2022).

## Fire

Fire is a major disturbance across the Yukon-Kuskokwim Highlands. Low severity fires destroy the canopy but leave the organic mat and rootstock mostly undisturbed. The vegetative community progresses directly back to a forest. Severe forest fires are stand replacement events. Post-fire communities typically pass through an herbaceous meadow community before ericaceous shrubs, birch, and willows colonize. Drier soils may support a deciduous aspen or birch forest, while moist soils support cottonwoods and spruce. On all forest and woodland ecological sites, post-fire succession leads to a relatively rapid accumulation of organic matter and mosses on the surface. This accumulation results in decreases in soil temperature, biologic activity, and nutrient availability and a gradual decrease in site productivity.

## **LRU notes**

MLRA 230X contains three life zones defined by the physiological limits of plant communities along an elevational gradient. The boreal life zone is the elevational band where forest communities dominate. Non-forested areas in the boreal life zone are often hydrologically driven, and are either too wet (i.e., bogs) or too dry (i.e., river bluffs) to support forest communities. Subalpine and alpine vegetation dominates at higher elevations. The subalpine zone is a transitional band between the boreal and the alpine life zones, and is characterized by sparse, stunted trees. Shrub height can be over four feet. Trees are absent from the alpine, and all shrubs are dwarf or prostrate. In general, the boreal life zone occurs below 1,200 feet; the subalpine life zone occurs between 1,200 and 1,600 feet; and the alpine life zone occurs above 1,600 feet.

Within each life zone, there are plant assemblages associated with cold and warm slopes. Slope temperature is a factor of slope steepness, aspect, and shading from surrounding ridges and mountains. Warm slopes occur on southeast to west aspects that are moderate to very steep and are not shaded by the surrounding landscape. Cold slopes occur on northwest to east facing slopes, occur in shaded slope positions, or occur in low-lying areas that are cold air sinks. Examples of shaded positions include head slopes, low relief backslopes of hills, and the base of hills and mountains shaded by adjacent mountain peaks. Warm boreal slopes have a cryic soil temperature regime and lack permafrost. White spruce forests are an indicator of warm boreal slopes. Cold boreal slope soils have a gelic soil temperature regime and commonly have permafrost. In this area, black spruce forests and woodlands are an indicator of cold boreal slopes.

## **Classification relationships**

Alaska Vegetation Classification:

Open needleleaf forest (I.A.2 - level III) / Open black spruce forest (I.A.2.f- level IV)  
(Vioreck et al., 1992)

Circumboreal Vegetation Map:

Alaska-Yukon Wet Black Spruce Woodlands and Scrub  
(Jorgensen and Meidinger, 2015)

BioPhysical Settings:

7316220 – Western North American Boreal Black Spruce Wet-Mesic Slope Woodland  
(Landfire, 2009)

## **Ecological site concept**

Ecological Site characteristics:

- Occurs in the boreal life zone on cold slopes of hills, mountains, plains, and terraces
- While soils are very deep, root-restricting permafrost occurs shallow to moderately deep depths

- Soils commonly formed in loess and colluvium and are capped with 9- to 13-inch-thick organic horizons
- Soils do not pond or flood. These poorly drained soils have a shallow water table throughout the growing season
- The reference plant community is an open black spruce forest with low shrubs, graminoids, and mosses dominating the understory
- Fire is the major disturbance in this ecological site and is responsible for three plant communities.

## Associated sites

R230XY620AK	<b>Boreal Peat Frozen Flats Complex</b> R230XY620AK describes the tussock tundra. They are located on the nearly level hill and mountain toeslopes down slope from F230XY611AK.
F230XY614AK	<b>Boreal Black Spruce Unfrozen Well Drained Slopes</b> Ecological site F230XY614AK occurs on the same hills and mountains but does not support permafrost in the soil.
F230XY615AK	<b>Boreal Deciduous Forest Loamy Steep Slopes</b> Ecological site F230XY615AK occurs on adjacent slopes that are much steeper and are associated with paper birch forests.
F230XY612AK	<b>Boreal Forest Loamy Slopes</b> Ecological site F230XY612AK describes white spruce forest on warm slopes. These ecological sites abut each other on the same slopes, but F230XY611AK is mostly restricted to north-facing slopes.

## Similar sites

F230XY614AK	<b>Boreal Black Spruce Unfrozen Well Drained Slopes</b> F230XY614AK and F230XY611AK both support black spruce forests. F230XY611AK is distinguished by poorly drained soils with permafrost which influences the vegetative community and fire dynamics.
-------------	---

**Table 1. Dominant plant species**

Tree	(1) <i>Picea mariana</i>
Shrub	(1) <i>Betula nana</i> (2) <i>Ledum palustre ssp. decumbens</i>
Herbaceous	(1) <i>Carex bigelowii</i>

## Physiographic features

- This boreal ecological site occurs on cold slopes of hills, mountains, plains, and terraces
- Elevation typically ranges between 100 and 1,200 feet
- Slope gradients are nearly level to steep (0 to 25 percent)

- This ecological site occurs on cold slopes. It is normally restricted to north-facing slopes, or slopes shaded by adjacent hills and mountains
- Flooding and ponding do not occur. A shallow water table (0 to 10 inches deep) is from May through September and often is present year-round.

**Table 2. Representative physiographic features**

Slope shape across	(1) Linear
Slope shape up-down	(1) Linear
Hillslope profile	(1) Backslope (2) Footslope
Geomorphic position, hills	(1) Side Slope
Geomorphic position, mountains	(1) Mountainbase (2) Mountainflank
Geomorphic position, terraces	(1) Tread
Landforms	(1) Hills > Plain (2) Mountains > Mountain (3) Hills > Hill (4) Valley > Terrace
Runoff class	Negligible to high
Flooding frequency	None
Ponding frequency	None
Elevation	30–366 m
Slope	0–25%
Water table depth	0–25 cm
Aspect	NW, N, NE, E


## Climatic features

The Yukon-Kuskokwim Highlands MLRA has short, warm summers and long, cold winters. Mean annual temperature ranges from 25 to 32 degrees Fahrenheit, with temperatures typically below freezing from October through April. June through August are the warmest months of the year and constitute the heart of the growing season. Approximately 60 percent of total annual precipitation occurs from June through September (PRISM, 2018; SNAP, 2014a). Across the MLRA, snowfall ranges from 80 to 100 inches (USDA, 2022).


**Table 3. Representative climatic features**

Frost-free period (characteristic range)	75-95 days
Freeze-free period (characteristic range)	65-85 days


Precipitation total (characteristic range)	330-508 mm
Frost-free period (actual range)	60-110 days
Freeze-free period (actual range)	50-100 days
Precipitation total (actual range)	254-1,016 mm
Frost-free period (average)	80 days
Freeze-free period (average)	70 days
Precipitation total (average)	381 mm




**Figure 1. Monthly precipitation range**




**Figure 2. Monthly minimum temperature range**




**Figure 3. Monthly maximum temperature range**



**Figure 4. Monthly average minimum and maximum temperature**



**Figure 5. Annual precipitation pattern**



**Figure 6. Annual average temperature pattern**

## Influencing water features

Precipitation and ground water are the main sources of water (Smith et al. 1995). Depth to the water table may decrease following summer storm events or spring snowmelt and increase during extended dry periods

## Wetland description

This ecological site is classified as a slope wetland in the Hydrogeomorphic (HGM) classification system (Smith et al. 1995; USDA-NRCS 2008).

## Soil features

- Common Gelisol great groups are Historthels and Histoturbels (Soil Survey Staff, 2013)
- Soils formed in loess over colluvium. Organic material is often intermixed with the mineral material due to cryoturbation.
- Rock fragments do not occur on the soil surface.
- Soils have a thick, 9- to 13-inch organic cap
- Soils are very deep but contain a shallow to moderately deep restrictive layer in the form of permafrost
- Subsurface rock fragments are generally absent except where cryoturbation has occurred
- Soil pH is extremely to moderately acidic near the surface and strongly acidic to neutral in permafrost
- Soils are poorly drained

**Table 4. Representative soil features**

Parent material	(1) Organic material (2) Loess (3) Eolian deposits (4) Colluvium
-----------------	---

Surface texture	(1) Peat (2) Silt loam
Drainage class	Poorly drained
Permeability class	Moderate
Depth to restrictive layer	25–66 cm
Soil depth	152 cm
Surface fragment cover ≤3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	21.59–28.96 cm
Soil reaction (1:1 water) (0-25.4cm)	3.9–5.7
Subsurface fragment volume ≤3" (0-152.4cm)	0%
Subsurface fragment volume >3" (0-152.4cm)	0%

**Table 5. Representative soil features (actual values)**

Drainage class	Not specified
Permeability class	Not specified
Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover ≤3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	Not specified
Soil reaction (1:1 water) (0-25.4cm)	Not specified
Subsurface fragment volume ≤3" (0-152.4cm)	0–20%
Subsurface fragment volume >3" (0-152.4cm)	0–12%

## Ecological dynamics

Fire

Fire has a major influence on vegetation across the Yukon-Kuskokwim Highlands MLRA. Wildland fire is a common and natural event that is unmanaged except near towns and villages. Fires are patchy and different locations within a single burn experience different levels of fire intensity, or no fire at all. In general, cooler and wetter areas experience low-severity fires, while warmer and drier areas experience high-severity fires. Because the soils have a thick organic cap and are poorly drained, the typical fire scenario for this ecological site is considered to result in a low severity burn. Other factors such as fire load and recent weather also affect fire characteristics. A typical fire event considerably alters vegetation, above ground biomass, and the organic soil cap.

Fire is the major natural disturbance for this ecological site. The fire return interval ranges between 50 and 150 years, with an average of less than 100 years (Viereck et al., 1992; Fryer, 2014). Fire is facilitated by a high fuel load, flammable chemical contents within the plant, and the multilayer community structure of black spruce (Landfire, 2009; Fryer, 2014). Ericaceous shrubs in the lower canopy act as ladder fuels. Black spruce forests are susceptible to all fire severity levels. This tree species is fire adapted, and seed release is facilitated by fire. Post-fire seeding rates are exceptionally high, and reestablishment occurs within five years (Fryer, 2014).


Low-severity fires have modest impacts on soil organic matter thickness, depth to permafrost, and soil drainage. While a low-severity fire can consume the bulk of above ground vegetation, minimal proportions of the organic mat are typically removed. Organic matter continues to insulate these cold soils. If permafrost remains at similar depths after a fire event, then soil drainage is unlikely to improve post-fire.

The normal fire cycle in black spruce is a stand-replacement event. There is no single pathway regarding boreal forest fire dynamics; post-fire communities depend on various factors including pre-fire community composition, fire severity, weather, and the presence/absence of permafrost (Fryer, 2014). In general, a typical post-fire successional pathway begins with an herbaceous meadow before progressing through a low shrubland and then onto a black spruce woodland or forest.


The information in this Ecological Dynamics section, including the state-and-transition model (STM), was developed based on current field data, professional experience, and a review of the scientific literature. As a result, all possible scenarios or plant species may not be included. Key indicator plant species, disturbances, and ecological processes are described to inform land management decisions.

## **State and transition model**

## Ecosystem states



## State 1 submodel, plant communities



**1.1a** - Fire; occurs 0-19 years after fire

**1.2a** - Fire recovery; occurs 40+ years after fire

**1.2b** - Fire

**1.3a** - Fire recovery; occurs 20 to 40 years after fire

## State 1 Reference State



The reference state describes three distinct vegetative communities on cold, boreal slopes with permafrost. Fire is the major disturbance. Factors such as a cyclical fire regime, soil acidity and soil drainage contribute to the community composition and dynamics of this ecological site. The reference state is developed and characterized using available vegetation models, including Landfire BpS and the Alaska vegetation classification system (Landfire, 2009; Viereck et al., 1992).

### **Dominant plant species**

- black spruce (*Picea mariana*), tree
- dwarf birch (*Betula nana*), shrub
- marsh Labrador tea (*Ledum palustre ssp. decumbens*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- Bigelow's sedge (*Carex bigelowii*), grass
- tussock cottongrass (*Eriophorum vaginatum*), grass
- sphagnum (*Sphagnum*), other herbaceous
- splendid feather moss (*Hylocomium splendens*), other herbaceous

### **Community 1.1**

***Picea mariana* / *Betula nana* - *Ledum palustre ssp. decumbens* / *Carex bigelowii***



The reference plant community is an open black spruce forest (Viereck et al. 1992). White spruce may be present on slightly warmer slopes. Tree cover is primarily in the medium tree stratum (between 15 and 40 feet in height). The dominant vegetative strata are medium trees, stunted trees, low shrubs, and mosses. The soil surface is primarily covered with mosses, lichens, and herbaceous litter. The binomial and vernacular name of common plants are listed in the dominant plant species table.

### **Dominant plant species**

- black spruce (*Picea mariana*), tree
- dwarf birch (*Betula nana*), shrub
- marsh Labrador tea (*Ledum palustre ssp. decumbens*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- tealeaf willow (*Salix pulchra*), shrub
- netleaf willow (*Salix reticulata*), shrub
- red fruit bearberry (*Arctostaphylos rubra*), shrub

- Bigelow's sedge (*Carex bigelowii*), grass
- cottongrass (*Eriophorum*), grass
- bluejoint (*Calamagrostis canadensis*), grass
- wideleaf polargrass (*Arctagrostis latifolia*), grass
- arctic bluegrass (*Poa arctica*), grass
- horsetail (*Equisetum*), other herbaceous
- cloudberry (*Rubus chamaemorus*), other herbaceous
- sphagnum (*Sphagnum*), other herbaceous
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- reindeer lichen (*Cladina*), other herbaceous

## Community 1.2

### ***Picea mariana* / *Betula nana* - *Ledum palustre* ssp. *decumbens* / *Carex bigelowii***

This community is a black spruce woodland (Viereck et al. 1992). The dominant vegetative strata are medium and regenerative trees, low and dwarf shrubs, medium graminoids, and mosses. Black spruce is a mix of saplings and young mature trees. The understory is a dense mix of birch and ericaceous shrubs. Willows such as tealeaf willow may be present. Ground cover is primarily mosses and herbaceous litter. The binomial and vernacular name of common plants are listed in the dominant plant species table.

#### **Dominant plant species**

- black spruce (*Picea mariana*), tree
- dwarf birch (*Betula nana*), shrub
- marsh Labrador tea (*Ledum palustre* ssp. *decumbens*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- tealeaf willow (*Salix pulchra*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- Bigelow's sedge (*Carex bigelowii*), grass
- Altai fescue (*Festuca altaica*), grass
- horsetail (*Equisetum*), other herbaceous
- sphagnum (*Sphagnum*), other herbaceous
- splendid feather moss (*Hylocomium splendens*), other herbaceous

## Community 1.3

### ***Betula nana* - *Ledum palustre* ssp. *decumbens* / *Carex* spp. - *Equisetum sylvaticum***



This community is an open low scrubland (Viereck et al. 1992). The dominant vegetative strata depend on fire disturbance dynamics, but typically include low shrubs, medium graminoids, and tall and medium forbs. Extant shrub species from the pre-fire community may be present, as may black spruce seedlings. Ground cover is primarily fresh and burned herbaceous litter, woody litter, and bare soil. The binomial and vernacular name of common plants are listed in the dominant plant species table.

### Dominant plant species

- marsh Labrador tea (*Ledum palustre ssp. decumbens*), shrub
- dwarf birch (*Betula nana*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- Bigelow's sedge (*Carex bigelowii*), grass
- bluejoint (*Calamagrostis canadensis*), grass
- sedge (*Carex*), grass
- horsetail (*Equisetum*), other herbaceous
- fireweed (*Chamerion angustifolium*), other herbaceous

### Pathway 1.1a

#### Community 1.1 to 1.3



*Picea mariana* / *Betula nana* -  
*Ledum palustre ssp. decumbens* / *Carex bigelowii*



*Betula nana* - *Ledum palustre ssp. decumbens* / *Carex spp.* -  
*Equisetum sylvaticum*

A fire sweeps through and incinerates much of the above ground vegetation. Because of

the associated cold and wet soils, this site commonly experiences low-severity fires. Minimal proportions of the organic mat are typically removed. The pre-fire vegetation generally reestablishes quickly from below ground root systems and rhizomes. This community develops immediately after a fire and persists for up to 20 years (Landfire, 2009).

### **Pathway 1.2a**

#### **Community 1.2 to 1.1**

Black spruce seedlings and saplings mature into a woodland. This community develops 40 or more years post-wildfire (Landfire, 2009). Total herbaceous cover drops as competition from spruce and shrubs increases. The organic mat made up of mosses grows and thickens, providing more insulation for permafrost to develop closer to the surface.

### **Pathway 1.2b**

#### **Community 1.2 to 1.3**

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated cold and wet soils, this site commonly experiences low-severity fires. Minimal proportions of the organic mat are typically removed. The pre-fire vegetation generally reestablishes quickly from below ground root systems and rhizomes.

### **Pathway 1.3a**

#### **Community 1.3 to 1.2**

Shrubs colonize the community either via rootstock or seeding. Black spruce seedlings and saplings begin to establish. This community develops 20 to 40 years after fire (Landfire, 2009).

## **Additional community tables**

### **Inventory data references**

Vegetative communities and transitions are described using existing models and expert knowledge. There are no vegetation inventory data points in NASIS associated with this ecological site.

External model data sources:

The Alaska-Yukon Region of the Circumboreal Vegetation Map (CBVM) (Jorgensen and Meidinger, 2015)

LANDFIRE Biophysical Settings Models (Landfire, 2009)

Viereck, L.A., C.T. Dyrness, A.R. Batten, and K.J. Wenzlick. 1992. The Alaska vegetation

classification. Gen. Tech. Rep. PNW-GTR-286. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 278 p. (Viereck et al., 1992)

## References

Vioreck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

## Other references

Fryer, Janet L. 2014. *Picea mariana*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <https://www.fs.usda.gov/database/feis/plants/tree/picmar/all.html> [2025, March 26].

Jorgensen, T., and D. Meidinger. 2015. The Alaska Yukon Region of the Circumboreal Vegetation Map (CBVM). CAFF Strategies Series Report. Conservation of Arctic Flora and Fauna, Akureyri, Iceland. ISBN: 978-9935-431-48-6.

Landfire. 2009. Biophysical Setting 7316220 – Western North American Boreal Black Spruce Wet-Mesic Slope Woodland. In: LANDFIRE Biophysical Setting Model: Map zone 73, [Online]. In: Vegetation Dynamics Models. In: LANDFIRE. Washington, DC: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory; U.S. Geological Survey; Arlington, VA: The Nature Conservancy (Producers). Available: <https://www.landfire.gov/vegetation/bps>. Accessed March 18, 2025

PRISM Climate Group (PRISM). 2018. “Alaska – average monthly and annual precipitation and minimum, maximum, and mean temperature for the period 1981-2010.” Oregon State University, Corvallis, Oregon. <https://prism.oregonstate.edu/projects/alaska.php>. Accessed Sept 17, 2024.

Scenarios network for Alaska and arctic planning (SNAP). 2014. “Historical Monthly Temperature – 1km, 1901-2009”. <http://ckan.snap.uaf.edu/dataset/>. Accessed Sept 17, 2024.

Scenarios network for Alaska and arctic planning (SNAP). 2014. “Historical monthly and derived precipitation products downscaled from CRU TS data via the delta methods – 2km, 1901-2009”. <http://ckan.snap.uaf.edu/dataset/>. Accessed Sept 17, 2024.

Soil Survey Staff. 2013. Simplified Guide to Soil Taxonomy. USDA-Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE.

United States Department of Agriculture, Natural Resources Conservation Service. 2022. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

Western Regional Climate Center (WRCC). 2021. Climate of Alaska. Retrieved from [https://wrcc.dri.edu/Climate/narrative\\_ak.php](https://wrcc.dri.edu/Climate/narrative_ak.php). Accessed November 15, 2024.

Western Regional Climate Center (WRCC). 2025. "McGrath WB Airport, Alaska 'Freeze Free' Season Probabilities." <https://wrcc.dri.edu/cgi-bin/cliTFrezD.pl?akmcgr>. Accessed Mar 6, 2025.

## Contributors

Phil Barber  
Blaine Spellman  
Marji Patz  
Claire Benton

## Acknowledgments

This ecological site description (ESD) fulfills the requirements of the Provisional Ecological Site (PES) national initiative. This ESD is published to fit current site-soil correlations as they are currently mapped and understood. Further data collection may provide the information to update this ESD from the provisional level to the approved level.

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	03/28/2026
Approved by	Blaine Spellman
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

# Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile**

features which may be mistaken for compaction on this site):

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 

14. **Average percent litter cover (%) and depth ( in):**
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 

17. **Perennial plant reproductive capability:**
-

