

## Ecological site R231XY105AK Alpine Dwarf Scrub Gravelly Alkaline Slopes

Last updated: 2/13/2024  
Accessed: 04/23/2024

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 231X–Interior Alaska Highlands

The Interior Alaska Uplands (MLRA 231X) is in the Interior Region of Alaska and includes the extensive hills, mountains, and valleys between the Tanana River to the south and the Brooks Range to the north. These hills and mountains surround the Yukon Flats Lowlands (MLRA 232X). MLRA 231X makes up about 69,175 square miles. The hills and mountains of the area tend to be moderately steep to steep resulting in high-relief slopes. The mountains are generally rounded at lower elevations and sharp-ridged at higher elevations. Elevation ranges from about 400 feet in the west, along the boundary with the Interior Alaska Lowlands (MLRA 229X), to 6,583 feet at the summit of Mt. Harper, in the southeast. Major tributaries include large sections of the Yukon, Koyukuk, Kanuti, Charley, Coleen, and Chatanika Rivers. This area is traversed by several major roads, including the Taylor Highway in the east and the Steese, Elliott, and Dalton Highways north of Fairbanks. The area is mostly undeveloped wild land that is sparsely populated. The largest community along the road system is Fairbanks with smaller communities like Alatna, Allakaket, Chicken, Eagle, Eagle Village, Hughes, and Rampart occurring along the previously mentioned rivers and highways.

The vast majority of this MLRA was unglaciated during the Pleistocene epoch with the exceptions being the highest mountains and where glaciers extended into the area from the Brooks Range. For the most part, glacial moraines and drift are limited to the upper elevations of the highest mountains. Most of the landscape is mantled with bedrock colluvium originating from the underlying bedrock. Valley bottoms are filled with Holocene fluvial deposits and colluvium from the adjacent mountain slopes. Silty loess, which originated from unvegetated flood plains in and adjacent to this area, covers much of the surface. On hill and mountain slopes proximal to major river valleys (e.g., Tanana and Yukon Rivers), the loess is many feet thick. As elevation and distance from major river valleys increases, loess thickness decreases significantly. Bedrock is commonly exposed on the highest ridges.

This area is in the zone of discontinuous permafrost. Permafrost commonly is close to the surface in areas of the finer textured sediments throughout the MLRA. Isolated masses of ground ice occur in thick deposits of loess on terraces and the lower side slopes of hills. Solifluction lobes, frost boils, and circles and stripes are periglacial features common on mountain slopes in this area. Pingos, thermokarst pits and mounds, ice-wedge polygons, and earth hummocks are periglacial features common on terraces, lower slopes of hills and mountains, and in upland valleys in the area.

The dominant soil orders in this area are Gelisols, Inceptisols, Spodosols, and Entisols. The soils in the area have a subgelic or cryic soil temperature regime, an aquic or udic soil moisture regime, and mixed mineralogy. Gelisols are common on north facing slopes, south facing footslopes, valley bottoms, and stream terraces. Gelisols are typically shallow or moderately deep to permafrost (10 to 40 inches) and are poorly or very poorly drained. Wildfires can disturb the insulating organic material at the surface, lowering the permafrost layer, eliminating perched water tables from Gelisols, and thus changing the soil classification. Inceptisols and Spodosols commonly form on south facing hill and mountain slopes. Entisols are common on flood plains and high elevation mountain slopes. Miscellaneous (non-soil) areas make up about 2 percent of this MLRA. The most common miscellaneous areas are rock outcrop and rubble land. In many valleys placer mine tailings are common.

Short, warm summers and long, cold winters characterize the subarctic continental climate of the area. The mean annual temperature of the area ranges from 22 to 27 degrees F. The mean annual temperature of the southern half of the area is approximately 3 degrees warmer compared to the northern half (PRISM 2018). The warmest months span June through August with mean monthly temperatures ranging from 50 to 56 degrees F. The coldest months span November through February with mean monthly temperatures ranging from -5 to 3 degrees F. When compared to the high-elevation alpine and subalpine life zones, the lower elevation boreal life zone tends to be 2-3 degrees F colder during the coldest months and 1-2 degrees F warmer during the warmest months (PRISM 2018). The freeze-free period at the lower elevations averages about 60 to 100 days, and the temperature usually remains above freezing from June through mid-September.

Precipitation is limited across this area, with the average annual precipitation ranging from 12 to 19 inches. The southern half of the areas receives approximately 2.5 inches more annual precipitation than the northern half (PRISM 2018). The lower elevation boreal life zone receives approximately 2.5 inches less annual precipitation than the high-elevation alpine and subalpine life zones (PRISM 2018). Approximately 3/5th of the annual precipitation occurs during the months of June through September with thunderstorms being common. The average annual snowfall ranges from about 45 to 100 inches. The ground is consistently covered with snow from November through March.

Most of this area is forested below an elevation of about 2500 feet. Dominant tree species on slopes are white spruce and black spruce. Black spruce stands are most common on north-facing slopes, stream terraces, and other sites with poor drainage and permafrost. White spruce stands are most common on warm slopes with dry soils. At lower elevations, lightning-caused wildfires are common, often burning many thousands of acres during a single fire. Following wildfires, forbs, grasses, willow, ericaceous shrubs, paper birch, and quacking aspen communities are common until they are eventually replaced by stands of spruce. Tall willow and alder scrub is extensive on low flood plains. White spruce and balsam poplar are common on high flood plains.

With increasing elevation, the forests and woodlands give way to subalpine communities dominated by krummholz spruce, shrub birch, willow, and ericaceous shrubs. At even higher elevations, alpine communities prevail which are characterized by diverse forbs, dwarf ericaceous shrubs, and eightpetal mountain-avens. Many of these high elevation communities have a considerable amount of lichen cover and bare ground.

## **LRU notes**

This area supports three life zones defined by the physiological limits of plant communities along an elevational gradient: boreal, subalpine, and alpine. The boreal life zone is the elevational band where forest communities dominate. Not all areas in the boreal life zone are forest communities, however, particularly in places with too wet or dry soil to support tree growth (e.g., bogs or river bluffs). Above the boreal band of elevation, subalpine and alpine vegetation dominate. The subalpine zone is typically a narrow transitional band between the boreal and the alpine life zones, and is characterized by sparse, stunted trees. In the subalpine, certain types of birch and willow shrub species grow at  $\geq 1$  m in height (commonly *Betula glandulosa* and *Salix pulchra*). In the alpine, trees no longer occur, and all shrubs are dwarf or lay prostrate on the ground. In this area, the boreal life zone occurs below 2500 feet elevation on average. The transition between boreal and alpine vegetation can occur within a range of elevations, and is highly dependent on slope, aspect, and shading from adjacent mountains.

Within each life zone, there are plant assemblages that are typically associated with cold slopes and warm slopes. Cold slopes and warm slopes are created by the combination of the steepness of the slope, the aspect, and shading from surrounding ridges and mountains. Warm slope positions typically occur on southeast to west facing slopes that are moderate to very steep ( $>10\%$  slope) and are not shaded by the surrounding landscape. Cold slopes typically occur on northwest to east facing slopes, occur in shaded slope positions, or occur in low-lying areas that are cold air sinks. Examples of shaded positions include head slopes, low relief backslopes of hills, and the base of hills and mountains shaded by adjacent mountain peaks. Warm boreal slope soils have a cryic soil temperature regime and lack permafrost. In this area, white spruce forests are an indicator of warm boreal slopes. Cold boreal slope soils typically have a gelic soil temperature regime and commonly have permafrost. In this area, black spruce forests and woodlands are an indicator of cold boreal slopes. The boreal life zone can occur at higher elevations on warm slopes, and lower elevations on cold slopes.

## Classification relationships

Landfire BPS – 7416310 – Western North American Boreal Alpine Dwarf-Shrub Summit

## Ecological site concept

This site occurs on warm alpine slopes with dry, gravelly, alkaline soils. This site is associated with the summits and backslopes of limestone mountains at high elevation. Circles and solifluction lobes are periglacial features occasionally associated with this site. The soils lack permafrost, do not have a water table during the growing season, and are considered well drained. The soils formed in silty loess and gravelly and alkaline colluvium. The pH of the colluvium commonly ranges from neutral to moderately alkaline.

The alpine life zone has a harsh climate that limits growth of vegetation and prevents the establishment of many species common at lower elevations. In this area, alpine vegetation is characterized as dwarf and prostrate shrubs intermixed with low-lying herbaceous plants. These unique plant communities are the result of high winds, a short growing season, deep and persistent snow beds, and cold soils. These climatic factors prevent the establishment and growth of many dominant boreal species like white spruce and black spruce.

Two plant communities occur within the reference state and the vegetation in each community differs in large part due to fire. When the reference state vegetation burns, the post-fire plant community is dominantly graminoids, forbs, and weedy mosses. With time and lack of another fire event, the post-fire vegetation goes through succession. For this site, the reference plant community is the most stable with the longest time since the vegetation was burned. This community is typically characterized as dryas dwarf scrub (Viereck et al. 1992) and is highly diverse. Common species include eight-petal mountain-avens, arctic willow, white arctic mountain heather, netleaf willow, northern singlespike sedge, various fescue, curled snow lichen, and crinkled snow lichen. The vegetative strata that characterize this community are dwarf shrubs (less than 8 inches), medium graminoids (between 4 and 24 inches), and foliose and fruticose lichen.

## Associated sites

R231XY103AK	<b>Alpine Dwarf Scrub Gravelly Frozen Alkaline Slopes</b> Occurs on colder alpine slopes with wet soils that have permafrost.
R231XY104AK	<b>Alpine Dwarf Scrub Gravelly Alkaline Cold Slopes</b> Occurs on colder alpine slopes with dry, gravelly soils that lack permafrost.
R231XY106AK	<b>Alpine Dwarf Scrub Gravelly Frozen Alkaline Slopes</b> Occurs on colder alpine slopes with wet and gravelly soils that do not have permafrost.
R231XY152AK	<b>High-elevation scrub gravelly drainageways</b> Occurs downslope in high elevation drainageways.

## Similar sites

R231XY101AK	<b>Alpine dwarf scrub gravelly slopes</b> Both sites occur on alpine slopes with dry and gravelly soils. Site 105 has alkaline soils resulting in different kinds and amounts of vegetation.
R231XY104AK	<b>Alpine Dwarf Scrub Gravelly Alkaline Cold Slopes</b> Both sites occur on the same mountains on dry, alkaline soils. Site 104 occurs on colder slopes with snowpack that persists for longer durations of time. These differences result in different kinds and amounts of vegetation.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Dryas octopetala</i> ssp. <i>octopetala</i> (2) <i>Salix arctica</i>
Herbaceous	(1) <i>Carex scirpoidea</i> (2) <i>Flavocetraria cucullata</i>

## Physiographic features

This alpine site occurs on limestone hills and mountains at high elevation. This site is associated with summits, shoulders, and backslopes of hills and mountains. Sorted circles, nonsorted circles, stripes, and solifluction lobes (Schoeneberger and Wysocki 2017) are periglacial microfeatures occasionally associated on mountain slopes. Elevation typically ranges between 2500 and 3000 feet but can go as low as 1800 feet on certain windswept hill summits and as high as 4925 feet on the largest limestone mountains in the area. This site occurs on steep and warm slopes that are southeast to west facing. During the growing season, a water table does not typically occur in the soil profile. This site does not experience flooding or ponding, but rather generates moderate runoff to adjacent, downslope ecological sites.



Figure 1. A sorted rock stripe associated with this site.

Table 2. Representative physiographic features

Hillslope profile	(1) Summit (2) Shoulder (3) Backslope
Landforms	(1) Mountains > Mountain slope (2) Mountains > Hill (3) Mountains > Mountain slope > Nonsorted circle (4) Mountains > Mountain slope > Circle (5) Mountains > Mountain slope > Stripe (6) Mountains > Mountain slope > Solifluction lobe
Runoff class	Medium
Flooding frequency	None
Ponding frequency	None
Elevation	2,500–3,000 ft
Slope	20–45%
Water table depth	60 in
Aspect	W, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	1,800–4,950 ft
Slope	18–75%

Water table depth	40 in
-------------------	-------

## Climatic features


When compared to the boreal life zone, this high-elevation site has a harsh climate. In this MLRA, snow first blankets and persists the longest in the alpine and subalpine life zones. From spring through fall (April through September), it is consistently 1 to 2 degrees F colder in the alpine and subalpine. These small differences in temperature are exacerbated due to constant and strong winds. Winds are much more intense in these high elevation areas because of limited trees providing windbreaks. When compared to the boreal life zone, this site has a much shorter growing season and the growing season is significantly colder for associated vegetation.

Short, warm summers and long, cold winters characterize the subarctic continental climate associated with this high-elevation site. The mean annual temperature of the site ranges from 23 to 27 degrees F. The warmest months span June through August with mean normal maximum monthly temperatures ranging from 57 to 63 degrees F. The coldest months span November through February with mean normal minimum temperatures ranging from -9 to -1 degrees F. The freeze-free period for the site ranges from 80 to 120 days, and the temperature usually remains above freezing from late May through mid-September.


The area receives minimal annual precipitation with the summer months being the wettest. Average annual precipitation in the alpine across the area typically ranges between 14 to 21 inches. Approximately 3/5th of the annual precipitation occurs during the months of June through September with thunderstorms common. The average annual snowfall ranges from about 45 to 100 inches. The ground is consistently covered with snow from mid-October through March.

**Table 4. Representative climatic features**


Frost-free period (characteristic range)	16-78 days
Freeze-free period (characteristic range)	76-114 days
Precipitation total (characteristic range)	14-21 in
Frost-free period (actual range)	4-87 days
Freeze-free period (actual range)	48-120 days
Precipitation total (actual range)	10-25 in
Frost-free period (average)	53 days
Freeze-free period (average)	90 days
Precipitation total (average)	17 in




**Figure 2. Monthly precipitation range**




**Figure 3. Monthly minimum temperature range**



**Figure 4. Monthly maximum temperature range**



**Figure 5. Monthly average minimum and maximum temperature**



**Figure 6. Annual precipitation pattern**




Figure 7. Annual average temperature pattern

## Climate stations used

- (1) EAGLE AP [USW00026422], Tok, AK
- (2) CHICKEN [USC00501684], Tok, AK
- (3) MILE 42 STEESE [USC00505880], Fairbanks, AK
- (4) BETTLES AP [USW00026533], Bettles Field, AK
- (5) CIRCLE HOT SPRINGS [USC00501987], Central, AK
- (6) FT KNOX MINE [USC00503160], Fairbanks, AK
- (7) GILMORE CREEK [USC00503275], Fairbanks, AK
- (8) FOX 2SE [USC00503181], Fairbanks, AK
- (9) ESTER DOME [USC00502868], Fairbanks, AK
- (10) ESTER 5NE [USC00502871], Fairbanks, AK
- (11) COLLEGE 5 NW [USC00502112], Fairbanks, AK
- (12) COLLEGE OBSY [USC00502107], Fairbanks, AK
- (13) KEYSTONE RIDGE [USC00504621], Fairbanks, AK

## Influencing water features

Due to its landscape position, this site is neither associated with or influenced by streams or wetlands. Precipitation and throughflow are the main source of water for this ecological site. Surface runoff and throughflow contribute some water to downslope ecological sites.

## Wetland description

n/a

## Soil features

Soils formed in windblown silt over gravelly parent material and do not have permafrost. Surface rock fragments are common and typically range between 5 and 50 percent cover. These are mineral soils often capped with 0 to 2 inches of organic material. The mineral soil below the organic material is a silt loam formed from wind-blown loess, which is at times mixed with some rock fragments due to cryoturbation, and has higher water holding capacity. The thickness of this loess layer is variable ranging from 0 to 4 inches. Below this loess layer is gravelly colluvium with rock fragments that typically range between 30 and 60 percent of the soil profile by volume and has lower water holding capacity. Soils are often very deep without restrictions. At times, soils with extremely gravelly colluvium have strongly contrasting textural stratification at very shallow depths (2 to 8 inches) and/or contract bedrock at moderate to very deep depths (40 to 60 inches). The pH of the soil profile typically ranges from neutral to moderately alkaline. The soils are dry for the growing season and are considered well drained.



Figure 8. A typical soil profile associated with this site.

Table 5. Representative soil features

Parent material	(1) Loess (2) Eolian deposits (3) Colluvium–limestone and dolomite
Surface texture	(1) Silt loam (2) Very gravelly silt loam (3) Very channery silt loam
Family particle size	(1) Loamy-skeletal
Drainage class	Well drained
Permeability class	Moderately rapid to very rapid
Depth to restrictive layer	40 in
Soil depth	40 in
Surface fragment cover ≤3"	5–50%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	1.5–4.1 in
Calcium carbonate equivalent (10-40in)	0–12%
Clay content (0-20in)	5–10%
Electrical conductivity (10-40in)	0–9 mmhos/cm
Sodium adsorption ratio (10-40in)	0
Soil reaction (1:1 water) (10-40in)	6.7–8.3
Subsurface fragment volume ≤3" (0-60in)	15–25%
Subsurface fragment volume >3" (0-60in)	15–35%

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Permeability class	Not specified



Depth to restrictive layer	2 in
Soil depth	Not specified
Surface fragment cover <=3"	0–50%
Surface fragment cover >3"	0–25%
Available water capacity (0-40in)	1–4.5 in
Calcium carbonate equivalent (10-40in)	Not specified
Clay content (0-20in)	Not specified
Electrical conductivity (10-40in)	Not specified
Sodium adsorption ratio (10-40in)	0–2
Soil reaction (1:1 water) (10-40in)	5.6–8.3
Subsurface fragment volume <=3" (0-60in)	10–50%
Subsurface fragment volume >3" (0-60in)	10–40%

## Ecological dynamics

### Climate

Located in the alpine life zone, this site is exposed to a variety of harsh environmental conditions. In this area, snowfall first appears and persists the longest in the alpine. As a result, snowpack tends to be deeper and persist for longer durations of time compared to lower-elevation sites and alpine vegetation has a comparatively shorter growing season. When this site is snow-free, cold soil temperatures and high winds also inhibit plant growth and vigor. This harsh climate maintains the dwarfed vegetation within this site and prevents the establishment and/or growth of dominant boreal species like white spruce and black spruce.

### Fire

Within this area, fire is considered a natural and common event that typically is unmanaged. Fire suppression is limited, and generally occurs adjacent to Fairbanks and the various villages spread throughout the area or on allotments with known structures, all of which have a relatively limited acre footprint. Most fires are caused by lightning strikes. From 2000 to 2020, 596 known fire events occurred in the Interior Alaska Uplands area and the burn perimeter of the fires totaled about 13.8 million acres (AICC 2022). Fire-related disturbances are highly patchy and can leave undisturbed areas within the burn perimeter. During this time frame, 80% of the fire events were smaller than 20,000 acres but 18 fire events were greater than 200,000 acres in size (AICC 2022). Over this period of 20 years, these burn perimeters cover approximately 30 percent of the Interior Alaska Uplands area.

The fire regime within Interior Alaska follows two general scenarios—low-severity burns and high-severity burns. It should be noted, however, that the fire regime in Interior Alaska can be much more complex (Johnstone et al. 2008). Burn severity refers to the proportion of the vegetative canopy and organic material consumed in a fire event (Chapin et al. 2006). Fires in cool and moist habitat tend to result in low-severity burns, while fires in warm and dry habitat tend to result in high-severity burns. Because the soils have a thin organic cap and are well drained, the typical fire scenario for this ecological site is considered to result in a high severity burn.


Large portions of the organic mat are consumed during a high-severity fire event, commonly exposing pockets of mineral soil. The loss of this organic mat, which insulates the mineral soil, and the decrease in site albedo tends to cause overall soil temperatures to increase (Hinzman et al. 2006). These alterations to soil temperature may result in increased depths of seasonal frost in the soil profile. High-severity fire events also destroy a majority of the

vascular and nonvascular biomass above ground.


Field data suggest that the scrub dominant community 1.1 burns and that fire events will cause a transition to the pioneering stage of fire succession. This stage (community 1.2) is a mix of species that either regenerate in place (e.g., subterranean root crowns for willow and rhizomes for graminoids) and/or from wind-dispersed seed or spores that colonize exposed mineral soil (e.g., fireweed [*Chamerion angustifolium*] and Ceratodon moss [*Ceratodon purpureus*]). The pioneering stage of fire succession is primarily composed of dwarf scrubs, grasses, forbs, grasses, and weedy bryophytes. This stage of succession currently persists for an unknown amount of time but is thought to last 10 to 30 years post-fire. Eightpetal mountain-avens and other dwarf shrubs, graminoids, and lichen continue to colonize and grow until they become dominant in the plant community, which marks the transition to the reference plant community (community 1.1).

## State and transition model

### Ecosystem states



### State 1 submodel, plant communities



**1.1a** - A high-severity fire sweeps through and incinerates much of the above ground vegetation.

**1.2a** - Time without fire.

## State 1 Reference State



**Figure 9.** A dwarf scrub community associated with this site.

The reference plant community is dryas dwarf scrub (Viereck et al. 1992). There are two plant communities in the reference state related to fire. Solifluction is a process associated with this state but does not happen to a degree resulting in a mosaic of vegetation. Solifluction is the slow, viscous downslope flow of water-saturated soil (Shoeneberger and Wysocki 2017). This process is most active for this site during spring thaw where the upper band of soil material slips on a seasonally frozen layer. Solifluction is a common process associated with several ecological sites in this area and this site can at times have small solifluction lobes. Since these small solifluction

lobes are uncommon and do not typically result in a vegetation mosaic, no alternative solifluction state was developed for this site (see R231XY113AK for a site that does have this alternate state). Cryoturbation is process associated with this state that results in the formation of non-sorted circles. Cryoturbation is a collective term used to describe all soil movements due to frost action, characterized by folded, broken and dislocated beds and lenses of unconsolidated deposits (Schoeneberger and Wysocki 2017). Since non-sorted circles are uncommon for this site and data did not typically support these features having a vegetation mosaic, no alternative state was developed for this site (see R231XY134AK for a site that does have this alternate state).

### Dominant plant species

- eightpetal mountain-avens (*Dryas octopetala ssp. octopetala*), shrub
- white arctic mountain heather (*Cassiope tetragona*), shrub
- northern singlespike sedge (*Carex scirpoidea*), grass
- fescue (*Festuca*), grass
- (*Flavocetraria cucullata*), other herbaceous
- whiteworm lichen (*Thamnolia vermicularis*), other herbaceous

### Community 1.1

**eightpetal mountain-avens - arctic willow / northern singlespike sedge - fescue / curled snow lichen - whiteworm lichen**



Figure 10. A typical plant community associated with community 1.1.

The reference plant community is characterized as dryas dwarf scrub (Viereck et al. 1992) and is highly diverse. Stunted white spruce occasionally occur but have limited cover. Common species include eight-petal mountain-avens, arctic willow, white arctic mountain heather, netleaf willow, northern singlespike sedge, various fescue, curled snow lichen (*Flavocetraria cucullata*), and crinkled snow lichen (*F. nivalis*). The vegetative strata that characterize this community are dwarf shrubs (less than 8 inches), medium graminoids (between 4 and 24 inches), and foliose and fruticose lichen. The soil surface is primarily covered with herbaceous litter, surface rock fragments, lichens, and mosses.

**Forest understory.** The mountain-avens most commonly associated with this site is *Dryas ajanensis* (DROCO). The Flora of North America no longer recognizes eightpetal mountain-avens (*Dryas octopetala*) as occurring in Alaska and has split this species concept into several new species (Springer and Parfitt 2015).

### Dominant plant species

- eightpetal mountain-avens (*Dryas octopetala ssp. octopetala*), shrub
- arctic willow (*Salix arctica*), shrub
- white arctic mountain heather (*Cassiope tetragona*), shrub
- netleaf willow (*Salix reticulata*), shrub
- northern singlespike sedge (*Carex scirpoidea*), grass
- Altai fescue (*Festuca altaica*), grass
- alpine fescue (*Festuca brachyphylla*), grass
- (*Flavocetraria cucullata*), other herbaceous
- (*Flavocetraria nivalis*), other herbaceous

## Community 1.2

northern singlespike sedge - fescue / fireweed / Ceratodon moss - juniper Polytrichum moss



Figure 11. A typical plant community associated with community 1.2.


Community 1.2 is in the pioneering stage of fire-induced secondary succession for this ecological site. It is characterized as either a mesic forb herbaceous or mesic graminoid herbaceous community (Viereck et al. 1992). Commonly observed species include eightpetal mountain-avens, netleaf willow, northern singlespike sedge, Altai fescue, bluegrass, fireweed, Ceratodon moss, and juniper Polytrichum moss. The soil surface is primarily covered with a mixture of weedy bryophyte species and surface rock fragments.

### Dominant plant species

- eightpetal mountain-avens (*Dryas octopetala* ssp. *octopetala*), shrub
- netleaf willow (*Salix reticulata*), shrub
- northern singlespike sedge (*Carex scirpoidea*), grass
- Altai fescue (*Festuca altaica*), grass
- bluegrass (*Poa*), grass
- fireweed (*Chamerion angustifolium*), other herbaceous
- ceratodon moss (*Ceratodon purpureus*), other herbaceous
- juniper polytrichum moss (*Polytrichum juniperinum*), other herbaceous

### Pathway 1.1a

#### Community 1.1 to 1.2



eightpetal mountain-avens -  
arctic willow / northern  
singlespike sedge - fescue /  
curled snow lichen -  
whiteworm lichen



northern singlespike sedge -  
fescue / fireweed / Ceratodon  
moss - juniper Polytrichum  
moss

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank.

### Pathway 1.2a

#### Community 1.2 to 1.1



northern singlespike sedge -  
fescue / fireweed / Ceratodon  
moss - juniper Polytrichum  
moss

eightpetal mountain-avens -  
arctic willow / northern  
singlespike sedge - fescue /  
curled snow lichen -  
whiteworm lichen

Time without fire results in decreases to graminoid and forb cover and increases to shrub and lichen cover.

## Additional community tables

Table 7. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)	Diameter (In)	Basal Area (Square Ft/Acre)
<b>Tree</b>							
white spruce	PIGL	<i>Picea glauca</i>	Native	–	0–1	–	–

Table 8. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
<b>Grass/grass-like (Graminoids)</b>					
Altai fescue	FEAL	<i>Festuca altaica</i>	Native	0.3–1	0–20
alpine fescue	FEBR	<i>Festuca brachyphylla</i>	Native	0.1–0.3	0–20
northern singlespike sedge	CASC10	<i>Carex scirpoidea</i>	Native	0.3–1	0–15
Bigelow's sedge	CABI5	<i>Carex bigelowii</i>	Native	0.3–1	0–2
<b>Forb/Herb</b>					
arctic lupine	LUAR2	<i>Lupinus arcticus</i>	Native	0.3–1	0–10
blackish oxytrope	OXNI	<i>Oxytropis nigrescens</i>	Native	0.1–0.3	0–10
Drummond's anemone	ANDR	<i>Anemone drummondii</i>	Native	0.1–0.3	0–7
purple mountain saxifrage	SAOP	<i>Saxifraga oppositifolia</i>	Native	0.1–0.3	0–5
alpine bistort	POVI3	<i>Polygonum viviparum</i>	Native	0.1–0.3	0–5
moss campion	SIAC	<i>Silene acaulis</i>	Native	0.1–0.3	0–5
stitchwort	MINUA	<i>Minuartia</i>	Native	0.1–0.3	0–5
Langsdorf's lousewort	PELA3	<i>Pedicularis langsdorffii</i>	Native	0.3–1	0–3
northern asphodel	TOCO	<i>Tofieldia coccinea</i>	Native	0.1–0.3	0–3
American throw wax	BUAM2	<i>Bupleurum americanum</i>	Native	0.3–1	0–1
northern Indian paintbrush	CAHY6	<i>Castilleja hyperborea</i>	Native	0.1–0.3	0–0.1
Yukon bellflower	CAAU	<i>Campanula aurita</i>	Native	0.1–0.3	0–0.1
alpine meadow-rue	THAL	<i>Thalictrum alpinum</i>	Native	0.1–0.3	0–0.1
<b>Shrub/Subshrub</b>					
eightpetal mountain-avens	DROCO	<i>Dryas octopetala ssp. octopetala</i>	Native	0.1–0.3	7–70
arctic willow	SAAR27	<i>Salix arctica</i>	Native	0.1–0.3	0–35
netleaf willow	SARE2	<i>Salix reticulata</i>	Native	0.1–0.3	0–25
white arctic mountain heather	CATE11	<i>Cassiope tetragona</i>	Native	0.1–0.3	0–15
bog blueberry	VAUL	<i>Vaccinium uliginosum</i>	Native	0.1–0.3	0–10
Lapland rosebay	RHLA2	<i>Rhododendron lapponicum</i>	Native	0.1–0.3	0–10
<b>Nonvascular</b>					
	FLCU	<i>Flavocetraria cucullata</i>	Native	0.1–0.3	0.1–27
whiteworm lichen	THVE60	<i>Thamnolia vermicularis</i>	Native	0.1–0.3	0–10
	FLNI	<i>Flavocetraria nivalis</i>	Native	0.1–0.3	0–10
cetraria lichen	CEER6	<i>Cetraria ericetorum</i>	Native	0.1–0.3	0–10
arctic dactylina lichen	DAAR60	<i>Dactylina arctica</i>	Native	0.1–0.3	0–10
	VUTI	<i>Vulpicida tilesii</i>	Native	0.1–0.3	0–10
witch's hair lichen	ALOC60	<i>Alectoria ochroleuca</i>	Native	0.1–0.3	0–8
bryocaulon lichen	BRDI60	<i>Bryocaulon divergens</i>	Native	0.1–0.3	0–8
greengreen reindeer lichen	CLRA60	<i>Cladina rangiferina</i>	Native	0.1–0.3	0–5

## Animal community

n/a

## Hydrological functions

n/a

## Recreational uses

n/a

## Wood products

n/a

## Other products

n/a

## Other information

n/a

## Inventory data references

Tier 2 sampling plots used to develop the reference state. Plot numbers as recorded in NASIS with associated community phase.

### Community 1.1

10NP02201, 10NP02205, 10NP02206, 10NP03501, 10NP03503, 10NP03505, 10TC02303, 10TC02305, 10TC02701, 13NR00602, 13NR00901, 2015AK290551, 2016AK290458, 2016AK290632

### Community 1.2

13BA00501, 2016AK290572

## References

Chapin, F.S., L.A. Viereck, P.C. Adams, K.V. Cleve, C.L. Fastie, R.A. Ott, D. Mann, and J.F. Johnstone. 2006. Successional processes in the Alaskan boreal forest. Page 100 in *Alaska's changing boreal forest*. Oxford University Press.

Hinzman, L.D., L.A. Viereck, P.C. Adams, V.E. Romanovsky, and K. Yoshikawa. 2006. Climate and permafrost dynamics of the Alaskan boreal forest. *Alaska's changing boreal forest* 39–61.

Johnstone, J.F. 2008. A key for predicting postfire successional trajectories in black spruce stands of interior Alaska. US Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Schoeneberger, P.J. and D.A. Wysocki. 2012. Geomorphic Description System. Natural Resources Conservation Service, 4.2 edition. National Soil Survey Center, Lincoln, NE.

United States Department of Agriculture, . 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

## Other references

Alaska Interagency Coordination Center (AICC). 2022. <http://fire.ak.blm.gov/>

J. C. Springer and B.D. Parfitt. 2015. Dryas. Flora of North America, Volume 9, Rosaceae. [http://www.efloras.org/florataxon.aspx?flora\\_id=1&taxon\\_id=110971](http://www.efloras.org/florataxon.aspx?flora_id=1&taxon_id=110971) [accessed 01/11/2023].

LANDFIRE. 2009. Western North American Boreal Alpine Dwarf-Shrub Summit. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.

PRISM Climate Group. 2018. Alaska – average monthly and annual precipitation and minimum, maximum, and mean temperature for the period 1981-2010. Oregon State University, Corvallis, Oregon. <https://prism.oregonstate.edu/projects/alaska.php>. (Accessed 4 September 2019).

United States Department of Agriculture-Natural Resources Conservation Service. 2016. U.S. General Soil Map (STATSGO2). Web Soil Survey. Available online at <http://websoilsurvey.nrcs.usda.gov>. Accessed (Accessed 3 March 2021).

## Contributors

Blaine Spellman  
Jamin Johanson  
Stephanie Shoemaker  
Phillip Barber

## Approval

Kirt Walstad, 2/13/2024

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/23/2024
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**
- 

2. **Presence of water flow patterns:**
-



3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---