

## **Ecological site R231XY115AK**

### **Alpine sedge silty frozen slopes**

Last updated: 2/13/2024  
Accessed: 05/03/2024

---

#### **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

#### **MLRA notes**

Major Land Resource Area (MLRA): 231X—Interior Alaska Highlands

The Interior Alaska Uplands (MLRA 231X) is in the Interior Region of Alaska and includes the extensive hills, mountains, and valleys between the Tanana River to the south and the Brooks Range to the north. These hills and mountains surround the Yukon Flats Lowlands (MLRA 232X). MLRA 231X makes up about 69,175 square miles. The hills and mountains of the area tend to be moderately steep to steep resulting in high-relief slopes. The mountains are generally rounded at lower elevations and sharp-ridged at higher elevations. Elevation ranges from about 400 feet in the west, along the boundary with the Interior Alaska Lowlands (MLRA 229X), to 6,583 feet at the summit of Mt. Harper, in the southeast. Major tributaries include large sections of the Yukon, Koyukuk, Kanuti, Charley, Coleen, and Chatanika Rivers. This area is traversed by several major roads, including the Taylor Highway in the east and the Steese, Elliott, and Dalton Highways north of Fairbanks. The area is mostly undeveloped wild land that is sparsely populated. The largest community along the road system is Fairbanks with smaller communities like Alatna, Allakaket, Chicken, Eagle, Eagle Village, Hughes, and Rampart occurring along the previously mentioned rivers and highways.

The vast majority of this MLRA was unglaciated during the Pleistocene epoch with the exceptions being the highest mountains and where glaciers extended into the area from the Brooks Range. For the most part, glacial moraines and drift are limited to the upper elevations of the highest mountains. Most of the landscape is mantled with bedrock colluvium originating from the underlying bedrock. Valley bottoms are filled with Holocene fluvial deposits and colluvium from the adjacent mountain slopes. Silty loess, which originated from unvegetated flood plains in and adjacent to this area, covers much of the surface. On hill and mountain slopes proximal to major river valleys (e.g., Tanana and Yukon Rivers), the loess is many feet thick. As elevation and distance from major river valleys increases, loess thickness decreases significantly. Bedrock is commonly exposed on the highest ridges.

This area is in the zone of discontinuous permafrost. Permafrost commonly is close to the surface in areas of the finer textured sediments throughout the MLRA. Isolated masses of ground ice occur in thick deposits of loess on terraces and the lower side slopes of hills. Solifluction lobes, frost boils, and circles and stripes are periglacial features common on mountain slopes in this area. Pingos, thermokarst pits and mounds, ice-wedge polygons, and earth hummocks are periglacial features common on terraces, lower slopes of hills and mountains, and in upland valleys in the area.

The dominant soil orders in this area are Gelisols, Inceptisols, Spodosols, and Entisols. The soils in the area have a subgelic or cryic soil temperature regime, an aquic or udic soil moisture regime, and mixed mineralogy. Gelisols are common on north facing slopes, south facing footslopes, valley bottoms, and stream terraces. Gelisols are typically shallow or moderately deep to permafrost (10 to 40 inches) and are poorly or very poorly drained. Wildfires can disturb the insulating organic material at the surface, lowering the permafrost layer, eliminating perched water tables from Gelisols, and thus changing the soil classification. Inceptisols and Spodosols commonly form on south facing hill and mountain slopes. Entisols are common on flood plains and high elevation mountain slopes. Miscellaneous (non-soil) areas make up about 2 percent of this MLRA. The most common miscellaneous areas are rock outcrop and rubble land. In many valleys placer mine tailings are common.

Short, warm summers and long, cold winters characterize the subarctic continental climate of the area. The mean annual temperature of the area ranges from 22 to 27 degrees F. The mean annual temperature of the southern half of the area is approximately 3 degrees warmer compared to the northern half (PRISM 2018). The warmest months span June through August with mean monthly temperatures ranging from 50 to 56 degrees F. The coldest months span November through February with mean monthly temperatures ranging from -5 to 3 degrees F. When compared to the high-elevation alpine and subalpine life zones, the lower elevation boreal life zone tends to be 2-3 degrees F colder during the coldest months and 1-2 degrees F warmer during the warmest months (PRISM 2018). The freeze-free period at the lower elevations averages about 60 to 100 days, and the temperature usually remains above freezing from June through mid-September.

Precipitation is limited across this area, with the average annual precipitation ranging from 12 to 19 inches. The southern half of the areas receives approximately 2.5 inches more annual precipitation than the northern half (PRISM 2018). The lower elevation boreal life zone receives approximately 2.5 inches less annual precipitation than the high-elevation alpine and subalpine life zones (PRISM 2018). Approximately 3/5th of the annual precipitation occurs during the months of June through September with thunderstorms being common. The average annual snowfall ranges from about 45 to 100 inches. The ground is consistently covered with snow from November through March.

Most of this area is forested below an elevation of about 2500 feet. Dominant tree species on slopes are white spruce and black spruce. Black spruce stands are most common on north-facing slopes, stream terraces, and other sites with poor drainage and permafrost. White spruce stands are most common on warm slopes with dry soils. At lower elevations, lightning-caused wildfires are common, often burning many thousands of acres during a single fire. Following wildfires, forbs, grasses, willow, ericaceous shrubs, paper birch, and quaking aspen communities are common until they are eventually replaced by stands of spruce. Tall willow and alder scrub is extensive on low flood plains. White spruce and balsam poplar are common on high flood plains.

With increasing elevation, the forests and woodlands give way to subalpine communities dominated by krummholz spruce, shrub birch, willow, and ericaceous shrubs. At even higher elevations, alpine communities prevail which are characterized by diverse forbs, dwarf ericaceous shrubs, and eightpetal mountain-avens. Many of these high elevation communities have a considerable amount of lichen cover and bare ground.

## **LRU notes**

This area supports three life zones defined by the physiological limits of plant communities along an elevational gradient: boreal, subalpine, and alpine. The boreal life zone is the elevational band where forest communities dominate. Not all areas in the boreal life zone are forest communities, however, particularly in places with too wet or dry soil to support tree growth (e.g., bogs or river bluffs). Above the boreal band of elevation, subalpine and alpine vegetation dominate. The subalpine zone is typically a narrow transitional band between the boreal and the alpine life zones, and is characterized by sparse, stunted trees. In the subalpine, certain types of birch and willow shrub species grow at  $\geq 1$  m in height (commonly *Betula glandulosa* and *Salix pulchra*). In the alpine, trees no longer occur, and all shrubs are dwarf or lay prostrate on the ground. In this area, the boreal life zone occurs below 2500 feet elevation on average. The transition between boreal and alpine vegetation can occur within a range of elevations, and is highly dependent on slope, aspect, and shading from adjacent mountains.

Within each life zone, there are plant assemblages that are typically associated with cold slopes and warm slopes. Cold slopes and warm slopes are created by the combination of the steepness of the slope, the aspect, and shading from surrounding ridges and mountains. Warm slope positions typically occur on southeast to west facing slopes that are moderate to very steep ( $>10\%$  slope) and are not shaded by the surrounding landscape. Cold slopes typically occur on northwest to east facing slopes, occur in shaded slope positions, or occur in low-lying areas that are cold air sinks. Examples of shaded positions include head slopes, low relief backslopes of hills, and the base of hills and mountains shaded by adjacent mountain peaks. Warm boreal slope soils have a cryic soil temperature regime and lack permafrost. In this area, white spruce forests are an indicator of warm boreal slopes. Cold boreal slope soils typically have a gelic soil temperature regime and commonly have permafrost. In this area, black spruce forests and woodlands are an indicator of cold boreal slopes. The boreal life zone can occur at higher elevations on warm slopes, and lower elevations on cold slopes.

## Classification relationships

Landfire BPS – 7416330 – Western North American Boreal Alpine Mesic Herbaceous Meadow

## Ecological site concept

This site occurs on alpine slopes with wet, silty, frozen soils. This site is generally associated with the summits and shoulders of rounded mountains at high elevation. Turf hummocks are common periglacial features associated with this site. Soils do not flood but commonly pond. These poorly to very poorly drained soils have a high-water table that remain wet at very shallow depth throughout the growing season. Permafrost typically occurs in the soil profile at moderate depth. These soils formed in silty loess that occasionally have limited rock fragments.

The alpine life zone has a harsh climate that limits growth of vegetation and prevents the establishment of many species common at lower elevations. In this area, alpine vegetation is characterized as dwarf and prostrate shrubs intermixed with low-lying herbaceous plants. These unique plant communities are the result of high winds, a short growing season, deep and persistent snow beds, and cold soils. These climatic factors prevent the establishment and growth of many dominant boreal species like white spruce and black spruce.

Two plant communities occur within the reference state and the vegetation in each community differs in large part due to fire. When the reference state vegetation burns, the post-fire plant community is dominantly ericaceous shrubs and sedges. With time and lack of another fire event, the post-fire vegetation goes through succession. For this site, the reference plant community is the most stable with the longest time since the vegetation was burned. This community is typically characterized as mixed shrub-sedge tussock tundra (Viereck et al. 1992) with bog blueberry, marsh Labrador tea, Bigelow's sedge, tussock cottongrass, and curled snow lichen the dominant vegetation. The vegetative strata that characterize this community are dwarf shrubs (less than 8 inches), medium graminoids (between 4 and 24 inches), and foliose and fruticose lichen.

## Associated sites

R231XY101AK	<b>Alpine dwarf scrub gravelly slopes</b> Occurs in the alpine but on dry and gravelly soils without permafrost.
R231XY113AK	<b>Alpine Dwarf Scrub Gravelly Moist Slopes</b> Occurs in the alpine but on moist and gravelly soils without permafrost.
R231XY134AK	<b>Alpine Dwarf Scrub Gravelly Frozen Slopes</b> Occurs in the alpine but on wet, gravelly, and frozen soils.
R231XY152AK	<b>High-elevation scrub gravelly drainageways</b> Occurs downslope in high elevation drainageways.

## Similar sites

R231XY103AK	<b>Alpine Dwarf Scrub Gravelly Frozen Alkaline Slopes</b> Both sites occur on alpine slopes with wet and frozen soils. Site 103 has alkaline soils resulting in different kinds and amounts of vegetation.
R231XY134AK	<b>Alpine Dwarf Scrub Gravelly Frozen Slopes</b> Site 134 has much gravelly soils. Plant communities associated with 134 tend to have more dwarf shrub cover and less sedge cover.
R231XY128AK	<b>Boreal Tussock Peat Frozen Slopes</b> Occurs at lower elevations in the boreal life zone resulting in different kinds and amounts of vegetation. Site 128 has comparatively less lichen cover which may result from differences in fire frequency.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Carex bigelowii</i> (2) <i>Flavocetraria cucullata</i>

Physiographic features

This site occurs at high elevation on slopes that commonly have extensive turf hummocks in the alpine life zone. This site is most associated with the summits and shoulders of rounded mountains. On occasion, the site occurs on nearly level slopes of rugged mountains, plateaus, and plains. On rare occasions the site occurs on rounded mountains with high-centered polygons. Turf hummocks are common periglacial features on all associated landforms and consist of vegetation and organic matter with or without a core of mineral soil and are typically 10 to 50 cm in height and 20 to 90 cm in diameter (Schoeneberger and Wysocki 2017). Elevation typically ranges between 2500 and 3100 feet but can go as low as 2050 feet and as high as 4150 feet on some summits of mountains. Summits are nearly level, shoulders are gently sloping, and these slopes occur on all aspects. The site does not flood. In general, ponding occurs frequently to occasionally for brief to long durations of time. A water table occurs at very shallow depth for much of the growing season. This site generates very limited runoff to adjacent, downslope sites.



Figure 1. Turf hummocks commonly associated with this site.



Figure 2. High-centered polygons with turf hummocks associated with this site.

Table 2. Representative physiographic features

Hillslope profile	(1) Summit (2) Shoulder
Landforms	(1) Mountains > Mountain slope > Turf hummock (2) Mountains > Plain > Turf hummock (3) Mountains > Plateau > Turf hummock (4) Mountains > Mountain slope > Polygon
Runoff class	Negligible to very low
Flooding frequency	None

Ponding duration	Brief (2 to 7 days) to long (7 to 30 days)
Ponding frequency	Occasional to frequent
Elevation	2,500–3,100 ft
Slope	0–4%
Ponding depth	2–12 in
Water table depth	0 in
Aspect	W, NW, N, NE, E, SE, S, SW

**Table 3. Representative physiographic features (actual ranges)**

Runoff class	Not specified
Flooding frequency	Not specified
Ponding duration	Not specified
Ponding frequency	Not specified
Elevation	2,050–4,150 ft
Slope	0–15%
Ponding depth	0–12 in
Water table depth	0–10 in

## Climatic features

When compared to the boreal life zone, this high-elevation site has a harsh climate. In this MLRA, snow first blankets and persists the longest in the alpine and subalpine life zones. From spring through fall (April through September) , it is consistently 1 to 2 degrees F colder in the alpine and subalpine. These small differences in temperature are exacerbated due to constant and strong winds. Winds are much more intense in these high elevation areas because of limited trees providing windbreaks. When compared to the boreal life zone, this site has a much shorter growing season and the growing season is significantly colder for associated vegetation.

Short, warm summers and long, cold winters characterize the subarctic continental climate associated with this high-elevation site. The mean annual temperature of the site ranges from 23 to 27 degrees F. The warmest months span June through August with mean normal maximum monthly temperatures ranging from 57 to 63 degrees F. The coldest months span November through February with mean normal minimum temperatures ranging from -9 to -1 degrees F. The freeze-free period for the site ranges from 80 to 120 days, and the temperature usually remains above freezing from late May through mid-September.

The area receives minimal annual precipitation with the summer months being the wettest. Average annual precipitation in the alpine across the area typically ranges between 14 to 21 inches. Approximately 3/5th of the annual precipitation occurs during the months of June through September with thunderstorms common. The average annual snowfall ranges from about 45 to 100 inches. The ground is consistently covered with snow from mid-October through March.

**Table 4. Representative climatic features**

Frost-free period (characteristic range)	16-78 days
Freeze-free period (characteristic range)	76-114 days
Precipitation total (characteristic range)	14-21 in
Frost-free period (actual range)	4-87 days
Freeze-free period (actual range)	48-120 days
Precipitation total (actual range)	10-25 in
Frost-free period (average)	53 days

Freeze-free period (average)	90 days
Precipitation total (average)	17 in




Figure 3. Monthly precipitation range




Figure 4. Monthly minimum temperature range




Figure 5. Monthly maximum temperature range






Figure 6. Monthly average minimum and maximum temperature



**Figure 7. Annual precipitation pattern**



**Figure 8. Annual average temperature pattern**

## Climate stations used

- (1) EAGLE AP [USW00026422], Tok, AK
- (2) CHICKEN [USC00501684], Tok, AK
- (3) MILE 42 STEESE [USC00505880], Fairbanks, AK
- (4) BETTLES AP [USW00026533], Bettles Field, AK
- (5) CIRCLE HOT SPRINGS [USC00501987], Central, AK
- (6) FT KNOX MINE [USC00503160], Fairbanks, AK
- (7) GILMORE CREEK [USC00503275], Fairbanks, AK
- (8) FOX 2SE [USC00503181], Fairbanks, AK
- (9) ESTER DOME [USC00502868], Fairbanks, AK
- (10) ESTER 5NE [USC00502871], Fairbanks, AK
- (11) COLLEGE 5 NW [USC00502112], Fairbanks, AK
- (12) COLLEGE OBSY [USC00502107], Fairbanks, AK
- (13) KEYSTONE RIDGE [USC00504621], Fairbanks, AK

## Influencing water features

This site is classified as a slope wetland under the Hydrogeomorphic (HGM) classification system (Smith et al. 1995; USDA-NRCS 2008). Precipitation and ground water are the main sources of water (Smith et al. 1995).

Depth to the water table may decrease following summer storm events or spring snowmelt and increase during extended dry periods.

## Wetland description

n/a

## Soil features

Soils formed in windblown silt over gravelly colluvium and have permafrost. Surface rock fragments are not present. These are mineral soils capped with 9 to 24 inches of saturated organic material. The mineral soil below the organic



material is a silt loam formed from wind-blown loess, which typically lacks rock fragments and has high water holding capacity. This silty layer is generally very thick and often cryoturbated. On occasion, cryoturbated material has large rock fragments in the soil profile with rock fragments commonly ranging between 0 and 15 percent of the soil profile by volume. While soils are very deep, permafrost typically occurs at moderate depths (20 to 32 inches). The pH of the soil profile ranges from extremely acidic to moderately acidic. The soils are wet for long portions of the growing season and are considered very poorly to poorly drained.



**Figure 9. A typical soil profile associated with this site.**

**Table 5. Representative soil features**

Parent material	(1) Loess (2) Eolian deposits (3) Colluvium
Surface texture	(1) Peat
Family particle size	(1) Coarse-silty (2) Coarse-loamy
Drainage class	Very poorly drained to poorly drained
Permeability class	Moderately rapid
Depth to restrictive layer	20–32 in
Soil depth	60 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	4.7–9.4 in
Calcium carbonate equivalent (10-40in)	0%
Clay content (0-20in)	5–10%
Electrical conductivity (10-40in)	0–3 mmhos/cm
Sodium adsorption ratio (10-40in)	0
Soil reaction (1:1 water) (10-40in)	3.3–5.6
Subsurface fragment volume <=3" (0-60in)	0–10%



Subsurface fragment volume >3" (0-60in)	0–3%
--------------------------------------------	------

**Table 6. Representative soil features (actual values)**

Drainage class	Not specified
Permeability class	Not specified
Depth to restrictive layer	12–35 in
Soil depth	Not specified
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-40in)	3.7–15.8 in
Calcium carbonate equivalent (10-40in)	Not specified
Clay content (0-20in)	Not specified
Electrical conductivity (10-40in)	Not specified
Sodium adsorption ratio (10-40in)	Not specified
Soil reaction (1:1 water) (10-40in)	3.3–6.5
Subsurface fragment volume <=3" (0-60in)	0–20%
Subsurface fragment volume >3" (0-60in)	0–10%

## Ecological dynamics

### Climate

Located in the alpine life zone, this site is exposed to a variety of harsh environmental conditions. In this area, snowfall first appears and persists the longest in the alpine. As a result, snowpack tends to be deeper and persist for longer durations of time compared to lower-elevation sites and alpine vegetation has a comparatively shorter growing season. When this site is snow-free, cold soil temperatures and high winds also inhibit plant growth and vigor. This harsh climate maintains the dwarfed vegetation within this site and prevents the establishment and/or growth of dominant boreal species like white spruce and black spruce.

### Fire

Within this area, fire is considered a natural and common event that typically is unmanaged. Fire suppression is limited, and generally occurs adjacent to Fairbanks and the various villages spread throughout the area or on allotments with known structures, all of which have a relatively limited acre footprint. Most fires are caused by lightning strikes. From 2000 to 2020, 596 known fire events occurred in the Interior Alaska Uplands area and the burn perimeter of the fires totaled about 13.8 million acres (AICC 2022). Fire-related disturbances are highly patchy and can leave undisturbed areas within the burn perimeter. During this time frame, 80 percent of the fire events were smaller than 20,000 acres but 18 fire events were greater than 200,000 acres in size (AICC 2022). Over this period of 20 years, these burn perimeters cover approximately 30 percent of the Interior Alaska Uplands area.

The fire regime within Interior Alaska follows two general scenarios—low-severity burns and high-severity burns. It should be noted, however, that the fire regime in Interior Alaska can be much more complex (Johnstone et al. 2008). Burn severity refers to the proportion of the vegetative canopy and organic material consumed in a fire event (Chapin et al. 2006). Fires in cool and moist habitat tend to result in low-severity burns, while fires in warm and dry

habitat tend to result in high-severity burns. Because the soils have a thick organic cap and are poorly to very poorly drained, the typical fire scenario for this ecological site is considered to result in a low severity burn.


While low-severity fires have a range of impacts to vegetation and soils for this site, permafrost generally remains in the soil profile. While a low-severity fire can consume the bulk of above ground vegetation, minimal proportions of the organic mat are removed. Organic matter continues to insulate these cold soils. Field data support that each plant community has permafrost and data from similar sites suggest that the associated low-severity fire event has a negligible impact on the depth of permafrost. If permafrost remains at similar depths after a fire event, then soil drainage is unlikely to improve post-fire. For this site, additional plots and environmental co-variate data will help clarify the variability in fire severity (e.g., timing of fire, soil organic matter moisture content, and pre-fire vegetation) and its effects to soil organic thickness, depth to permafrost, and drainage.

When minimal proportions of the organic mat are consumed, many species regenerate asexually from belowground root systems or rhizomes. Species known to regenerate after low-severity fire events include various graminoids (e.g. *Carex* spp. and *Eriophorum* spp.), forbs (e.g. *Equisetum* sp.), and shrubs (e.g. *Ledum groenlandicum*, *Vaccinium uliginosum*, *Salix* sp.) (Johnstone et al. 2010).


Because the dominant vegetation in community 1.1 is a mixture of quick growing species (sedges and ericaceous shrubs) and slow growing species (lichens), an unknown timeframe is required for postfire recovery back to the reference plant community. After a low-severity fire event, field data suggest that sedge and ericaceous shrubs regenerate quickly and are dominant (community 1.2). Lichen continues to colonize and grow until it also becomes dominant in the plant community, which marks the transition back to the reference plant community (community 1.1). Lichen are a slow growing species and post-fire recovery may take greater than 10 to 30 years.

State and transition model

Ecosystem states



State 1 submodel, plant communities



1.1a - A low-severity fire sweeps through and incinerates much of the above ground vegetation.

1.2a - Time without fire.

State 1  
Reference State



**Figure 10. A sedge dominant community associated with this alpine site.**

The reference plant community is mixed shrub-sedge tussock tundra (Viereck et al. 1992). There are two plant communities in the reference state related to fire.

### **Dominant plant species**

- bog blueberry (*Vaccinium uliginosum*), shrub
- marsh Labrador tea (*Ledum palustre ssp. decumbens*), shrub
- Bigelow's sedge (*Carex bigelowii*), grass
- tussock cottongrass (*Eriophorum vaginatum*), grass
- (*Flavocetraria cucullata*), other herbaceous
- sphagnum (*Sphagnum*), other herbaceous

### **Community 1.1**

**bog blueberry - marsh Labrador tea / Bigelow's sedge - tussock cottongrass / curled snow lichen - Sphagnum**



**Figure 11. A typical plant community associated with community 1.1.**

The reference plant community is characterized as mixed shrub-sedge tussock tundra (Viereck et al. 1992) with Bigelow's sedge, tussock cottongrass, and curled snow lichen the dominant vegetation. Stunted white spruce occasionally occur but have limited cover. Other common species include marsh Labrador tea, bog blueberry, scrub birch, lingonberry, cloudberry, various reindeer lichen, and Sphagnum. The vegetative strata that characterize this community are dwarf shrubs (less than 8 inches), medium graminoids (between 4 and 24 inches), and foliose and fruticose lichen. The soil surface is primarily covered with moss and lichen.

### **Dominant plant species**

- marsh Labrador tea (*Ledum palustre ssp. decumbens*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- resin birch (*Betula glandulosa*), shrub

- lingonberry (*Vaccinium vitis-idaea*), shrub
- Bigelow's sedge (*Carex bigelowii*), grass
- tussock cottongrass (*Eriophorum vaginatum*), grass
- (*Flavocetraria cucullata*), other herbaceous
- sphagnum (*Sphagnum*), other herbaceous
- greygreen reindeer lichen (*Cladina rangiferina*), other herbaceous
- cloudberry (*Rubus chamaemorus*), other herbaceous
- reindeer lichen (*Cladina stygia*), other herbaceous

## Community 1.2

### Bigelow's sedge - tussock cottongrass / Sphagnum



Figure 12. A typical plant community associated with community 1.2.


Community 1.2 is in the early stage of fire-induced secondary succession for this ecological site. Community 1.2 is characterized as mixed shrub-sedge tussock tundra (Viereck et al. 1992) with Bigelow's sedge, tussock cottongrass, and Sphagnum the dominant vegetation. Other common species include marsh Labrador tea, lingonberry, scrub birch, bog rosemary, alpine bearberry, wideleaf polargrass, and cloudberry. The vegetative strata that characterize this community are low shrubs (between 8 and 36 inches), medium graminoids (between 4 and 24 inches), and mosses. The soil surface is primarily covered with herbaceous litter and moss.

### Dominant plant species

- marsh Labrador tea (*Ledum palustre* ssp. *decumbens*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- resin birch (*Betula glandulosa*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- alpine bearberry (*Arctostaphylos alpina*), shrub
- bog rosemary (*Andromeda polifolia*), shrub
- Bigelow's sedge (*Carex bigelowii*), grass
- tussock cottongrass (*Eriophorum vaginatum*), grass
- wideleaf polargrass (*Arctagrostis latifolia*), grass
- sphagnum (*Sphagnum*), other herbaceous
- cloudberry (*Rubus chamaemorus*), other herbaceous

## Pathway 1.1a

### Community 1.1 to 1.2




bog blueberry - marsh  
Labrador tea / Bigelow's sedge  
- tussock cottongrass / curled  
snow lichen - Sphagnum

Bigelow's sedge - tussock  
cottongrass / Sphagnum

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated cold and wet soils, this site commonly experiences low-severity fires. Minimal proportions of the organic mat are typically removed. The pre-fire vegetation generally reestablishes quickly from below ground root systems and rhizomes.

Pathway 1.2a  
Community 1.2 to 1.1



Bigelow's sedge - tussock  
cottongrass / Sphagnum

bog blueberry - marsh  
Labrador tea / Bigelow's sedge  
- tussock cottongrass / curled  
snow lichen - Sphagnum

Time without fire results in decreases to graminoid and moss cover and increases to lichen cover.

Additional community tables

Table 7. Community 1.1 forest overstory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)	Diameter (In)	Basal Area (Square Ft/Acre)
Tree							
white spruce	PIGL	<i>Picea glauca</i>	Native	1–3	0–0.1	–	–

Table 8. Community 1.1 forest understory composition

Common Name	Symbol	Scientific Name	Nativity	Height (Ft)	Canopy Cover (%)
<b>Grass/grass-like (Graminoids)</b>					
Bigelow's sedge	CABI5	<i>Carex bigelowii</i>	Native	0.3–1	0–40
tussock cottongrass	ERVA4	<i>Eriophorum vaginatum</i>	Native	0.3–1	0–40
<b>Forb/Herb</b>					
cloudberry	RUCH	<i>Rubus chamaemorus</i>	Native	0.1–0.3	0–10
<b>Shrub/Subshrub</b>					
lingonberry	VAVI	<i>Vaccinium vitis-idaea</i>	Native	0.1–0.3	1–30
marsh Labrador tea	LEPAD	<i>Ledum palustre ssp. decumbens</i>	Native	0.8–2	2–20
bog blueberry	VAUL	<i>Vaccinium uliginosum</i>	Native	0.8–2	0.1–20
resin birch	B EGL	<i>Betula glandulosa</i>	Native	0.8–2	0–15
alpine bearberry	ARAL2	<i>Arctostaphylos alpina</i>	Native	0.1–0.3	0–10
black crowberry	EMNI	<i>Empetrum nigrum</i>	Native	0.1–0.3	0–10
bog rosemary	ANPO	<i>Andromeda polifolia</i>	Native	0.1–0.3	0–4
<b>Nonvascular</b>					
	FLCU	<i>Flavocetraria cucullata</i>	Native	0.1–0.3	25–75
sphagnum	SPHAG2	<i>Sphagnum</i>	Native	0.1–0.3	0–35
greygreen reindeer lichen	CLRA60	<i>Cladina rangiferina</i>	Native	0.1–0.3	0–30
reindeer lichen	CLST5	<i>Cladina stygia</i>	Native	0.1–0.3	0–10
whiteworm lichen	THSU60	<i>Thamnolia subuliformis</i>	Native	0.1–0.3	0–8
polytrichum moss	POLYT5	<i>Polytrichum</i>	Native	0.1–0.3	0–5
cup lichen	CLADO3	<i>Cladonia</i>	Native	0.1–0.3	0–5
arctic dactylina lichen	DAAR60	<i>Dactylina arctica</i>	Native	0.1–0.3	0–4
cetraria lichen	CELA60	<i>Cetraria laevigata</i>	Native	0.1–0.3	0–3
aulacomnium moss	AUPA70	<i>Aulacomnium palustre</i>	Native	0.1–0.3	0–2

Animal community

n/a

Hydrological functions

n/a

Recreational uses

n/a

Wood products

n/a

Other products

n/a

Other information

n/a



## Inventory data references

Tier 2 sampling plots used to develop the reference state. Plot numbers as recorded in NASIS with associated community phase.

### Community 1.1

09TC00104, 12CP00501, 12NR03802, 12NR03901, 12NR03902, 12SN02001, 12SN02201, 12SN03201, S2014AK290005

### Community 1.2

08TC00703, 08TC00705

## References

Chapin, F.S., L.A. Viereck, P.C. Adams, K.V. Cleve, C.L. Fastie, R.A. Ott, D. Mann, and J.F. Johnstone. 2006. Successional processes in the Alaskan boreal forest. Page 100 in *Alaska's changing boreal forest*. Oxford University Press.

Hinzman, L.D., L.A. Viereck, P.C. Adams, V.E. Romanovsky, and K. Yoshikawa. 2006. Climate and permafrost dynamics of the Alaskan boreal forest. *Alaska's changing boreal forest* 39–61.

Johnstone, J.F., T.N. Hollingsworth, F.S. CHAPIN III, and M.C. Mack. 2010. Changes in fire regime break the legacy lock on successional trajectories in Alaskan boreal forest. *Global change biology* 16:1281–1295.

Johnstone, J.F. 2008. A key for predicting postfire successional trajectories in black spruce stands of interior Alaska. US Department of Agriculture, Forest Service, Pacific Northwest Research Station.

Schoeneberger, P.J. and D.A. Wysocki. 2012. Geomorphic Description System. Natural Resources Conservation Service, 4.2 edition. National Soil Survey Center, Lincoln, NE.

United States Department of Agriculture, . 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

## Other references

Alaska Interagency Coordination Center (AICC). 2022. <http://fire.ak.blm.gov/>

J. C. Springer and B.D. Parfitt. 2015. *Dryas*. *Flora of North America*, Volume 9, Rosaceae. [http://www.efloras.org/florataxon.aspx?flora\\_id=1&taxon\\_id=110971](http://www.efloras.org/florataxon.aspx?flora_id=1&taxon_id=110971) [accessed 01/11/2023].

LANDFIRE. 2009. Western North American Boreal Alpine Mesic Herbaceous Meadow. In: *LANDFIRE National Vegetation Dynamics Models*. USDA Forest Service and US Department of Interior. Washington, DC.

PRISM Climate Group. 2018. Alaska – average monthly and annual precipitation and minimum, maximum, and mean temperature for the period 1981-2010. Oregon State University, Corvallis, Oregon.



<https://prism.oregonstate.edu/projects/alaska.php>. (Accessed 4 September 2019).

United States Department of Agriculture-Natural Resources Conservation Service. 2016. U.S. General Soil Map (STATSGO2). Web Soil Survey. Available online at <http://websoilsurvey.nrcs.usda.gov>. Accessed (Accessed 3 March 2021).

## Contributors

Blaine Spellman  
Jamin Johanson  
Stephanie Shoemaker  
Phillip Barber

## Approval

Kirt Walstad, 2/13/2024

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/03/2024
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**
- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state**

for the ecological site:

---

17. **Perennial plant reproductive capability:**

---