

Ecological site F233XY171AK

Boreal Woodland Loamy Frozen Terraces

Last updated: 6/10/2025

Accessed: 03/21/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 233X–Upper Kobuk and Koyukuk Hills and Valleys

The Upper Kobuk and Koyukuk Hills and Valleys MLRA (herein called area) occurs in Interior Alaska. This area makes up 8,405 square miles. The largest tributaries are the Kobuk and the Koyukuk Rivers. Major tributaries of the Kobuk are the Reed, Beaver, Mauneluk, and Pau Rivers. Major tributaries of the Koyukuk River are the Alatna, John, and Kanuti Rivers. This area is primarily undeveloped wildland and sparsely populated. The communities within or near this area are Bettles, Kobuk, and Shungnak.

The terrain of this area consists of broad, nearly level river valleys and basins and rolling uplands separated by isolated hills and low rounded mountains. In the river valleys, nearly level flood plains and stream terraces gradually transition to gently sloping to moderately steep slopes leading to the hills and mountains. Basins are on the Pau River Flats between the eastern Zane and Lockwood Hills, on the Kanuti Flats between the Kanuti and Koyukuk Rivers, and along the middle reaches of the Hogatza River. Basins and stream terraces are dotted with hundreds of lakes and interconnecting wetlands. Elevation ranges from about 150 feet in the western part of the area, at the confluence of the Kobuk and Mauneluk Rivers, to 4,765 feet at the summit of Fritts Mountain, in the Angaycuham Mountains.

Geology and Soils

The northern part of the area was covered repeatedly by Pleistocene glaciers originating in the Brooks Range to the north. Slightly modified to highly modified moraines and drift cover many of the rolling uplands. Glacial ice flowed over most of the hills and low mountains, removing existing deposits and leaving a thin layer of glacial deposits. Today,

the lower mountain slopes, hills, and valley bottoms are covered with a variety of material, including glacial drift, colluvium, slope alluvium, fluvial deposits, and silty loess. In the southern part of the area, basins and valleys are filled with Quaternary glaciofluvial and fluvial deposits. Hills and upland slopes are covered with bedrock colluvium and slope alluvium, which are mantled with loess in places. The bedrock geology underlying much of the area consists dominantly of Permian through Lower Cretaceous stratified sedimentary and volcanic rocks.

This area is in the zone of discontinuous permafrost. Permafrost is close to the surface in lands with finer textured sediments throughout the area. Isolated masses of ground ice occur on terraces and the lower side slopes of hills. Permafrost does not occur on flood plains, on steep south-facing slopes, or other lands with very gravelly soils. Periglacial features, such as thermokarst pits, peat plateaus, and earth hummocks, are on the lower hill and mountain slopes and in upland valleys.

The dominant soil orders in this area are Gelisols, Inceptisols, and Entisols. The Gelisols are shallow or moderately deep to permafrost, occur on finer textured sediments, and are poorly drained or very poorly drained. Common Gelisol suborders are Histels, Orthels, and Turbels. The Histels have thick accumulations of surface organic material and occur in depressions, lake margins, and peat plateau. The Orthels and Turbels have comparably thinner surface organic material and occur on stream terraces and hill and upland slopes. The Inceptisols and Entisols are typically associated with gravelly soils that do not have permafrost within their profile, are deep, and are somewhat poorly drained to well drained. The common Inceptisol suborders are Cryepts and Gelepts both of which occur on upland and mountain slopes. Cryepts occur under forested soils at lower elevations and Gelepts on alpine tundra at higher elevations. Common Entisol suborders are Cryofluvents and Cryorthents both of which occur on alluvium on flood plains. Miscellaneous (non-soil) areas make up about 8 percent of this MLRA. The most common are rock outcrop, rubble land, and water.

Wildfires disturb the insulating organic material at the soil surface and can change the presence and/or depth of permafrost in the soil profile. These fire related changes to permafrost can also change the depth and presence of perched water tables. Gelisols that burn in this area can change soil taxonomic classification. For instance, depending on fire-severity, Histels may change to Orthels and Orthels may change to Inceptisols. Depending on the frequency and intensity of fires, landform position, and soil texture, the soils may or may not revert back to their original taxonomic classification.

Climate

Short, warm summers and long, cold winters characterize the continental subarctic climate of the area. The average annual precipitation ranges from 15 to 19 inches on valley bottoms and basins and from 19 to 26 inches at the higher elevations in the hills and mountains (PRISM 2018). Most of the precipitation falls as rain between May and September. The average annual snowfall ranges from 65 to 80 inches. The average

annual temperature is 22 to 24 degrees Fahrenheit (PRISM 2018). The temperature normally remains above freezing from mid-June through August in river valleys and basins with a freeze-free period ranging from 109 to 125 days. The freeze-free period is significantly shorter on higher elevation mountain slopes.

Vegetation

Most of this area is forested below an elevation of 1600 feet. Dominant tree species on slopes are white spruce and black spruce. Black spruce stands dominate on north-facing slopes, stream terraces, and other sites with poor drainage and permafrost. White spruce stands dominate on steep, south-facing slopes with dry soils. At lower elevations, lightning-caused wildfires are common, often burning many thousands of acres during a single fire event. Following wildfires, forbs, grasses, willow, ericaceous shrubs, paper birch, and quacking aspen communities are common until they are eventually replaced by stands of spruce. Tall willow and alder scrub is extensive on low flood plains. White spruce and balsam poplar are common on high flood plains.

With increasing elevation, the forests and woodlands give way to subalpine communities dominated by krummholz spruce, shrub birch, willow, and ericaceous shrubs. At even higher elevations, alpine communities prevail which are characterized by diverse forbs, dwarf ericaceous shrubs, and eightpetal mountain-avens. Many of these high elevation communities have a considerable amount of lichen cover and bare ground.

LRU notes

In this area, we refer to three life zones that are defined by the physiological limits of plant communities along an elevational gradient: boreal, subalpine, and alpine. The boreal life zone is the elevational band where forest communities dominate. Not all areas in the boreal life zone are forest communities, however, particularly in places with too wet or dry soil to support tree growth (e.g., bogs or river bluffs). Above the boreal band of elevation, subalpine and alpine vegetation dominate. The subalpine zone is a narrow transitional band between the boreal and the alpine life zones, and is characterized by sparse, stunted trees that can be considered tree line. In the subalpine, certain types of birch and willow shrub species grow at greater than or equal to one meter in height (commonly *Betula glandulosa* and *Salix pulchra*). In the alpine, trees no longer occur, and all shrubs are dwarf or lay prostrate on the ground. In this area, the boreal life zone occurs below 1600 feet elevation on average. The transition between boreal and subalpine vegetation can occur within a range of approximately 350 feet of elevation, and is highly dependent on slope, aspect, and shading from adjacent mountains.

Within each life zone, there are plant assemblages that are associated with cold slopes and warm slopes. Cold slopes and warm slopes are created by the combination of the steepness of the slope, the aspect, and shading from surrounding ridges and mountains. Warm slope positions occur on southeast to west facing slopes that are moderate to very steep (greater than 10 percent slope) and are not shaded by the surrounding landscape.

Cold slopes occur on northwest to east facing slopes, occur in shaded slope positions, or occur in low-lying areas that are cold air sinks. Examples of shaded positions include head slopes, low relief backslopes of hills, and the base of hills and mountains shaded by adjacent mountain peaks. Warm boreal slope soils have a cryic soil temperature regime and lack permafrost. In this area, white spruce forests are an indicator of warm boreal slopes. Cold boreal slope soils have a gelic soil temperature regime and commonly have permafrost. In this area, black spruce forests and woodlands are an indicator of cold boreal slopes. The boreal life zone can occur at higher elevations on warm slopes, and lower elevations on cold slopes.

Classification relationships

Landfire BPS – 7416211 – Western North American Boreal Black Spruce Dwarf-tree Peatland – Boreal Complex (Landfire 2009)

Ecological site concept

- Occurs in the boreal life zone on broad flat areas of stream terraces.
- Flooding does not occur. Ponding occurs frequently to occasionally for brief to long durations.
- Soils formed in a thin layer of windblown silts over silty and sandy alluvium.
- Soils are very deep but have permafrost at shallow to moderate depths.
- These are wet soils that are considered very poorly to poorly drained.
- The reference plant community is dwarf tree scrub woodland (Viereck et al. 1992) with black spruce the dominant tree. Multiple plant communities occur within the reference state related to wildfire.

Associated sites

F233XY131AK	Boreal Forest Gravelly Floodplain Occurs on high flood plains with productive white spruce forests.
R233XY130AK	Boreal Scrubland Gravelly Floodplain Occurs on low flood plains with willow and alder dominant plant communities.
R233XY207AK	Boreal Sedge Peat Depressions Occurs on depressions of floodplains stream terraces with sedge dominant plant communities.

Similar sites

F231XY171AK	Boreal Woodland Loamy Frozen Terraces Occurs in an adjacent area (MLRA 231X) on similar soils and is provisionally thought to have similar vegetation and disturbance dynamics.
-------------	---

F233XY111AK	Boreal Forest Loamy Frozen Slopes Both ecological sites have similar soils and vegetation. Site 171 occurs on stream terraces, while site 111 occurs on slopes of low-elevation mountains.
-------------	--

Table 1. Dominant plant species

Tree	(1) <i>Picea mariana</i>
Shrub	(1) <i>Ledum palustre ssp. decumbens</i> (2) <i>Vaccinium vitis-idaea</i>
Herbaceous	(1) <i>Eriophorum vaginatum</i> (2) <i>Sphagnum</i>

Physiographic features

- Occurs on broad flat areas of stream terraces.
- Associated with the boreal life zone. The representative elevation ranges between 150 and 1300 feet. Associated with cold slopes. Slopes are nearly level and can occur on all slope aspects.
- Flooding does not occur.
- Ponding occurs frequently to occasionally for brief to long durations of time. Ponding depth ranges up to 24 inches above the soil surface.
- These are wet soils with a seasonal water table occurring between 0 and 10 inches.
- Associated with negligible to very low amounts of runoff to adjacent, downslope ecological sites.

Table 2. Representative physiographic features

Geomorphic position, terraces	(1) Tread
Landforms	(1) Alluvial plain > Stream terrace (2) Alluvial plain > Stream terrace > Turf hummock
Runoff class	Negligible to very low
Flooding frequency	None
Ponding duration	Brief (2 to 7 days) to long (7 to 30 days)
Ponding frequency	Occasional to frequent
Elevation	46–396 m
Slope	0–2%
Ponding depth	0–30 cm
Water table depth	0–25 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding frequency	Not specified
Ponding duration	Not specified
Ponding frequency	Not specified
Elevation	Not specified
Slope	Not specified
Ponding depth	0–61 cm
Water table depth	Not specified

Climatic features

Short, warm summers and long, cold winters characterize the subarctic continental climate associated with this boreal forest gravelly slopes ecological site. The mean annual temperature for MLRA 233X ranges from 22 to 24 degrees Fahrenheit (PRISM 2008). The warmest months span May through August with mean normal maximum monthly temperatures ranging from 51 to 64 degrees Fahrenheit. The coldest months span December through March with mean normal minimum temperatures ranging from -2 to 3 degrees Fahrenheit. The freeze-free period for this boreal ecological site ranges from 105 to 129 days, and the temperature generally remains above freezing from late May through early-September.

The area receives minimal annual precipitation with July through September being the wettest. Average annual precipitation across MLRA 233X ranges between 17 to 21 inches (PRISM 2008). Approximately half of the annual precipitation occurs during the months of July through September with seasonal thunderstorms. The average annual snowfall ranges from 65 to 80 inches (USDA 2022). The ground is consistently covered with snow from November through March.

Table 4. Representative climatic features

Frost-free period (characteristic range)	73-105 days
Freeze-free period (characteristic range)	105-129 days
Precipitation total (characteristic range)	432-533 mm
Frost-free period (actual range)	26-111 days
Freeze-free period (actual range)	98-133 days
Precipitation total (actual range)	356-610 mm
Frost-free period (average)	90 days
Freeze-free period (average)	115 days
Precipitation total (average)	457 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) BETTLES AP [USW00026533], Bettles Field, AK

Influencing water features

This site is classified as an organic soil flats wetland under the Hydrogeomorphic (HGM) classification system (Smith et al. 1995; USDA-NRCS 2008). Precipitation is the main source of water (Smith et al. 1995).

Depth to the water table may decrease following summer storm events or spring snowmelt and increase during extended dry periods.

Wetland description

n/a

Soil features

- Soils formed in loess over sandy and silty alluvium.
- Rock fragments do not occur on the soil surface or in the soil profile.
- Capped with up to 19 inches of peat.
- The surface mineral horizon is a silt loam formed from loess or silty alluvium. This silty soil horizon is very thick
- While soils are very deep, permafrost occurs at shallow to moderate depths (18 to 32 inches).
- The pH of the soil profile ranges from very strongly acidic to slightly acidic.
- These are wet soils that are considered very poorly to poorly drained.
- Soils are classified as Gelisols in the great groups Historthels and Histoturbels.

Table 5. Representative soil features

Parent material	(1) Loess (2) Alluvium
Surface texture	(1) Peat
Family particle size	(1) Coarse-silty (2) Coarse-loamy
Drainage class	Poorly drained
Permeability class	Moderately rapid
Depth to restrictive layer	46–81 cm
Soil depth	152 cm
Surface fragment cover ≤3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	14.99–23.62 cm
Calcium carbonate equivalent (25.4-101.6cm)	0%

Clay content (0-50.8cm)	5–10%
Electrical conductivity (25.4-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (25.4-101.6cm)	0
Soil reaction (1:1 water) (25.4-101.6cm)	5–6.5
Subsurface fragment volume ≤3" (0-152.4cm)	0%
Subsurface fragment volume >3" (0-152.4cm)	0%

Table 6. Representative soil features (actual values)

Drainage class	Very poorly drained to poorly drained
Permeability class	Not specified
Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover ≤3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	14.99–39.12 cm
Calcium carbonate equivalent (25.4-101.6cm)	Not specified
Clay content (0-50.8cm)	Not specified
Electrical conductivity (25.4-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (25.4-101.6cm)	0–3
Soil reaction (1:1 water) (25.4-101.6cm)	5–7.2
Subsurface fragment volume ≤3" (0-152.4cm)	0–8%
Subsurface fragment volume >3" (0-152.4cm)	0–2%

Ecological dynamics

In the Upper Kobuk and Koyukuk Hills and Valleys MLRA (herein called area), fire is a common and natural event that has a significant control on the vegetation dynamics across the landscape. A typical fire event in the lands associated with this boreal forest silty frozen slopes ecological site will reset plant succession and alter dynamic soil properties (e.g., soil organic matter and depth of permafrost). For this ecological site to progress from the earliest stages of post-fire succession to the oldest stages of succession, data suggest that 70-130 years or more must elapse without another fire event (Johnstone et al. 2010a).

Within this area, wildfire is considered a natural and common event that typically goes unmanaged. Fire suppression is limited and occurs adjacent to the small villages spread throughout the area or on allotments with known structures, all of which have a relatively limited acre footprint. Most fires are caused by lightning strikes. From 2000 to 2020, 124 known fire events occurred in this area and the burn perimeter of the fires totaled approximately one million acres (AICC 2022). Fire-related disturbances are highly patchy and can leave undisturbed areas within the burn perimeter. During this time frame, 90 percent of the fire events were smaller than 20,000 acres but three fire events were greater than 100,000 acres in size (AICC 2022). Over this period of 20 years, these burn perimeters cover approximately 20 percent of this area.

The fire regime within Interior Alaska follows two general scenarios—low-severity burns and high-severity burns. It should be noted, however, that the fire regime in Interior Alaska can be considered more complex (Johnstone et al. 2008). Burn severity refers to the proportion of the vegetative canopy and organic material consumed in a fire event (Chapin et al. 2006). Fires in cool and moist habitat tend to result in low-severity burns, while fires in warm and dry habitat tend to result in high-severity burns. Because the soils have a thick organic cap and are poorly drained, the typical fire scenario for this ecological site is considered to result in a low-severity burn.

The low-severity fire regime predicted for this ecological site will have modest impacts to soil organic matter thickness, depth of permafrost, and soil drainage. While a low-severity fire can consume the bulk of above ground vegetation, minimal proportions of the organic mat are typically removed. Organic matter continues to insulate these cold soils. Field data from similar ecological sites support that each plant community has permafrost and that the associated low-severity fire event had a negligible impact on the depth of permafrost. If permafrost remains at similar depths after a fire event, then soil drainage is unlikely to improve post-fire.

In areas prone to low-severity fire events, the pre-fire vegetative community generally reestablishes quickly and there is minimal long-term alteration to community composition (Johnstone et al. 2010; Bernhardt et al. 2011). When minimal proportions of the organic mat are consumed, many species regenerate asexually using below ground root systems and rhizomes. Species known to regenerate after low-severity fire events include various graminoids (e.g., *Carex* spp. and *Eriophorum* spp.), forbs (e.g., *Equisetum* sp.), and shrubs (e.g., *Ledum groenlandicum*, *Vaccinium uliginosum*, *Salix* sp.) (Johnstone et al.


2010). Black spruce is the Interior Alaska tree species best adapted to a low-severity fire regime. Black spruce have semi-serotinous cones and a low-severity fire often results in a flush of black spruce seedlings at the burned location.

The later stages of succession have an overstory that is a mix of broadleaf and immature needleleaf trees (community 1.2) or mature needleleaf trees (community 1.1). The recruitment of trees species during the early stages of post-fire succession largely controls the composition of the stand of trees in the later stages of post-fire succession (Johnstone et al. 2010a). During these later stages of succession, the slower growing black spruce seedlings mature and eventually replace the shade-intolerant broadleaf tree species. The typical fire return interval for black spruce stands in the boreal forest is 70-130 years (Johnstone et al. 2010a).


Lands in Interior Alaska are burning more frequently than in the past, which may result in alternative states of succession. The historic fire return interval for white spruce stands in Interior Alaska occurs approximately once every 150 years (Landfire 2009; Abrahamson 2014). Global climate change is warming the boreal forest and causing a longer growing season. As a result, stands of spruce in the Alaskan boreal forest are burning more frequently and intensely than their historic averages (Kelly et al. 2013). Increases to burn frequency favors forested stands dominated by quick growing deciduous trees (community 1.3). A major reason being that increased fire frequency decreases the presence and abundance of mature, cone-bearing trees. Less mature trees result in less spruce seedlings post-fire and an overall decreased abundance of spruce in the developing forest canopy. Increased fire frequency paired with decreased spruce propagules lowers the chances for the site to progress to post-fire successional communities 1.2 and 1.1. Increased burn frequency in the boreal forest may result in alternative states of post-fire succession with stands of deciduous trees persisting for longer than normal durations of time (Johnstone et al. 2010b).

State and transition model

Ecosystem states



State 1 submodel, plant communities



1.1a - A low-severity fire sweeps through and incinerates much of the above ground vegetation

1.2a - Occurs 50 years after wildfire

1.2b - A low-severity fire sweeps through and incinerates much of the above ground vegetation

1.3a - Occurs 10 years after wildfire

State 1 Reference State

The reference plant community is dwarf tree scrub woodland (Vioreck et al. 1992) with the dominant tree being black spruce. There are three plant communities within the reference state related to fire. The vegetation modeled for this site has limited data and is considered provisional.

Dominant plant species

- black spruce (*Picea mariana*), tree
- tussock cottongrass (*Eriophorum vaginatum*), grass
- Bigelow's sedge (*Carex bigelowii*), grass
- sphagnum (*Sphagnum*), other herbaceous
- reindeer lichen (*Cladina*), other herbaceous

Community 1.1

black spruce / tussock cottongrass - Bigelow's sedge / Sphagnum - reindeer lichen

The reference plant community is characterized as dwarf tree scrub woodland (Vioreck et al. 1992) with black spruce as the dominant tree. Black spruce tree cover primarily occurs in the stunted tree stratum (greater than 50 years of age and less than 15 feet). Live

deciduous trees, primarily resin birch, occasionally occur in the tree canopy but with limited cover. The soil surface is primarily covered with herbaceous litter, moss, and lichen. Common understory species include marsh Labrador tea, lingonberry, dwarf birch, tussock cottongrass, Bigelow's sedge, cloudberry, various reindeer lichen, curled snow lichen (*Flavocetraria cucullata*), Sphagnum, and Schreber's big red stem moss. The understory vegetative strata that characterize this community are medium graminoids (between 4 and 24 inches), mosses, and lichens.

Dominant plant species

- black spruce (*Picea mariana*), tree
- resin birch (*Betula neoalaskana*), tree
- marsh Labrador tea (*Ledum palustre ssp. decumbens*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- dwarf birch (*Betula nana*), shrub
- tussock cottongrass (*Eriophorum vaginatum*), grass
- Bigelow's sedge (*Carex bigelowii*), grass
- cloudberry (*Rubus chamaemorus*), other herbaceous
- reindeer lichen (*Cladina*), other herbaceous
- sphagnum (*Sphagnum*), other herbaceous
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous

Community 1.2

black spruce / tussock cottongrass - Bigelow's sedge / Sphagnum

Community 1.2. is in the late stage of fire-induced secondary succession for this ecological site. It is characterized as needleleaf woodland (Viereck et al. 1992). Black spruce seedlings are abundant and tree cover primarily occurs in regenerative tree stratum. Deciduous tree seedlings, primarily resin birch, occasionally occur but with limited cover. The soil surface is primarily covered with mosses and herbaceous litter. Common understory species include Siberian alder, marsh Labrador tea, lingonberry, bog blueberry, scrub birch (*Betula glandulosa*), dwarf birch, leatherleaf, bog Labrador tea, tussock cottongrass, Bigelow's sedge, cloudberry, and Sphagnum. The understory vegetative strata that characterize this community are tree regeneration, low shrubs (between 8 and 36 inches), medium graminoids (between 4 and 24 inches), and mosses.

Dominant plant species

- black spruce (*Picea mariana*), tree
- resin birch (*Betula neoalaskana*), tree
- Siberian alder (*Alnus viridis ssp. fruticosa*), shrub
- marsh Labrador tea (*Ledum palustre ssp. decumbens*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- resin birch (*Betula glandulosa*), shrub
- dwarf birch (*Betula nana*), shrub

- leatherleaf (*Chamaedaphne calyculata*), shrub
- bog Labrador tea (*Ledum groenlandicum*), shrub
- tealeaf willow (*Salix pulchra*), shrub
- small cranberry (*Vaccinium oxycoccos*), shrub
- tussock cottongrass (*Eriophorum vaginatum*), grass
- Bigelow's sedge (*Carex bigelowii*), grass
- cloudberry (*Rubus chamaemorus*), other herbaceous

Community 1.3

dwarf birch - marsh Labrador tea / tussock cottongrass - Bigelow's sedge / Sphagnum

Community 1.3 is in the early stage of fire-induced secondary succession for this ecological site. This community is characterized as open low scrub (Viereck et al. 1992). Seedlings of black spruce and resin birch are commonly observed but have limited cover. Common species include dwarf birch, tealeaf willow, marsh Labrador tea, lingonberry, bog blueberry, bog Labrador tea, leatherleaf, crowberry, tussock cottongrass, Bigelow's sedge, cloudberry, and Sphagnum. The strata that characterize this community are low shrubs (between 8 and 36 inches), medium graminoids (between 4 and 24 inches), and mosses.

Dominant plant species

- dwarf birch (*Betula nana*), shrub
- tealeaf willow (*Salix pulchra*), shrub
- marsh Labrador tea (*Ledum palustre ssp. decumbens*), shrub
- lingonberry (*Vaccinium vitis-idaea*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- bog Labrador tea (*Ledum groenlandicum*), shrub
- leatherleaf (*Chamaedaphne calyculata*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- small cranberry (*Vaccinium oxycoccos*), shrub
- Siberian alder (*Alnus viridis ssp. fruticosa*), shrub
- tussock cottongrass (*Eriophorum vaginatum*), grass
- Bigelow's sedge (*Carex bigelowii*), grass
- bluejoint (*Calamagrostis canadensis*), grass
- sphagnum (*Sphagnum*), other herbaceous
- cloudberry (*Rubus chamaemorus*), other herbaceous
- reindeer lichen (*Cladina*), other herbaceous
- juniper polytrichum moss (*Polytrichum juniperinum*), other herbaceous
- horsetail (*Equisetum*), other herbaceous
- (*Flavocetraria cucullata*), other herbaceous
- felt lichen (*Peltigera aphthosa*), other herbaceous

Pathway 1.1a

Community 1.1 to 1.3

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated cold and wet soils, this site commonly experiences low-severity fires. Minimal proportions of the organic mat are typically removed. The pre-fire vegetation generally reestablishes quickly from below ground root systems and rhizomes.

Pathway 1.2a

Community 1.2 to 1.1

Community pathway 1.2a is thought to occur 50 years after wildfire (Landfire 2009). Black spruce seedlings and saplings mature into a stunted woodland.

Pathway 1.2b

Community 1.2 to 1.3

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated cold and wet soils, this site commonly experiences low-severity fires. Minimal proportions of the organic mat are typically removed. The pre-fire vegetation generally reestablishes quickly from below ground root systems and rhizomes.

Pathway 1.3a

Community 1.3 to 1.2

Community pathway 1.3a is thought to occur 10 years after fire (Landfire 2009). Black spruce seedlings and sapling start to become a characteristic component of the plant community.

Additional community tables

Animal community

not available

Hydrological functions

not available

Recreational uses

not available

Wood products

not available

Other products

not available

Other information

not available

Inventory data references

The vegetation modeled for this site has limited data and is considered provisional. The associated model was largely developed from NRCS staff with working knowledge of the area and literature review.

References

- Bernhardt, E.L., T.N. Hollingsworth, and . 2011. Fire severity mediates climate-driven shifts in understory community composition of black spruce stands of interior Alaska. *Journal of Vegetation Science* 22:32–44.
- Chapin, F.S., L.A. Viereck, P.C. Adams, K.V. Cleve, C.L. Fastie, R.A. Ott, D. Mann, and J.F. Johnstone. 2006. Successional processes in the Alaskan boreal forest. Page 100 in *Alaska's changing boreal forest*. Oxford University Press.
- Hinzman, L.D., L.A. Viereck, P.C. Adams, V.E. Romanovsky, and K. Yoshikawa. 2006. Climate and permafrost dynamics of the Alaskan boreal forest. *Alaska's changing boreal forest* 39–61.
- Johnstone, J.F., T.N. Hollingsworth, F.S. CHAPIN III, and M.C. Mack. 2010. Changes in fire regime break the legacy lock on successional trajectories in Alaskan boreal forest. *Global change biology* 16:1281–1295.
- Johnstone, J.F., F.S. Chapin, T.N. Hollingsworth, M.C. Mack, V. Romanovsky, and M. Turetsky. 2010. Fire, climate change, and forest resilience in interior Alaska. *Canadian Journal of Forest Research* 40:1302–1312.
- Johnstone, J.F. 2008. A key for predicting postfire successional trajectories in black spruce stands of interior Alaska. US Department of Agriculture, Forest Service, Pacific Northwest Research Station.

- Kelly, R., M.L. Chipman, P.E. Higuera, I. Stefanova, L.B. Brubaker, and F.S. Hu. 2013. Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National Academy of Sciences* 110:13055–13060.
- Landfire. 2009. Biophysical Setting. LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior, Washington, DC..
- Schoeneberger, P.J. and D.A. Wysocki. 2012. Geomorphic Description System. Natural Resources Conservation Service, 4.2 edition. National Soil Survey Center, Lincoln, NE.
- Smith, R.D., A.P. Ammann, C.C. Bartoldus, and M.M. Brinson. 1995. An approach for assessing wetland functions using hydrogeomorphic classification, reference wetlands, and functional indices.
- United States Department of Agriculture Staff. 2008. Hydrogeomorphic Wetland Classification System: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service.
- United States Department of Agriculture, . 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Other references

- PRISM Climate Group. 2018. Alaska – average monthly and annual precipitation and minimum, maximum, and mean temperature for the period 1981-2010. Oregon State University, Corvallis, Oregon. <https://prism.oregonstate.edu/projects/alaska.php>. (Accessed 4 September 2019).
- Scenarios network for Alaska and arctic planning (SNAP). Historical Monthly Temperature – 1km, 1901-2009. <http://ckan.snap.uaf.edu/dataset/>. (Accessed 5 May 2021).
- SNAP. Historical monthly and derived precipitation products downscaled from CRU TS data via the delta methods – 2km, 1901-2009. <http://ckan.snap.uaf.edu/dataset/>. (Accessed 5 May 2021).

Contributors

Blaine Spellman

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	03/21/2026
Approved by	Blaine Spellman
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production,**

not just forage annual-production):

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-