

Ecological site R233XY130AK

Boreal Scrubland Gravelly Floodplain

Last updated: 6/10/2025

Accessed: 03/16/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 233X–Upper Kobuk and Koyukuk Hills and Valleys

The Upper Kobuk and Koyukuk Hills and Valleys MLRA (herein called area) occurs in Interior Alaska. This area makes up 8,405 square miles. The largest tributaries are the Kobuk and the Koyukuk Rivers. Major tributaries of the Kobuk are the Reed, Beaver, Mauneluk, and Pau Rivers. Major tributaries of the Koyukuk River are the Alatna, John, and Kanuti Rivers. This area is primarily undeveloped wildland and sparsely populated. The communities within or near this area are Bettles, Kobuk, and Shungnak.

The terrain of this area consists of broad, nearly level river valleys and basins and rolling uplands separated by isolated hills and low rounded mountains. In the river valleys, nearly level flood plains and stream terraces gradually transition to gently sloping to moderately steep slopes leading to the hills and mountains. Basins are on the Pau River Flats between the eastern Zane and Lockwood Hills, on the Kanuti Flats between the Kanuti and Koyukuk Rivers, and along the middle reaches of the Hogatza River. Basins and stream terraces are dotted with hundreds of lakes and interconnecting wetlands. Elevation ranges from about 150 feet in the western part of the area, at the confluence of the Kobuk and Mauneluk Rivers, to 4,765 feet at the summit of Fritts Mountain, in the Angaycuham Mountains.

Geology and Soils

The northern part of the area was covered repeatedly by Pleistocene glaciers originating in the Brooks Range to the north. Slightly modified to highly modified moraines and drift cover many of the rolling uplands. Glacial ice flowed over most of the hills and low mountains, removing existing deposits and leaving a thin layer of glacial deposits. Today,

the lower mountain slopes, hills, and valley bottoms are covered with a variety of material, including glacial drift, colluvium, slope alluvium, fluvial deposits, and silty loess. In the southern part of the area, basins and valleys are filled with Quaternary glaciofluvial and fluvial deposits. Hills and upland slopes are covered with bedrock colluvium and slope alluvium, which are mantled with loess in places. The bedrock geology underlying much of the area consists dominantly of Permian through Lower Cretaceous stratified sedimentary and volcanic rocks.

This area is in the zone of discontinuous permafrost. Permafrost is close to the surface in lands with finer textured sediments throughout the area. Isolated masses of ground ice occur on terraces and the lower side slopes of hills. Permafrost does not occur on flood plains, on steep south-facing slopes, or other lands with very gravelly soils. Periglacial features, such as thermokarst pits, peat plateaus, and earth hummocks, are on the lower hill and mountain slopes and in upland valleys.

The dominant soil orders in this area are Gelisols, Inceptisols, and Entisols. The Gelisols are shallow or moderately deep to permafrost, occur on finer textured sediments, and are poorly drained or very poorly drained. Common Gelisol suborders are Histels, Orthels, and Turbels. The Histels have thick accumulations of surface organic material and occur in depressions, lake margins, and peat plateau. The Orthels and Turbels have comparably thinner surface organic material and occur on stream terraces and hill and upland slopes. The Inceptisols and Entisols are typically associated with gravelly soils that do not have permafrost within their profile, are deep, and are somewhat poorly drained to well drained. The common Inceptisol suborders are Cryepts and Gelepts both of which occur on upland and mountain slopes. Cryepts occur under forested soils at lower elevations and Gelepts on alpine tundra at higher elevations. Common Entisol suborders are Cryofluvents and Cryorthents both of which occur on alluvium on flood plains. Miscellaneous (non-soil) areas make up about 8 percent of this MLRA. The most common are rock outcrop, rubble land, and water.

Wildfires disturb the insulating organic material at the soil surface and can change the presence and/or depth of permafrost in the soil profile. These fire related changes to permafrost can also change the depth and presence of perched water tables. Gelisols that burn in this area can change soil taxonomic classification. For instance, depending on fire-severity, Histels may change to Orthels and Orthels may change to Inceptisols. Depending on the frequency and intensity of fires, landform position, and soil texture, the soils may or may not revert back to their original taxonomic classification.

Climate

Short, warm summers and long, cold winters characterize the continental subarctic climate of the area. The average annual precipitation ranges from 15 to 19 inches on valley bottoms and basins and from 19 to 26 inches at the higher elevations in the hills and mountains (PRISM 2018). Most of the precipitation falls as rain between May and September. The average annual snowfall ranges from 65 to 80 inches. The average

annual temperature is 22 to 24 degrees Fahrenheit (PRISM 2018). The temperature normally remains above freezing from mid-June through August in river valleys and basins with a freeze-free period ranging from 109 to 125 days. The freeze-free period is significantly shorter on higher elevation mountain slopes.

Vegetation

Most of this area is forested below an elevation of 1600 feet. Dominant tree species on slopes are white spruce and black spruce. Black spruce stands dominate on north-facing slopes, stream terraces, and other sites with poor drainage and permafrost. White spruce stands dominate on steep, south-facing slopes with dry soils. At lower elevations, lightning-caused wildfires are common, often burning many thousands of acres during a single fire event. Following wildfires, forbs, grasses, willow, ericaceous shrubs, paper birch, and quacking aspen communities are common until they are eventually replaced by stands of spruce. Tall willow and alder scrub is extensive on low flood plains. White spruce and balsam poplar are common on high flood plains.

With increasing elevation, the forests and woodlands give way to subalpine communities dominated by krummholz spruce, shrub birch, willow, and ericaceous shrubs. At even higher elevations, alpine communities prevail which are characterized by diverse forbs, dwarf ericaceous shrubs, and eightpetal mountain-avens. Many of these high elevation communities have a considerable amount of lichen cover and bare ground.

LRU notes

In this area, we refer to three life zones that are defined by the physiological limits of plant communities along an elevational gradient: boreal, subalpine, and alpine. The boreal life zone is the elevational band where forest communities dominate. Not all areas in the boreal life zone are forest communities, however, particularly in places with too wet or dry soil to support tree growth (e.g., bogs or river bluffs). Above the boreal band of elevation, subalpine and alpine vegetation dominate. The subalpine zone is a narrow transitional band between the boreal and the alpine life zones, and is characterized by sparse, stunted trees that can be considered tree line. In the subalpine, certain types of birch and willow shrub species grow at greater than or equal to one meter in height (commonly *Betula glandulosa* and *Salix pulchra*). In the alpine, trees no longer occur, and all shrubs are dwarf or lay prostrate on the ground. In this area, the boreal life zone occurs below 1600 feet elevation on average. The transition between boreal and subalpine vegetation can occur within a range of approximately 350 feet of elevation, and is highly dependent on slope, aspect, and shading from adjacent mountains.

Within each life zone, there are plant assemblages that are associated with cold slopes and warm slopes. Cold slopes and warm slopes are created by the combination of the steepness of the slope, the aspect, and shading from surrounding ridges and mountains. Warm slope positions occur on southeast to west facing slopes that are moderate to very steep (greater than 10 percent slope) and are not shaded by the surrounding landscape.

Cold slopes occur on northwest to east facing slopes, occur in shaded slope positions, or occur in low-lying areas that are cold air sinks. Examples of shaded positions include head slopes, low relief backslopes of hills, and the base of hills and mountains shaded by adjacent mountain peaks. Warm boreal slope soils have a cryic soil temperature regime and lack permafrost. In this area, white spruce forests are an indicator of warm boreal slopes. Cold boreal slope soils have a gelic soil temperature regime and commonly have permafrost. In this area, black spruce forests and woodlands are an indicator of cold boreal slopes. The boreal life zone can occur at higher elevations on warm slopes, and lower elevations on cold slopes.

Classification relationships

Landfire BPS – 6916141 – Western North American Boreal Montane Floodplain Forest and Shrubland – Boreal

Landfire BPS – 7416150 – Western North American Boreal Lowland Large River Floodplain Forest and Shrubland (Landfire 2009)

Ecological site concept

- Occurs in the boreal life zone on low floodplains.
- Soils formed in alluvium.
- Flooding occurs frequently for very long to long durations. Ponding does not occur.
- Soils are considered poorly to somewhat poorly drained.
- Soils are very deep without restrictions. Permafrost does not occur in the soil profile.
- The reference plant community is closed tall scrub (Viereck et al. 1992) with feltleaf willow the dominant overstory shrub. Multiple plant communities occur within the reference state related to flooding.

Associated sites

F233XY131AK	Boreal Forest Gravelly Floodplain Occurs on high flood plains with white spruce dominant plant communities.
F233XY171AK	Boreal Woodland Loamy Frozen Terraces Occurs on stream terraces with stands of black spruce.
R233XY207AK	Boreal Sedge Peat Depressions Occurs on depressions of flood plains and stream terraces with sedge dominant plant communities.

Similar sites

R231XY130AK	Boreal Scrubland Gravelly Floodplain Occurs in an adjacent area (MLRA 231X) on similar soils and is provisionally thought to have similar vegetation and disturbance dynamics.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Salix alaxensis</i>
Herbaceous	(1) <i>Calamagrostis canadensis</i>

Physiographic features

- Occurs on low flood plains.
- Associated with the boreal life zone. Representative elevation occurs between 150 and 1300 feet.
- Slopes are nearly level and occur on all aspects.
- Ponding does not occur.
- Flooding occurs frequently for long to very long durations.
- These are moist to wet soils with a seasonal water table occurring between 0 and 20 inches.
- Associated with negligible amounts of runoff to adjacent, downslope ecological sites.

Table 2. Representative physiographic features

Landforms	(1) Alluvial plain > Flood plain
Runoff class	Negligible
Flooding duration	Long (7 to 30 days) to very long (more than 30 days)
Flooding frequency	Frequent
Ponding frequency	None
Elevation	46–396 m
Slope	0–2%
Water table depth	25–51 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
Flooding duration	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	46–488 m
Slope	Not specified
Water table depth	0–51 cm

Climatic features

Short, warm summers and long, cold winters characterize the subarctic continental climate associated with this boreal forest gravelly slopes ecological site. The mean annual temperature for MLRA 233X ranges from 22 to 24 degrees Fahrenheit (PRISM 2008). The warmest months span May through August with mean normal maximum monthly temperatures ranging from 51 to 64 degrees Fahrenheit. The coldest months span December through March with mean normal minimum temperatures ranging from -2 to 3 degrees Fahrenheit. The freeze-free period for this boreal ecological site ranges from 105 to 129 days, and the temperature generally remains above freezing from late May through early-September.

The area receives minimal annual precipitation with July through September being the wettest. Average annual precipitation across MLRA 233X ranges between 17 to 21 inches (PRISM 2008). Approximately half of the annual precipitation occurs during the months of July through September with seasonal thunderstorms. The average annual snowfall ranges from 65 to 80 inches (USDA 2022). The ground is consistently covered with snow from November through March.

Table 4. Representative climatic features

Frost-free period (characteristic range)	73-105 days
Freeze-free period (characteristic range)	105-129 days
Precipitation total (characteristic range)	432-533 mm
Frost-free period (actual range)	26-111 days
Freeze-free period (actual range)	98-133 days
Precipitation total (actual range)	356-610 mm
Frost-free period (average)	90 days
Freeze-free period (average)	115 days
Precipitation total (average)	457 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) BETTLES AP [USW00026533], Bettles Field, AK

Influencing water features

In the associated flood plains, overbank flow from the channel and subsurface hydraulic connections between the stream and adjacent wetlands are the main sources of water (Smith et al. 1995).

Depth to the water table may decrease following summer storm events or spring snowmelt and increase during extended dry periods.

Wetland description

This ecological site is classified as a riverine wetland under the Hydrogeomorphic (HGM) classification system (Smith et al. 1995; USDA-NRCS 2008).

Soil features

- Soils formed in alluvium.
- Rock fragments on the soil surface range between 0 and 75 percent cover.
- Capped with up to one inch of organic material.
- The surface mineral horizon is stratified silts and fine sands. The thickness of these finer sediment horizons is highly variable and occur over bands of sandy and gravelly alluvium.
- These gravelly soils have subsurface rock fragments ranging between 30 and 70 percent of the soil profile by volume.
- While soils are very deep, strong contrasting textural stratification occurs at very shallow to shallow depths (3 to 13 inches). This restriction can affect the movement and retention of water and/or nutrients.
- The pH of the soil profile ranges from slightly acidic to moderately alkaline.
- These are moist to wet soils that are considered poorly to somewhat poorly drained.

Table 5. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Fine sand (2) Silt
Family particle size	(1) Coarse-loamy over sandy or sandy-skeletal (2) Sandy-skeletal
Drainage class	Poorly drained to somewhat poorly drained
Permeability class	Rapid
Depth to restrictive layer	8–33 cm
Soil depth	152 cm
Surface fragment cover ≤3"	0–20%
Surface fragment cover >3"	3–55%

Available water capacity (0-101.6cm)	2.03–8.64 cm
Calcium carbonate equivalent (25.4-101.6cm)	0–5%
Clay content (0-50.8cm)	4–6%
Electrical conductivity (25.4-101.6cm)	0–1 mmhos/cm
Sodium adsorption ratio (25.4-101.6cm)	0
Soil reaction (1:1 water) (25.4-101.6cm)	6.2–8.4
Subsurface fragment volume ≤3" (0-152.4cm)	20–50%
Subsurface fragment volume >3" (0-152.4cm)	10–20%

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Permeability class	Not specified
Depth to restrictive layer	8–79 cm
Soil depth	Not specified
Surface fragment cover ≤3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	0.51–8.64 cm
Calcium carbonate equivalent (25.4-101.6cm)	Not specified
Clay content (0-50.8cm)	Not specified
Electrical conductivity (25.4-101.6cm)	Not specified
Sodium adsorption ratio (25.4-101.6cm)	0–3
Soil reaction (1:1 water) (25.4-101.6cm)	5.8–8.4
Subsurface fragment volume ≤3" (0-152.4cm)	0–50%

Subsurface fragment volume >3" (0-152.4cm)	0–35%
---	-------

Ecological dynamics

All montane streams and rivers in the Upper Kobuk and Koyukuk Hills and Valley MLRA (herein called area) have low and/or high flood plain ecological sites. These flood plain ecological sites represent major breaks in the flood regime and dominant vegetative type on associated tributaries. The low flood plain ecological site is thought to flood frequently (greater than 50 times in 100 years) for brief to long durations of time (2 to 30 days) and supports shrub dominant communities. The high flood plain ecological site floods occasionally to rarely (1 to 50 times in 100 years) for brief durations of time (2 to 7 days) and supports forested plant communities.


The shift of vegetative type from shrubland to forest represents riparian primary succession along major streams in the area. On other Interior Alaska flood plains, this successional process is thought to take between 200 and 300 years (Chapin et al. 2006). The flood regime, growth traits of vegetation, biotic competition, and a slew of other factors contribute to the dynamic nature of boreal flood plain succession. For more detailed information on boreal flood plain succession and successional drivers, refer to Walker et al. (1986) and Chapin et al. (2006).

Data indicates that differences in flood frequency and duration result in different plant communities for this ecological site. Low floodplain positions thought to flood more frequently have significant decreases to willow and alder height and cover. Given this observation, more frequently and severely flooded plant communities were incorporated into the reference state (community 1.2 and 1.3). These plant communities represent the successional transition from river wash to the reference plant community (community 1.1).


These montane streams and rivers have a stream terrace ecological site (see F231XY171AK). When compared to montane flood plains, stream terraces occur on higher landform positions that are often further away from the active stream channel. These montane stream terraces no longer flood. Stream terraces have thick peat layers, contact permafrost at shallow to moderate depths, commonly pond, and have wetter soils. Stream terraces support much less productive stands of black spruce (*Picea mariana*).

State and transition model

Ecosystem states



State 1 submodel, plant communities



1.1a - More frequent and longer duration flooding.

1.2a - Less frequent and shorter duration flooding.

1.2b - More frequent and longer duration flooding.

1.3a - Less frequent and shorter duration flooding.

State 1 Reference State

Multiple plant communities occur within the reference state and the vegetation differs in large part due to flooding. The frequent flooding that occurs for long durations of time prevents the establishment of trees and development of forested plant communities. For this site, plant community 1.1 has the least frequent and shortest duration flood events. This community is characterized as closed tall scrub (Vioreck et al. 1992) with the dominant tall scrub being felleaf willow. If flooding becomes less frequent and lasts for shorter durations of time, balsam poplar and white spruce gain dominance and the ecological site shifts to the high flood plain. The vegetation modeled for this site has limited data and is considered provisional.

Dominant plant species

- felleaf willow (*Salix alaxensis*), shrub
- bluejoint (*Calamagrostis canadensis*), grass

Community 1.1 felleaf willow / bluejoint

The reference plant community is characterized as closed tall scrub (Vioreck et al. 1992), which is primarily composed of felleaf willow. Within the reference state for this site, this

community has the least severe flood regime. Balsam poplar and white spruce seedling, saplings, and immature trees are common but are not dominant in the overstory. Other commonly observed species include thinleaf alder, prickly rose, squashberry, bluejoint, northern bedstraw, Tilesius' wormwood, tall bluebells, and horsetail. The vegetative strata that characterize this community phase are tall shrubs (greater than 10 feet) and tall graminoids (greater than 2 feet). The soil surface is primarily covered with herbaceous litter and woody debris, but large patches of exposed bare soil can occur (as much as 60 percent of plot).

Dominant plant species

- feltleaf willow (*Salix alaxensis*), shrub
- prickly rose (*Rosa acicularis*), shrub
- thinleaf alder (*Alnus incana ssp. tenuifolia*), shrub
- squashberry (*Viburnum edule*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- northern bedstraw (*Galium boreale*), other herbaceous
- Tilesius' wormwood (*Artemisia tilesii*), other herbaceous
- horsetail (*Equisetum*), other herbaceous

Community 1.2

feltleaf willow / bluejoint / northern bedstraw - Tilesius' wormwood

Community 1.2 is characterized as closed low scrub (Vioreck et al. 1992) primarily composed of feltleaf willow. Balsam poplar seedlings and saplings are common but are not a dominant overstory species. Commonly observed species include thin leaf alder, Siberian alder, prickly rose, bluejoint, northern bedstraw, fireweed, Tilesius' wormwood, and field horsetail. The vegetative strata that characterize this community are medium shrubs (between 3 and 10 feet), medium graminoids (between 4 and 24 inches), and medium forbs (between 4 and 24 inches). The soil surface is primarily covered with herbaceous litter and woody debris, but large patches of exposed bare soil and surface rock fragments can occur (as much as 100 percent of plot).

Dominant plant species

- feltleaf willow (*Salix alaxensis*), shrub
- Siberian alder (*Alnus viridis ssp. fruticosa*), shrub
- thinleaf alder (*Alnus incana ssp. tenuifolia*), shrub
- prickly rose (*Rosa acicularis*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- northern bedstraw (*Galium boreale*), other herbaceous
- fireweed (*Chamerion angustifolium*), other herbaceous
- Tilesius' wormwood (*Artemisia tilesii*), other herbaceous
- field horsetail (*Equisetum arvense*), other herbaceous

Community 1.3

feltleaf willow - thinleaf alder / dwarf fireweed - Tilesius' wormwood

Community 1.3 is sparsely vegetated and is often characterized as open low scrub (Viereck et al. 1992) with dominant shrubs being feltleaf willow and thinleaf alder. Within the reference state for this site, this community has the most severe flood regime. Balsam poplar and white spruce seedlings and saplings are common but are not a dominant overstory species. This community is highly diverse. Commonly observed species include shrubby cinquefoil, bluejoint, various bluegrass species, tufted wheatgrass, tufted hairgrass, dwarf fireweed, Tilesius' wormwood, alpine sweetvetch, and arctic lupine. The vegetative strata that characterize this community are regenerating trees (less than 15 feet), low shrubs (between 8 and 36 inches), medium forbs (between 4 and 24 inches), and medium graminoids (between 4 and 24 inches). Large patches of exposed bare soil and rock fragments are common (as much as 100 percent of plot).

Dominant plant species

- feltleaf willow (*Salix alaxensis*), shrub
- thinleaf alder (*Alnus incana* ssp. *tenuifolia*), shrub
- shrubby cinquefoil (*Dasiphora fruticosa*), shrub
- Siberian alder (*Alnus viridis* ssp. *fruticosa*), shrub
- prickly rose (*Rosa acicularis*), shrub
- bluegrass (*Poa*), grass
- tufted hairgrass (*Deschampsia cespitosa*), grass
- tufted wheatgrass (*Elymus macrourus*), grass
- bluejoint (*Calamagrostis canadensis*), grass
- spike trisetum (*Trisetum spicatum*), grass
- dwarf fireweed (*Chamerion latifolium*), other herbaceous
- Tilesius' wormwood (*Artemisia tilesii*), other herbaceous
- alpine sweetvetch (*Hedysarum alpinum*), other herbaceous
- arctic lupine (*Lupinus arcticus*), other herbaceous
- common yarrow (*Achillea millefolium*), other herbaceous
- fireweed (*Chamerion angustifolium*), other herbaceous
- elegant hawkbeard (*Crepis elegans*), other herbaceous
- field horsetail (*Equisetum arvense*), other herbaceous
- dwarf scouringrush (*Equisetum scirpoides*), other herbaceous
- bitter fleabane (*Erigeron acris*), other herbaceous
- arctic aster (*Eurybia sibirica*), other herbaceous
- marsh grass of Parnassus (*Parnassia palustris*), other herbaceous
- merckia (*Wilhelmsia physodes*), other herbaceous

Pathway 1.1a

Community 1.1 to 1.2

More frequent and longer duration flooding. The reference state for this ecological site

floods frequently for long periods of time (> 50 times in 100 years). Areas that are thought to flood less frequently are represented by community 1.1 and areas that are thought to flood more frequently are represented by community 1.2 and 1.3. When compared to community 1.1, the more frequently flooded plant community (community 1.2) shorter shrubs and less willow cover.

Pathway 1.2a

Community 1.2 to 1.1

Less frequent and shorter duration flooding. Areas that are thought to flood less frequently are represented by community 1.1 and areas that are thought to flood more frequently are represented by community 1.2. When compared to community 1.1, the more frequently flooded plant community has shorter shrubs and less willow cover.

Pathway 1.2b

Community 1.2 to 1.3

More frequent and longer duration flooding. Areas that are thought to flood less frequently are represented by community 1.2 and areas that are thought to flood more frequently are represented by community 1.3. When compared to community 1.2, the more frequently flooded plant communities have less shrub cover.

Pathway 1.3a

Community 1.3 to 1.2

Less frequent and shorter duration flooding. Areas that are thought to flood less frequently are represented by community 1.2 and areas that are thought to flood more frequently are represented by community 1.3. When compared to community 1.2, the more frequently flooded plant community has less willow and alder cover.

Additional community tables

Animal community

not available

Hydrological functions

not available

Recreational uses

not available

Wood products

not available

Other products

not available

Other information

not available

Inventory data references

The vegetation modeled for this site has limited data and is considered provisional. The associated model was largely developed from NRCS staff with working knowledge of the area and literature review.

References

Abrahamson, I.L. 2014. Fire Regimes of Alaskan White Spruce Communities.

Chapin, F.S., L.A. Viereck, P.C. Adams, K.V. Cleve, C.L. Fastie, R.A. Ott, D. Mann, and J.F. Johnstone. 2006. Successional processes in the Alaskan boreal forest. Page 100 in Alaska's changing boreal forest. Oxford University Press.

Landfire. 2009. Biophysical Setting. LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior, Washington, DC..

Schoeneberger, P.J. and D.A. Wysocki. 2017. Geomorphic Description System, Version 5.0..

United States Department of Agriculture, . 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Walker, L.R., J.C. Zasada, and F.S. Chapin III. 1986. The role of life history processes in primary succession on an Alaskan floodplain. *Ecology* 67:1243–1253.

Other references

PRISM Climate Group. 2018. Alaska – average monthly and annual precipitation and minimum, maximum, and mean temperature for the period 1981-2010. Oregon State University, Corvallis, Oregon. <https://prism.oregonstate.edu/projects/alaska.php>. (Accessed 4 September 2019).

Scenarios network for Alaska and arctic planning (SNAP). Historical Monthly Temperature – 1km, 1901-2009. <http://ckan.snap.uaf.edu/dataset/>. (Accessed 5 May 2021).

SNAP. Historical monthly and derived precipitation products downscaled from CRU TS data via the delta methods – 2km, 1901-2009. <http://ckan.snap.uaf.edu/dataset/>. (Accessed 5 May 2021).

Contributors

Blaine Spellman

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	03/16/2026
Approved by	Blaine Spellman
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-