

Ecological site R234XY702AK

Subalpine scrub gravelly slopes

Last updated: 6/09/2025

Accessed: 03/18/2026

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 234X–Interior Brooks Range Mountains

Geography

The Interior Brooks Range mountains area consists of predominantly steep, jagged mountains and narrow valleys that drain the southern side of the Brooks Range. This area is bordered by the Northern Brooks Range Mountains (Major Land Resource Area (MLRA) 244) to the north, the Western Brooks Range Mountains Foothills and Valleys (MLRA 243) to the west, and the Upper Kobuk and Koyukuk Hills and Valleys (MLRA 233) and the Interior Alaska Highlands (MLRA 231) to the south. The Brooks Range represents a drainage divide that is also the dividing line between MLRA 234 and MLRA 244. Rivers draining to the north wind their way along the North Slope until they reach the Arctic Ocean, while rivers that drain to the south reach the northern Pacific Ocean via the Bering Sea. MLRA 234 covers an area of 22,479 miles and is sparsely populated (USDA, 2022).

Geology

The Brooks Range is the northernmost extension of the Rocky Mountains and the highest range within the Arctic Circle, with high peaks in the eastern part of the range reaching elevations of nearly 9,000 feet. This area falls within the zone of discontinuous permafrost, with the continuous permafrost zone primarily occurring north of the Brooks Range. Wide, U-shaped valleys are evidence of extensive glaciation in the Early and Mid-Pleistocene, with most glaciers retreating to their current, high-elevation positions by the Late Pleistocene. The characteristically sharp upper peaks give way to lower mountain slopes comprised of alluvial and colluvial fans before reaching the gently sloping flood plains and terraces of the valley bottoms. While Paleozoic and Jurassic igneous and volcanic rocks can be found in the eastern part of the range, most of the lower slopes of this area are comprised of modified glacial material, alluvial, and colluvial deposits. Many rivers and

streams, such as the Koyukuk, have their headwaters in the Brooks Range and drain to the Bering Sea and North Pacific Ocean via the Yukon River.

Soils

The dominant soils orders in this MLRA are Gelisols, Entisols, and Inceptisols. Soils in the area have a gelic (subgelic) or cryic temperature regime, a udic or aquic moisture regime, and mixed minerology. Gelisols are common on soils that are shallow or moderately deep to permafrost and are somewhat poorly to very poorly drained. Gelisols are more common on cold slopes and stream terraces. In some cases, higher-intensity wildfires lead to loss of insulation when the surface organic layer is burned. This can lead to permafrost loss or active layer expansion and ultimately alter hydrology and taxonomic classification. Entisols and Inceptisols lacking in permafrost range from excessively-well to poorly drained. Entisols and Inceptisols are more common on rocky terrain, warm boreal slopes, and flood plains. Miscellaneous areas such as glaciers, riverwash, rock outcrop, and rubble land make up 63 percent of the MLRA.

Vegetation

The continental subarctic climate of the Brooks Range in conjunction with shallow, rocky soils leads to a sparsely forested landscape in this MLRA. Spruce-hardwood forests and woodlands tend to be relegated to lower elevations where deeper soils form on foot slopes and terraces. This is contrasted by the ericaceous dwarf shrub communities that are abundant on shallow, rocky slopes and ridges. Exposed sites are predominantly covered in lichen and sporadic forbs. Black spruce (*Picea mariana*) woodlands and tussock-forming sedge communities are on high stream terraces and foot slopes where permafrost occurs, as are wet sedge meadows. Floodplains tend to be dominated by low to tall willow scrub communities.

Land use

Except for areas along the Dalton Highway, access to most of this MLRA is extremely limited, lending itself to intact natural vegetation communities. For this reason, land use primarily takes the form of subsistence hunting, gathering, and fishing by local communities. In addition to subsistence activity, the Brooks Range is also a popular recreation destination, with many users utilizing air taxi and guiding services to access remote parts of the area. As is the case with much of interior Alaska, major resource concerns involve the persistence of permafrost, the degradation of which can lead to various changes in hydrology and nutrient cycling.

LRU notes

This area supports three life zones defined by the physiological limits of plant communities along an elevational gradient: boreal, subalpine, and alpine. The boreal life zone is the elevational band where forest communities dominate. Not all areas in the boreal life zone are forest communities, however, particularly in places where soil is too wet or dry to support tree growth (e.g., bogs or river bluffs). Above the boreal band of elevation, subalpine and alpine vegetation dominate. The subalpine zone is typically a narrow

transitional band between the boreal and the alpine life zones, and is characterized by sparse, stunted trees. In the subalpine, certain types of birch and willow shrub species grow at taller than one meter in height (commonly *Betula glandulosa* and *Salix pulchra*). In the alpine, trees no longer occur, and all shrubs are dwarf or lay prostrate on the ground. In this area, the boreal life zone occurs below 2,500 feet elevation on average. The transition between boreal and alpine vegetation can occur within a range of elevations, and is highly dependent on slope, aspect, and shading from adjacent mountains.

Within each life zone, there are plant assemblages that are typically associated with cold slopes and warm slopes. Cold slopes and warm slopes are created by the combination of the steepness of the slope, the aspect, and shading from surrounding ridges and mountains. Warm slope positions typically occur on southeast to west facing slopes that are moderate to very steep (greater than ten percent slope) and are not shaded by the surrounding landscape. Cold slopes typically occur on northwest to east facing slopes, occur in shaded slope positions, or occur in low-lying areas that are cold air sinks. Examples of shaded positions include head slopes, low relief backslopes of hills, and the base of hills and mountains shaded by adjacent mountain peaks. Warm boreal slope soils have a cryic soil temperature regime and lack permafrost. In this area, white spruce (*Picea glauca*) forests are an indicator of warm boreal slopes. Cold boreal slope soils typically have a gelic soil temperature regime and commonly have permafrost. In this area, black spruce forests and woodlands are an indicator of cold boreal slopes. The boreal life zone can occur at higher elevations on warm slopes, and lower elevations on cold slopes.

Classification relationships

Alaska Vegetation Classification
Closed low shrub (II.C.1. – level III)
(Viereck et al. 1992)

LANDFIRE Biophysical Settings
7416011 – Western North American Boreal Treeline White Spruce Woodland - Boreal
(LANDFIRE biophysical settings, 2009)

Ecological site concept

Key soil and site characteristics

- Occurs in the subalpine zone at elevations ranging 2,050 to 3,360 feet
- Warm slopes (southeast to west facing)
- Loamy-skeletal, well drained
- Moderate slopes (4 to 23 percent)

Associated sites

F234XY714AK	Boreal forest gravelly slopes Occurs downslope on similar soils in the boreal life zone.
-------------	----------------------------------------------------------------------------------------------------

Similar sites

F234XY714AK	Boreal forest gravelly slopes This ecological site occurs in the boreal life zone. R234XY702AK and F234XY714AK share similar vegetation but F234XY714AK has more productive white spruce (<i>Picea glauca</i>) and forested stands.
-------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Betula glandulosa</i> (2) <i>Vaccinium uliginosum</i>
Herbaceous	(1) <i>Pleurozium schreberi</i> (2) <i>Hylocomium splendens</i>

Physiographic features

This subalpine ecological site occurs on mountain slopes and glaciated plains at high elevations typically ranging 2,050 to 3,360 feet. Slopes range from 4 to 23 percent. This ecological site is typically associated with warm slopes that are southeast to west facing. Flooding and ponding are not known to occur and there is no high-water table in the soil profile.

Table 2. Representative physiographic features

Geomorphic position, mountains	(1) Center third of mountainflank (2) Lower third of mountainflank
Landforms	(1) Mountains > Mountain slope (2) Plains > Plain
Runoff class	Very low
Flooding frequency	None
Ponding frequency	None
Elevation	625–1,024 m
Slope	4–23%
Aspect	W, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Runoff class	Not specified
--------------	---------------

Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	271–1,804 m
Slope	0–65%

Climatic features

Characterized by a continental subarctic climate, cool, short summers interspace long, cool winters in this Major Land Resource Area (MLRA). Average annual temperatures range from 8 to 16 degrees, with freezing temperatures possible throughout the year. Precipitation ranges from 10 inches at low elevations up to 30 inches in the high elevations, with average snowfall ranging from 60 to 100 inches, annually. On average, there are only four frost free days per year at lower elevations, with frost possible throughout the year at higher elevations. The average high temperature in July (the warmest month, on average) is 65 degrees F, while the average high temperature in January is -1 degrees F. Extreme lows are common throughout interior Alaska. The lowest temperature recorded in Bettles Field, the location of one of the weather stations below, reached -69 degrees F.

Table 4. Representative climatic features

Frost-free period (characteristic range)	3-4 days
Freeze-free period (characteristic range)	48-59 days
Precipitation total (characteristic range)	254-356 mm
Frost-free period (actual range)	3-4 days
Freeze-free period (actual range)	46-61 days
Precipitation total (actual range)	229-356 mm
Frost-free period (average)	4 days
Freeze-free period (average)	54 days
Precipitation total (average)	305 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) WISEMAN [USC00509869], Bettles Field, AK
- (2) CHANDALAR SHELF DOT [USC00501497], Southerly North Slope Bo, AK
- (3) CHANDALAR LAKE [USC00501492], Yukon Flats Nat Wildlife, AK

Influencing water features

Due to landscape position, no streams or wetlands are associated with this ecological site. Precipitation and throughflow are the main sources of water.

Soil features

The soils of this ecological site formed in windblown silts and colluvium and do not have permafrost. Surface rock fragments are not typically high, but in some cases make up 33 percent cover. The mineral soil is capped with a thin organic horizon of one to two inches. The mineral soil below is a silt loam and contains higher amount of rock fragments than at the surface. Below the silty layer is gravelly colluvium which can act as a strongly contrasting textural stratification when rock fragments are present, causing root restriction within four to ten inches of the soil surface, but soils range from deep to very deep. Soils are extremely acidic to slightly acidic and are well drained.

Table 5. Representative soil features

Parent material	(1) Loess (2) Colluvium
Surface texture	(1) Sandy loam
Family particle size	(1) Loamy-skeletal
Drainage class	Well drained
Permeability class	Very rapid
Depth to restrictive layer	10–25 cm
Soil depth	112–152 cm
Surface fragment cover ≤3"	0–33%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	11.68–17.02 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (25.4-101.6cm)	4.1–6.3
Subsurface fragment volume ≤3" (0-152.4cm)	12–36%

Subsurface fragment volume >3" (0-152.4cm)	5–54%
-----------------------------------------------	-------

Table 6. Representative soil features (actual values)

Drainage class	Not specified
Permeability class	Not specified
Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	Not specified
Calcium carbonate equivalent (0-101.6cm)	Not specified
Electrical conductivity (0-101.6cm)	Not specified
Sodium adsorption ratio (0-101.6cm)	Not specified
Soil reaction (1:1 water) (25.4-101.6cm)	Not specified
Subsurface fragment volume <=3" (0-152.4cm)	10–46%
Subsurface fragment volume >3" (0-152.4cm)	0–54%

Ecological dynamics

Climate

Located in the subalpine life zone, this ecological site is exposed to a variety of harsh environmental conditions. Compared to the boreal life zone, snowpack tends to be deeper and persist for longer durations of time. As a result, subalpine vegetation has a comparatively shorter season to grow and reproduce. When this ecological site is snow-free, cold soil temperatures and high winds also inhibit plant growth and performance with krummholz white spruce (*Picea glauca*) being common. This harsh climate maintains vegetation within this ecological site and prevents the establishment and/or growth of dominant boreal species like white spruce and black spruce (*Picea mariana*).

The cover and density of white spruce associated with this ecological site is highly variable but is largely controlled by the harsh subalpine climate. This ecological site occurs at elevations just above the boreal life zone. Given the nearby seed source and high

winds, this ecological site receives ample seed rain from forested stands of white spruce. As a result, ocular cover of white spruce is highly variable, commonly ranging from 0 to 15 percent canopy cover or more. While trees may be present, their ability to produce viable seed is limited by the harsh high-elevation climate (Roland et al. 2013). The subalpine climate combined with a fire regime greatly limits forest potential for this ecological site.

Fire


Within this area, fire is considered a natural and common event that typically is unmanaged. Fire suppression is limited and generally occurs adjacent to various villages spread throughout the area or on allotments with known structures, all of which have a relatively limited acre footprint. Most fires are caused by lightning strikes. Fire-related disturbances are highly patchy and can leave undisturbed areas within the burn perimeter.

The fire regime within interior Alaska follows two general scenarios—low-severity burns, and high-severity burns. It should be noted, however, that the fire regime in interior Alaska can be much more complex (Johnstone et al. 2008). Burn severity refers to the proportion of the vegetative canopy and organic material consumed in a fire event (Chapin et al. 2006). Fires in cool and moist habitat tend to result in low-severity burns, while fires in warm and dry habitat tend to result in high-severity burns. Because the soils have a thin organic cap and are well to somewhat excessively drained, the typical fire scenario for this ecological site is considered to result in a high severity burn. Large portions of the organic mat are consumed during a high-severity fire event, commonly exposing pockets of mineral soil. The loss of this organic mat, which insulates the mineral soil, and the decrease in site albedo tends to cause overall soil temperatures to increase (Hinzman et al. 2006). These alterations to soil temperature may result in increased depths of seasonal frost in the soil profile. High-severity fire events also destroy a majority of the vascular and nonvascular biomass above ground.


Successional dynamics suggest that the scrub dominant community 1.1 burns and that fire events will cause a transition to the pioneering stage of fire succession. This stage (community 1.2) is a mix of species that either regenerate in place (e.g., subterranean root crowns for willow and rhizomes for graminoids) and/or from wind-dispersed seed or spores that colonize exposed mineral soil (e.g., quaking aspen [*Populus tremuloides*] and *Ceratodon* moss [*Ceratodon purpureus*]). The pioneering stage of fire succession is primarily composed of forbs, grasses, and weedy bryophytes. This stage of succession currently persists for an unknown amount of time but is thought to last 10 to 30 years post-fire. Scrub birch (*Betula* spp.) and ericaceous shrubs continue to colonize and grow in stature on recently burned sites until they become dominant in the overstory, which marks the transition to the reference plant community (community 1.1).

State and transition model

Ecosystem states



State 1 submodel, plant communities



1.1a - Fire

1.2a - Time without fire

State 1 Reference state

The reference plant community is closed low scrub (Viereck et al. 1992) with the dominant shrub being scrub birch (*Betula glandulosa*). There are two plant communities within the reference state related to fire.

Dominant plant species

- resin birch (*Betula glandulosa*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous
- splendid feather moss (*Hylocomium splendens*), other herbaceous

Community 1.1

Scrub birch - bog blueberry / Schreber's big red stem moss - splendid feathermoss

The reference plant community is characterized as closed low scrub (Viereck et al. 1992) with scrub birch (*Betula glandulosa*) the dominant overstory vegetation. White spruce (*Picea glauca*) is common but typically have limited cover. Other common species include bog blueberry (*Vaccinium uliginosum*), bog Labrador tea (*Ledum groenlandicum*), marsh Labrador tea (*Ledum palustre* ssp. *decumbens*), crowberry (*Empetrum nigrum*), lingonberry (*Vaccinium vitis-idaea*), various reindeer lichen (*Cladina* spp.), Schreber's big redstem moss (*Pleurozium schreberi*), and splendid feathermoss (*Hylocomium*

splendens). The soil surface is primarily covered with herbaceous litter, lichen, and moss. The vegetative strata that characterize this community are medium shrubs

Dominant plant species

- white spruce (*Picea glauca*), tree
- resin birch (*Betula glandulosa*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- bog Labrador tea (*Ledum groenlandicum*), shrub
- marsh Labrador tea (*Ledum palustre* ssp. *decumbens*), shrub
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- greygreen reindeer lichen (*Cladina rangiferina*), other herbaceous
- reindeer lichen (*Cladina stygia*), other herbaceous

Community 1.2

Bluejoint / fireweed

Community 1.2 is in the pioneering stage of fire-induced secondary succession for this ecological site. It is characterized as either a mesic forb herbaceous or mesic graminoid herbaceous community (Viereck et al. 1992). White spruce (*Picea glauca*) occasionally occurs but have limited cover. Although small areas of exposed bare soil are common, the soil surface is primarily covered with a mixture of weedy bryophyte species, woody debris, and herbaceous litter. Commonly observed species include various willow (*Salix* spp.), fireweed (*Chamerion angustifolium*), bluejoint (*Calamagrostis canadensis*), juniper polytrichum moss (*Polytrichum juniperium*), and common liverwort (*Marchantia polymorpha*). The vegetative strata that characterize this community are medium forbs (between 4 and 24 inches), medium graminoids (between 4 and 24 inches), and mosses.

Dominant plant species

- willow (*Salix*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- fireweed (*Chamerion angustifolium*), other herbaceous
- juniper polytrichum moss (*Polytrichum juniperinum*), other herbaceous
- (*Marchantia polymorpha*), other herbaceous

Pathway 1.1a

Community 1.1 to 1.2

A fire sweeps through and incinerates much of the above ground vegetation. Because of the associated dry soils, this ecological site commonly experiences high-severity fires. A significant proportion of organic matter is consumed, leaving exposed mineral soil. Vegetation usually resprouts from surviving individuals or is recruited from nearby areas via seed or seedbank

Pathway 1.2a

Community 1.2 to 1.1

Time without fire. Grass and forb cover decreases, while scrub birch (*Betula* spp.) and ericaceous shrub cover increases.

Additional community tables

Inventory data references

The vegetation modeled for this ecological site has limited data and is considered provisional. The associated model was largely developed from NRCS (Natural Resources Conservation Service) staff with working knowledge of the area and literature review.

Plant community composition is largely based on ecological sites from Major Land Resource Area (MLRA) 231X: Interior Alaska Highlands.

References

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286..

Other references

Chapin, F.S., L.A. Viereck, P.C. Adams, K.V. Cleve, C.L. Fastie, R.A. Ott, D. Mann, and J.F. Johnstone. 2006. Successional processes in the Alaskan boreal forest. Page 100 in *Alaska's changing boreal forest*. Oxford University Press.

Hinzman, L.D., L.A. Viereck, P.C. Adams, V.E. Romanovsky, and K. Yoshikawa. 2006. Climate and permafrost dynamics of the Alaskan boreal forest. *Alaska's changing boreal forest* 39–61.

Johnstone, J.F. 2008. A key for predicting postfire successional trajectories in black spruce stands of interior Alaska. US Department of Agriculture, Forest Service, Pacific Northwest Research Station.

LANDFIRE. 2009. Western North American Boreal Treeline White Spruce Woodland - Boreal. In: *LANDFIRE National Vegetation Dynamics Models*. USDA Forest Service and US Department of Interior. Washington, DC.

Roland, C.A., J.H. Schmidt, and J.F. Johnstone. 2014. Climate sensitivity of reproduction in a mast-seeding boreal conifer across its distributional range from lowland to treeline

forests. *Oecologia* 174:665–677.

Schoeneberger, P.J. and D.A. Wysocki. 2012. Geomorphic Description System. Natural Resources Conservation Service, 4.2 edition. National Soil Survey Center, Lincoln, NE.

United States Department of Agriculture, . 2022. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.

Viereck, L.A., C. T. Dyrness, A. R. Batten, and K. J. Wenzlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286.

Contributors

Tyler Annetts
Phil Barber

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	03/18/2026
Approved by	Blaine Spellman
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

-
3. **Number and height of erosional pedestals or terracettes:**

 4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

 5. **Number of gullies and erosion associated with gullies:**

 6. **Extent of wind scoured, blowouts and/or depositional areas:**

 7. **Amount of litter movement (describe size and distance expected to travel):**

 8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

 9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

 10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

 11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

 12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-