

Ecological site R242XY405AK

Arctic scrub loamy floodplain

Last updated: 5/29/2025
Accessed: 12/22/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 242X–Northern Seward Peninsula-Selawik Lowlands

The Northern Seward Peninsula-Selawik Lowlands (MLRA 242X) includes the mosaic of coastal lowlands, river deltas, gently sloping uplands, and isolated hills and low mountains along the northern Seward Peninsula and in the lower Selawik Basin at the head of Kotzebue Sound. To the east, the area extends to the lower slopes of the Purcell Mountains, Zane Hills, and Sheklukshuk Range. MLRA 242X makes up 8,445 square miles. Lakes, ponds, and saturated soils occur throughout most of this area. The MLRA is mostly undeveloped wildland and is sparsely populated. It is in the zone of continuous permafrost.

Land ownership:

MLRA 242X encompasses the northernmost portion of the Bering Land Bridge National Preserve, the westernmost portion of Cape Krusenstern National Monument, and the majority of the Selawik National Wildlife Refuge.

The Bering Land Bridge National Preserve is located on the Seward Peninsula. The preserve is one of the most remote protected areas in the United States. The preserve is used by native Alaskans for subsistence hunting and is home to several archaeological sites. The preserve is around 2.6 million acres.

The Cape Krusenstern National Monument is located in northwestern Alaska, bordering the Chukchi Sea. Beach ridges within the monument safeguard evidence of 5,000 years of occupation by the Inupiat people, and more than 9,000 years of human occupation (NPS 2025). The Inupiat people maintain subsistence camps within the monument to this day. The monument is around 650,000 acres in size.

The Selawik National Wildlife Refuge is located in the Waring Mountains of northwestern Alaska, and comprises a transitional zone marking the opening of boreal forests into Arctic tundra (USFWS 2025). Selawik Refuge is an important migratory and wintering habitat for the Western Arctic Caribou Herd, one of four herds found across Arctic Alaska. The refuge is around 2.1 million acres, with 240,000 of that set aside as federally designated wilderness.

The majority of MLRA 242X is private lands or those managed by the USFWS, NPS, BLM, and State of Alaska. The USFWS manages 1,604,744 acres, around 36 percent of the MLRA. The National Park Service manages 1,350,544 acres, around 30 percent of the MLRA. The BLM manages 606,464 acres, around 14 percent of the MLRA. The State of Alaska has a patent on 55,675 acres of the MLRA, around 1 percent of the MLRA. Much of the MLRA is within Alaska Native allotments or patents around 832,328 acres, or 19 percent of the MLRA.

Climate:

The arctic climate of MLRA 242X is characterized by brief, cool summers and long, very cold winters. The average annual precipitation ranges from 11 to 12 inches. Average annual snowfall ranges from about 40 to 100 inches. The average annual temperature ranges from 18 to 25 degrees Fahrenheit. The average freeze-free period is between 98 to 104 days.

Geology:

The western part of this MLRA was unglaciated during the Pleistocene Epoch. Most of the eastern part was covered by glacial ice originating in the Waring Mountains and Brooks Range to the north. Sediments across the vast majority of the area consist of fine textured, Holocene and Pleistocene deltaic and fluvial deposits on coastal lowlands, Holocene fluvial deposits on flood plains and stream terraces, and mixed colluvium and slope alluvium on mountain footslopes. The underlying bedrock geology consists primarily of stratified sedimentary rocks and volcanic rocks of Cretaceous, Tertiary, and Quaternary age.

Soils:

The dominant soil orders in this MLRA are Gelisols, with Histosols and Entisols covering a comparatively minor extent in coastal or estuarine zones. These Gelisols are shallow or moderately deep to permafrost and are typically poorly to very poorly drained. Miscellaneous (nonsoil) areas make up about 25 percent of this MLRA. The most common are water and beaches.

Gelisols are soils that have permafrost within 100 cm of the soil surface and/or have gelic materials within 100 cm of the soil surface and have permafrost within 200 cm. Gelic materials are mineral or organic soil materials that have evidence of cryoturbation (frost churning) and/or ice segregation in the active layer (seasonal thaw layer) and/or the upper

part of the permafrost (NRCS 2024). The common suborders of Gelisols within this MLRA are Turbels, Histels, and Orthels.

The Histels have thick accumulations of surface organic material and are associated with high-center polygons. The Orthels and Turbels have comparably thinner surface organic material and occur on high floodplains, stream terraces, low-center polygons, and the slopes of hills and plains. Turbels show signs of cryoturbation while Orthels do not.

Histosols have a high content of organic matter and no permafrost. Most are saturated year-round, but a few are freely drained. The most common suborder of Histosols in this MLRA are Fibrists. Fibrists are wet, slightly decomposed Histosols. Most of the soils support natural vegetation of widely spaced small trees, shrubs, forbs, and grasses and grass-like plants (NRCS 2025).

Entisols are soils that show little or no evidence of pedogenic horizon development and in this MLRA are associated with recently deposited sediments. The most common suborder of Entisols in this MLRA are Aquents, which are widely distributed throughout this MLRA. They are common in low lying estuarine areas bordering the Chukchi Sea and Kotzebue Sound.

Fire Dynamics and Succession:

Reported fire history in this MLRA spans 1957 to 2021. During this time period, there were 67 recorded fires. The mean fire size was approximately twelve thousand acres. The largest fire was the 1977 AUGUS fire, burning 235,584 acres (AICC 2025). The number of fires that burn in tundra systems are limited by ignitions, prevalence of atmospheric moisture, and fuel moisture content or fuel availability. Fires in the tundra happen less often than in boreal systems (NPS 2024). Fires can be sporadic and widely distributed. The fire return intervals vary widely in tundra systems, from 30 years to over 1,000 years. In the past, fires have burned more frequently on the Seward Peninsula and in the Noatak Valley than in other tundra regions. This is due to slightly warmer and drier conditions and higher amounts of plants and shrubs above ground such as tussock cottongrass (NPS 2024).

There are over 143 million acres of arctic and subarctic tundra in Alaska, most of which is designated in a limited fire management option. A limited fire management option is one of four fire management options outlined in the Alaska Interagency Wildland Fire Management Plan (AICC 2024). A 'limited' approach is the most hands-off management option, meaning the fire will be left to behave naturally and fill its natural ecological role (AICC 2024).

Arctic tundra areas are experiencing warmer temperatures and an increase in fire activity over the past twenty years. Climate model forecasts show more warming in the future, particularly in the high northern areas. This could affect the length of the growing season, how well plants and shrubs grow, and rain and snowfall. A longer and more robust growing season could likely impact trends regarding number and intensity of fires in the arctic tundra.

Vegetation dynamics:

Uplands are not common in this MLRA but where present they generally support dwarf scrub dominated by tussock tundra. On shallow, rocky soils and exposed sites, lichens and scattered herbs dominate the ground layer. Bare soil and bedrock generally are extensive. On mesic sites, halophytic sedges and grasses dominate. Depressions, drainageways, and other saturated areas support wet sedge meadows and wet sedge-moss meadows. The vegetation on flood plains consists of a mixture of wet sedge meadows and of tall scrub dominated by various willows and shrub birch.

Vascular and non-vascular plant succession dynamics vary in post-fire environments. Data collected in the footprint of the abnormally severe Anaktuvuk River fire shows that four years after the burn, above-ground net primary productivity of vascular plants was equal in burned and unburned areas, though total live biomass was less (Bret-Harte et al 2013). Graminoid biomass had recovered to unburned levels, but shrubs had not. Most of the vascular plant biomass had resprouted from surviving underground parts. Much greater changes were observed in the biomass and composition of the non-vascular plant community (Bret-Harte et al 2013). Lichen biomass appears to take decades to centuries to recover from the disturbances of fires, particularly under a warming climate (Racine et al 2004).

Tussock cottongrass is the primary tussock forming plant species in arctic tundra. Tussock cottongrass burns easily yet is very hardy. It grows quickly and is very productive, forming a tussock made of fine dead fuels from the previous year. These fine fuels burn quickly and easily. Tundra fires are usually wind driven and move rapidly, burning through years of accumulated tussock grass thatch. The tussock growth protects the cottongrass roots and plant tissues from fire. These plants and shrubs recover and grow vigorously after fire, benefitting from nutrients and warmer soils after a fire (NPS 2024).

Fire impacts soil composition and structure (Li et al 2021). Fires will often partially or completely consume the organic material layer, depending on the severity of the fire. Fire can also impact permafrost. On average, forest fires reduce the permafrost extent by up to 9-16 percent and accelerate permafrost thaw by five years. The effects of wildfire on permafrost are much larger in forested areas than in tundra, bogs, and fens (Li et al 2021).

Classification relationships

Landfire Biophysical Settings –

6816980 – Alaska Arctic Wet Sedge Meadow (Landfire 2009)

Viereck Communities:

Wet sedge meadow tundra – III.A.3.a (Viereck et al. 1992)

Ecological site concept

- Arctic climate
- Associated landforms are low flood plains
- Soils are derived from organic material over alluvium.
- Soils are considered somewhat poorly drained
- Soils have permafrost at 31 inches
- The reference plant community is characterized as a wet sedge meadow tundra (Vioreck et al 1992) with the dominant plants being willows, sedges, and tussock cottongrass.

Associated sites

R242XY406AK	Arctic scrub loamy frozen floodplain Ecological sites 405 and 406 are both found on floodplains. Ecological site 406 is found on terraces, while site 405 is found in troughs. This leads to differences in soil moisture regime, flooding frequency, and depth to restriction which all impact the type and distribution of vegetation found at the two sites.
-------------	--

Similar sites

R242XY401AK	Arctic silty polygon complex Ecological site 401 is found on low center polygons, while ecological site 405 is found on low floodplains. Although they are located on different parts of the landscape, the vegetation associated with both can be described as a wet sedge meadow tundra.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Salix fuscescens</i> (2) <i>Salix pulchra</i>
Herbaceous	(1) <i>Carex aquatilis</i> (2) <i>Eriophorum angustifolium</i>

Physiographic features

The Arctic scrub loamy floodplain ecological site occurs on low flood plains. Elevation ranges between 10 and 450 feet. Associated foothills are nearly level. This site occurs on all aspects and shows no preference for north-facing or south-facing slopes. Flooding occurs occasionally for brief durations. Soils are considered somewhat poorly drained. The flooding duration is brief, and there is no ponding at this site. The water table associated with this site is 10 to 20 inches below the soil surface.

Table 2. Representative physiographic features

Landforms	(1) Alluvial plain > Flood plain
-----------	----------------------------------

Runoff class	Medium
Flooding duration	Brief (2 to 7 days)
Flooding frequency	Occasional
Ponding frequency	None
Elevation	3–137 m
Slope	0–2%
Water table depth	25–51 cm
Aspect	W, NW, N, NE, E, SE, S, SW

Climatic features

Sea ice strongly influences the climate of MLRA 242X. A protective barrier of sea ice has typically formed along the Chukchi coast and Kotzebue sound by early November and often lasting until April. The Kotzebue sound and Chukchi sea moderate diurnal and monthly temperatures resulting in a maritime climate. Summer temperatures (June through August) are relatively stable with mean maximum monthly temperatures ranging between 50 to 58 degrees Fahrenheit. As sea ice forms on the coastline, temperatures decrease significantly with the area shifting to a continental climate. The coldest months (January through March) have mean monthly temperatures ranging from 4 to 7 degrees Fahrenheit. The coverage and formation of sea ice has been decreasing due to rising atmospheric temperatures (USDA Climate Hub 2025).

The Northern Seward Peninsula-Selawik Lowlands have summers that are short and cool and winters that are long and cold. Strong winds are common throughout the year. Mean annual air temperatures typically range from 18 to 25 degrees Fahrenheit. The warmest months are June, July, and August. During these summer months, the typical freeze free period for the area ranges from 103 to 99 days. The coldest months are January, February, and March.

This area is semi-arid with mean annual precipitation typically around 12 inches. The warmest months have overcast skies with frequent fog and precipitation while the coldest months have clear skies. The three wettest months are July, August, and September where the area may receive half of its total annual precipitation.

Table 3. Representative climatic features

Frost-free period (characteristic range)	59-75 days
Freeze-free period (characteristic range)	99-103 days
Precipitation total (characteristic range)	279-305 mm
Frost-free period (actual range)	54-80 days

Freeze-free period (actual range)	98-104 days
Precipitation total (actual range)	279-305 mm
Frost-free period (average)	67 days
Freeze-free period (average)	101 days
Precipitation total (average)	305 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) WALES [USW00026618], Wales, AK
- (2) KOTZEBUE RALPH WEIN AP [USW00026616], Kotzebue, AK

Influencing water features

This ecological site is considered a riverine wetland. Dominant water sources are often overbank flow from the channel or subsurface hydraulic connections between the stream channel and wetlands.

Wetland description

This ecological site is classified as a riverine wetland under the Hydrogeomorphic (HGM) classification system (Smith et al. 1995; USDA-NRCS 2008).

Soil features

- Soils formed from organic material and alluvium
- Subsurface rock fragments both bigger and smaller than three inches are present at this site, although they are not common
- The surface mineral horizon is a silt loam formed from sand and alluvium
- Soils are very deep
- Permafrost is a restriction that occurs at moderate depths (35 inches)
- These are considered wet soils that are somewhat poorly
- Soils range from very strongly acidic to moderately alkaline

Table 4. Representative soil features

Parent material	(1) Organic material (2) Alluvium
-----------------	--------------------------------------

Surface texture	(1) Silt loam
Family particle size	(1) Coarse-loamy
Drainage class	Somewhat poorly drained
Permeability class	Moderately rapid
Depth to restrictive layer	30–89 cm
Soil depth	152 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	10.16–16 cm
Calcium carbonate equivalent (0-25.4cm)	0%
Clay content (25.4-101.6cm)	0–5%
Electrical conductivity (25.4-101.6cm)	0–2 mmhos/cm
Subsurface fragment volume <=3" (0-152.4cm)	0–4%
Subsurface fragment volume >3" (0-152.4cm)	0–4%

Ecological dynamics

The Northern Seward Peninsula-Selawik Lowlands MLRA has a harsh climate and cold soils and occurs in the zone of continuous permafrost. This MLRA occurs in the arctic biome and has a growing season that is both short and cold. As a result, the vertical and horizontal structure of vegetation is severely limited. Vegetation within the arctic biome is typically restricted to dwarf shrubs, mosses, and lichens.

Vegetation over most of this MLRA consists of mesic graminoid herbaceous and dwarf scrub communities. Open low scrub occurs along drainages. Mesic graminoid herbaceous communities dominated by tussock-forming sedges are widespread. Typical species are tussock cottongrass (*Eriophorum vaginatum*) and Bigelow's sedge (*Carex bigelowii*). Low shrubs, such as dwarf arctic birch (*Betula nana*), crowberry (*Empetrium nigrum*), Labrador-tea (*Ledum palustre* ssp. *decumbens*), and mountain cranberry (*Viburnum edule*) often occur and may co-dominate with sedges. Mosses and lichens are commons between tussocks. Dwarf scrub communities are dominated by mat forming *Dryas* species accompanied by ericaceous species (Gallant et al 1995).

State and transition model

Ecosystem states

1. Reference state

State 1 submodel, plant communities

1.1. bog willow -
tealeaf willow / water
sedge - tall cottongrass
/ sphagnum

State 1 Reference state

The reference plant community is characterized as wet sedge meadow tundra (Viereck et al. 1992). This ecological site has no known associated disturbance regimes and has one plant community within the reference state. All plant communities associated with this ecological site have limited data, so the state-and-transition model is provisional.

Dominant plant species

- Alaska bog willow (*Salix fuscescens*), shrub
- tealeaf willow (*Salix pulchra*), shrub
- bog rosemary (*Andromeda polifolia*), shrub
- tall cottongrass (*Eriophorum angustifolium*), grass
- lesser saltmarsh sedge (*Carex glareosa*), grass
- round sedge (*Carex rotundata*), grass
- sphagnum (*Sphagnum*), other herbaceous

Community 1.1

bog willow - tealeaf willow / water sedge - tall cottongrass / sphagnum

This plant community is characterized as a wet sedge meadow (Viereck et al. 1992) with the dominant plants being bog willow (*Salix fuscescens*), tealeaf willow (*Salix pulchra*), bog rosemary (*Andromeda polifolia*), water sedge (*Carex aquatilis*), tall cottongrass (*Eriophorum angustifolium*), saltmarsh sedge, and Sphagnum. The vegetative strata with the highest cover are low shrubs, sedges, and moss (Landfire 2009).

Dominant plant species

- Alaska bog willow (*Salix fuscescens*), shrub

- tealeaf willow (*Salix pulchra*), shrub
- bog rosemary (*Andromeda polifolia*), shrub
- tall cottongrass (*Eriophorum angustifolium*), grass
- lesser saltmarsh sedge (*Carex glareosa*), grass
- round sedge (*Carex rotundata*), grass
- sphagnum (*Sphagnum*), other herbaceous
- drepanocladus moss (*Drepanocladus*), other herbaceous

Additional community tables

Animal community

Mammals common to the area include brown bear, caribou, moose, musk ox, black bear, wolf, red fox, a variety of other furbearers, and rodents. Many species of migratory waterfowl and shore birds' nest in the ponds and wetlands. Raptors include gyrfalcon, peregrine falcon, golden eagle, hawks, and owls. Arctic char and Arctic grayling are in most of the rivers. Lake trout and northern pike are common in many lakes.

Hydrological functions

n/a

Recreational uses

Local residents use this remote area primarily for subsistence hunting, fishing, and gathering. Hunting and other kinds of wildland recreation are increasingly important. Most visitors are served by air taxi, guiding, and outfitting companies operating out of the major Alaska communities. Most of the communities are along the major rivers or lakes or on the coast.

Wood products

n/a

Other products

n/a

Other information

The major soil resource concern is disturbance of the fragile permafrost-affected soils. Disturbance of the insulating organic material at the surface results in thawing of the upper soil layers. This thawing can lead to ponding, soil subsidence, erosion, and disruption of surface drainage. All management activities need to include protection of the organic surface material and the thermal balance of the soils.

Inventory data references

The vegetation modeled for this site has limited data and is considered provisional. The associated model was largely developed from NRCS staff with working knowledge of the area and literature review.

Other references

“Alaska Fire History Chart.” AICC - Predictive Services - Intelligence / Reports, Alaska Interagency Coordination Center, fire.ak.blm.gov/predsvcs/intel.php. Accessed 29 Jan. 2025.

Bret-Harte MS, Mack MC, Shaver GR, Huebner DC, Johnston M, Mojica CA, Pizano C, Reiskind JA. 2013 The response of Arctic vegetation and soils following an unusually severe tundra fire. *Phil Trans R Soc B* 368: 20120490.
<http://dx.doi.org/10.1098/rstb.2012.0490>

“Cape Krusenstern Animals.” National Parks Service, U.S. Department of the Interior, 27 Sept. 2027, www.nps.gov/cakr/learn/nature/animals.htm.

“Existing Vegetation Cover.” LandFire, USDA / DOI, www.landfire.gov/vegetation/evc. Accessed 29 Jan. 2025.

“FEIS - Syntheses about Fire Ecology and Fire Regimes in the United States.” Fire Effects Information System, www.feis-crs.org/feis/. Accessed 29 Jan. 2025.

“Fire in Ecosystems: Arctic Tundra (U.S. National Park Service).” National Parks Service, U.S. Department of the Interior, www.nps.gov/articles/000/fire-in-ecosystems-arctic-tundra.htm. Accessed 29 Jan. 2025.

Gallant, Alisa L., et al. “Ecoregions of Alaska.” Professional Paper, 1995, <https://doi.org/10.3133/pp1567>.

Higuera, Philip E., et al. “Variability of tundra fire regimes in Arctic Alaska: Millennial-scale patterns and ecological implications.” *Ecological Applications*, vol. 21, no. 8, 1 Dec. 2011, pp. 3211–3226, <https://doi.org/10.1890/11-0387.1>.

Hydrogeomorphic Wetland Classification System: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service, USDA - NRCS, Feb. 2008.

“History & Culture.” National Parks Service, U.S. Department of the Interior, 18 Nov. 2024, www.nps.gov/cakr/learn/historyculture/index.htm.

News from the Western Arctic Caribou Herd Working Group, Alaska Department of Fish

and Game, July 2014,
www.adfg.alaska.gov/static/home/library/pdfs/wildlife/caribou_trails/caribou_trails_2014.pdf. Accessed 6 Mar. 2025.

Li, Xiao-Ying, et al. "Influences of forest fires on the Permafrost Environment: A Review." *Advances in Climate Change Research*, vol. 12, no. 1, Feb. 2021, pp. 48–65, <https://doi.org/10.1016/j.accre.2021.01.001>.

Racine, C., Jandt, R., Meyers, C., & Dennis, J. (2004). Tundra Fire and Vegetation Change along a Hillslope on the Seward Peninsula, Alaska, U.S.A. *Arctic, Antarctic, and Alpine Research*, 36(1), 1–10. 036[0001:TFAVCA]2.0.CO;2

Rickbeil, Gregory J.M., et al. "The grazing impacts of four barren ground caribou herds (*Rangifer tarandus groenlandicus*) on their summer ranges: An application of archived remotely sensed vegetation productivity data." *Remote Sensing of Environment*, vol. 164, July 2015, pp. 314–323, <https://doi.org/10.1016/j.rse.2015.04.006>.

Selawik National Wildlife Refuge, US Fish and Wildlife Service, www.fws.gov/refuge/selawik. Accessed 5 Mar. 2025.

Smith, R. Daniel, et al. "An Approach for Assessing Wetland Functions Using Hydrogeomorphic Classification, Reference Wetlands, and Functional Indices." *Wetlands Research Program Technical Report WRP-DE-9 - US Army Corps of Engineers*.

Viereck, L.A., et al. "The Alaska Vegetation Classification." *General Technical Report PNW-GTR-286*, July 1992, <https://doi.org/10.2737/pnw-gtr-286>.

Contributors

Blaine T Spellman
Tyler Annetts
Matthew J Mayer

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
--------------------------	--

Contact for lead author	
Date	12/22/2025
Approved by	Blaine Spellman
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought**

or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. **Perennial plant reproductive capability:**
