

# Ecological site R243XY301AK

## Alpine scrub gravelly swales

Last updated: 5/29/2025

Accessed: 02/08/2026

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### MLRA notes

Major Land Resource Area (MLRA): 243X–Western Brooks Range Mountains, Foothills, and Valleys

Boundaries and important features:

The Western Brooks Range Mountains, Foothills, and Valleys MLRA (MLRA 243X) encompasses the southern slopes of the De Long Mountains, the Baird Mountains, the Noatak River drainage, and the lower Kobuk River drainage. The southern limit of the area includes the western Lockwood Hills, Sheklukshuk and Waring Mountains, and Kiana and Igichuk Hills. MLRA 243 makes up 22,705 square miles. This MLRA shares boundaries with MLRAs 242X, 244X, and 245X. MLRA 242X (Northern Seward Peninsula-Selawik Lowlands) has nearly level to rolling plains, river deltas, and extended mountain footslopes. MLRA 244X (Northern Brooks Range Mountains) has steep, rugged, high mountains and narrow valleys. MLRA 245X (Arctic Foothills) has broad, rounded hills and nearly level uplands at the base of the Brooks Range. MLRA 243X shares a less apparent boundary with MLRAs 233 (Upper Kobuk and Koyukuk Hills and Valleys) and 234X (Interior Brooks Range Mountains), where a continental subarctic climate prevails.

Kobuk Valley National Park is located in the south-central portion of MLRA 243X, along the Kobuk River. Kobuk Valley National Park has an area of around 1.75 million acres and was designated to preserve the 100 ft high Great Kobuk Sand Dunes and the surrounding area which includes the Onion Portage caribou migration route. The Noatak National Preserve is located in the north-central portion of the MLRA, along the Noatak River corridor. The Noatak River is the nation's largest unaltered river basin, and the preserve is around 6.5 million acres. 5.7 million acres of the preserve are designated as wilderness. The Noatak River is also a designated National Wild and Scenic River.

The Red Dog Mine is located in the northwestern portion of the MLRA. The Red Dog Mine is the world's largest producer of zinc and has the world's largest zinc reserves. The mine is the primary economic driver of the Northwest Arctic Borough. The Northwest Arctic Borough was formed in 1986 on the economic basis of taxing the mine.

#### Geology:

The entire area was glaciated during the Early and Middle Pleistocene, except for possibly small portions of the Baird Mountains. By the Late Pleistocene, glaciers had retreated from most of the area, except for the central, upper-elevation portions of the De Long Mountains in the northern part of the MLRA. The valley of the upper Noatak River was likely covered by extensive proglacial lakes during parts of the Pleistocene Epoch. In the mountains, glacial deposits have eroded away or been buried by mountain colluvium and alluvium, which accumulated during the Holocene Epoch across about 60 percent of the present landscape.

Glacial moraines, drift, and outwash deposits are extensive on the lower mountain slopes and in valleys at the mid and lower elevations. These deposits cover about 18 percent of the MLRA. Flood plains, stream terraces, and alluvial fans have recent and Pleistocene fluvial deposits. The underlying bedrock geology consists almost entirely of stratified sedimentary rocks of Paleozoic and Precambrian age and, in some cases, Cretaceous age.

This area is in the zone of continuous permafrost. In the mountains, permafrost is most evident in unconsolidated materials. In the valleys, thick layers of permafrost occur in both fine textured and coarse textured materials. Depth to the base of the permafrost layer may be 1,000 feet (305 meters) or more. In close proximity to water bodies, it may be 600 feet (185 meters) or more. Periglacial features, such as pingos, thermokarst pits, thaw lakes, solifluction lobes, and high- and low-center polygons, are common on-stream terraces, on the lower mountain slopes, and in swales on foothills.

#### Soils:

The dominant soil orders in this MLRA are Gelisols, Entisols, Inceptisols, and Mollisols. The soils in the area have a gelic soil temperature regime and an aquic or udic soil moisture regime. The Gelisols are shallow or moderately deep to permafrost, occur on fine to gravelly textured sediments, and are very poorly to moderately well drained. Common Gelisol suborders are Histels, Orthels, and Turbels. The Histels have thick accumulations of surface organic material and occur in depressions, lake margins, and shallow basins. The Orthels and Turbels have comparably thinner surface organic material with the Turbels being cryoturbated. These widespread soils occur on slopes of mountains, hills, and plains across the MLRA. Inceptisols (suborder Gelepts), Mollisols (suborder Gelolls), and Entisols (suborder Gelents) occur on upper mountain slopes and ridges formed in loamy to stony colluvium and residuum. These soils are shallow to deep and are generally well drained. Entisols (suborder Cryofluvents) on flood plains formed in stratified loamy,

sandy, and gravelly alluvium and are somewhat poorly to well drained.

### Vegetation Dynamics:

The mountain slopes and ridges in this area generally support dwarf scrub dominated by Dryas, ericaceous shrubs including crowberry, and dwarf willow. Lichens and scattered herbs dominate the ground layer on shallow, rocky soils and exposed sites. There are extensive areas of bare soil and bedrock. On the more mesic sites, sedges, forbs, and mosses cover most of the surface. Areas at the lower elevations and areas of the deeper soils on nearly level uplands, terraces, and basins generally support low willow and ericaceous shrub scrub and mesic graminoid herbaceous communities, commonly with extensive areas of tussock-forming sedges. Saturated sites support wet sedge meadows and wet sedge-moss meadows. Flood plains support a mixture of tall and low scrub dominated by various willows, shrub birch, and alder.

### Climate:

Short, generally cool summers and long, very cold winters characterize the continental arctic climate of the area. The average annual precipitation ranges from about 10 to 15 inches (255 to 380 millimeters) at the lower elevations in the western part of this MLRA and along the central Noatak River and from about 20 to 40 inches (510 to 1,015 millimeters) in the mountains. The average annual snowfall is about 35 to 100 inches (90 to 255 centimeters). The average annual temperature ranges from about 8 to 16 degrees F (-13 to -9 degrees C). Snow and freezing temperatures can occur in any month of the year, particularly at the higher elevations.

### LRU notes

This area supports two life zones defined by the physiological limits of plant communities along an elevational gradient: arctic lowlands and alpine. In this MLRA, the arctic lowland life zone typically occurs below 1000 feet elevation on average and is the elevational band where lowland vegetation dominates. For this MLRA, certain vascular plant species are common in the lowlands and much less common in the alpine (i.e. *Betula nana*, *Salix pulchra*, *Ledum palustre*). Above the arctic lowlands band of elevation, alpine vegetation dominates. For this MLRA, certain vascular plant and lichen species are common in the alpine and much less common in the lowlands (i.e. *Dryas octopetala*, *Saxifraga oppositifolia*, *Empetrum nigrum*). The transition between arctic and alpine vegetation can occur within a range of elevations, and is highly dependent on latitude, slope, aspect, and shading from adjacent mountains.

### Classification relationships

Landfire Biophysical Settings – 16390 - Arctic Mesic-Wet Willow Shrubland (Landfire 2009)

## Viereck Communities:

Open Low Willow-Sedge Shrub Tundra— II.C.2.h (Viereck et al. 1994)

### Ecological site concept

- Associated with swales on the flanks and base of rugged mountains.
- Associated with the alpine. Elevation ranges between 1000 and 5400 feet. Slopes are gently sloping to steep and occur on all aspects.
- Soils are derived from silty loess over gravelly colluvium.
- Soils are deep to very deep, with soil depth controlled by bedrock contact. Soils typically do not have permafrost. Where present, permafrost occurs at moderate depth (20 to 30 inches).
- These moist soils are considered somewhat poorly drained and neither pond or flood.
- The reference plant community is characterized as open low willow-sedge shrub tundra (Viereck et al., 1992) with dominant shrubs being tealeaf willow, feltleaf willow, and grayleaf willow. This ecological site has no known associated disturbance regimes and has one plant community within the reference state.

### Associated sites

R243XY404AK	<b>Alpine dwarf scrub gravelly limestone slopes</b> Ecological site 404 occurs upslope on dry soils derived from limestone and similar bedrock.
R243XY403AK	<b>Arctic dwarf scrub gravelly slopes</b> Ecological site 403 occurs upslope on dry soils derived from acidic bedrock.
R243XY302AK	<b>Alpine scrub gravelly frozen slopes</b> Occurs on the same mountain slopes but on wet and frozen soils.

### Similar sites

R243XY302AK	<b>Alpine scrub gravelly frozen slopes</b> Ecological site 302 occupies a similar ecological niche as 301 but does not occur in swales and has soils derived from limestone and similar bedrock. Differences in drainage, presence and depth of permafrost, and soil chemistry result in different kinds and amounts of vegetation.
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Salix pulchra</i> (2) <i>Salix alaxensis</i>
Herbaceous	(1) <i>Carex aquatilis</i> (2) <i>Pleurozium schreberi</i>

## Physiographic features

This alpine scrub ecological site (R243XY301AK) is associated with swales on the flanks and bases of rugged mountains. Swales are defined as shallow, open depressions that lack defined channels but can funnel overland or subsurface surface flow into a drainageway (Schoeneberger and Wysocki 2017). Representative elevation ranges between 1000 and 5400 feet but this ecological site occurs at lower elevations on exposed, windswept positions. Slopes range from gently sloping to steep and show no preference for north-facing or south-facing aspects. Flooding and ponding do not occur. A seasonal water table occurs in May and June at shallow depths (10 to 20 inches). There is low amounts of runoff to adjacent, downslope ecological sites.

**Table 2. Representative physiographic features**

Geomorphic position, mountains	(1) Mountainflank (2) Mountainbase
Hillslope profile	(1) Backslope (2) Foothslope
Landforms	(1) Mountains > Mountain slope > Swale
Runoff class	Low
Flooding frequency	None
Ponding frequency	None
Elevation	305–1,646 m
Slope	10–20%
Water table depth	25–51 cm
Aspect	W, NW, N, NE, E, SE, S, SW

**Table 3. Representative physiographic features (actual ranges)**

Runoff class	Not specified
Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	15–1,929 m
Slope	8–40%
Water table depth	0–51 cm

## Climatic features


MLRA 243X is characterized by an arctic continental climate having short, warm summers and long, cold, dark winters. The average annual mean temperature is 20.1 degrees

Fahrenheit. The warmest months span June through August, with normal maximum temperature ranging from 57 to 61 degrees Fahrenheit. The coldest months are December through February, with normal minimum temperature ranging from -15 to -10 degrees Fahrenheit. This MLRA receives high annual precipitation with the summer months being the wettest. Average annual precipitation ranges from 9 to 32 inches. Approximately 40 percent of the annual precipitation occurs during the months of July through September.


Snow persists in the alpine and arctic lowland life zones throughout much of the year. A lack of trees and tall shrubs also means that this site is subject to strong winds, further exacerbating cold temperatures. This site has a much shorter growing season than sites in lower elevation areas and the growing season is significantly colder for associated vegetation.

**Table 4. Representative climatic features**


Frost-free period (characteristic range)	55-85 days
Freeze-free period (characteristic range)	40-70 days
Precipitation total (characteristic range)	356-660 mm
Frost-free period (actual range)	45-90 days
Freeze-free period (actual range)	30-75 days
Precipitation total (actual range)	229-813 mm
Frost-free period (average)	69 days
Freeze-free period (average)	54 days
Precipitation total (average)	508 mm




**Figure 1. Monthly precipitation range**




**Figure 2. Monthly minimum temperature range**




**Figure 3. Monthly maximum temperature range**



**Figure 4. Monthly average minimum and maximum temperature**



**Figure 5. Annual precipitation pattern**



**Figure 6. Annual average temperature pattern**

## Influencing water features

Due to its landscape position, this site is neither associated with or influenced by streams or wetlands. Precipitation and throughflow are the main source of water for this ecological site. Surface runoff and throughflow contribute some water to downslope ecological sites.

## Wetland description

Not a wetland.

## Soil features

- Soils formed in loess over gravelly colluvium. The colluvium is derived from schist and igneous rock.
- Rock fragments on the soil surface range between zero and five percent cover.
- Capped with one to eight inches of peat.
- The surface mineral horizons are silt loams or channery silt loams with the silty material being derived from loess or silty colluvium. Where present, the silt loam surface layer is

thin.

- These gravelly soils have subsurface rock fragments ranging between 15 and 65 percent of the soil profile by volume.
- Soils are deep to very deep, with soil depth controlled by bedrock contact.
- Soil restrictions include strong contrasting textural stratification (between 3 and 12 inches) and bedrock (between 48 and 70 inches). On rare occasion, permafrost occurs at moderate depth (20 to 30 inches).
- Soils range from strongly acidic to slightly acidic.
- These moist soils are typically considered somewhat poorly drained. Soils with permafrost are poorly drained.

**Table 5. Representative soil features**

Parent material	(1) Eolian deposits (2) Colluvium–schist (3) Colluvium–igneous rock
Surface texture	(1) Very channery loam (2) Peat
Family particle size	(1) Loamy-skeletal
Drainage class	Somewhat poorly drained
Permeability class	Moderately rapid
Depth to restrictive layer	8–30 cm
Soil depth	122–178 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0–5%
Available water capacity (0-101.6cm)	5.84–12.95 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Soil reaction (1:1 water) (25.4-101.6cm)	5.3–6.5
Subsurface fragment volume <=3" (0-152.4cm)	10–20%
Subsurface fragment volume >3" (0-152.4cm)	5–45%

**Table 6. Representative soil features (actual values)**

Drainage class	Poorly drained to somewhat poorly drained
----------------	---

Permeability class	Not specified
Depth to restrictive layer	Not specified
Soil depth	Not specified
Surface fragment cover <=3"	Not specified
Surface fragment cover >3"	Not specified
Available water capacity (0-101.6cm)	Not specified
Calcium carbonate equivalent (0-101.6cm)	Not specified
Electrical conductivity (0-101.6cm)	Not specified
Soil reaction (1:1 water) (25.4-101.6cm)	Not specified
Subsurface fragment volume <=3" (0-152.4cm)	Not specified
Subsurface fragment volume >3" (0-152.4cm)	Not specified

## Ecological dynamics

The Western Brooks Range MLRA (MLRA243X) occurs in the arctic biome and this site is exposed to a variety of harsh environmental conditions. This area has cool, short summers and long cold winters. Limited warmth during the short summer months inhibits trees from occurring, and the expansive tundra is composed of a mosaic of low growing shrubs, sedges, moss, and lichen. Sites with greater than 25 percent lichen cover tend to be exposed to the wind and accumulate little winter snow (Viereck et al. 1992).

The successional status of this ecological site is unclear but appears to be stable in most situations. Drier site conditions may lead to the development of tussock tundra while wetter site conditions may lead to the loss of shrubs and development of a wet sedge meadow community (Landfire 2009).

## State and transition model

### Ecosystem states

1. Reference State

## State 1 submodel, plant communities

- 1.1. tealeaf willow -
- feltleaf willow / water
- sedge / splendid
- feathermoss -
- Schreber's big red
- stem moss

## State 1 Reference State

The reference plant community is characterized as open low willow-sedge shrub tundra (Viereck et al. 1992). This ecological site has no known disturbance regimes and has one plant community within the reference state. All plant communities associated with this ecological site have limited data, so the state-and-transition model is provisional.

### Dominant plant species

- tealeaf willow (*Salix pulchra*), shrub
- feltleaf willow (*Salix alaxensis*), shrub
- water sedge (*Carex aquatilis*), grass
- splendid feather moss (*Hylocomium splendens*), other herbaceous
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous

### Community 1.1

#### **tealeaf willow - feltleaf willow / water sedge / splendid feathermoss - Schreber's big red stem moss**

This plant community is characterized as open low willow-sedge shrub tundra (Viereck et al. 1992) with the dominant plants being tealeaf willow, feltleaf willow, grayleaf willow, water sedge, and various feathermosses. Richardson's willow, water sedge, tall cottongrass, and Sphagnum. The vegetative strata with the highest cover are low shrubs and moss (Landfire 2009).

### Dominant plant species

- tealeaf willow (*Salix pulchra*), shrub
- feltleaf willow (*Salix alaxensis*), shrub
- grayleaf willow (*Salix glauca*), shrub
- mountain alder (*Alnus viridis* ssp. *crispa*), shrub
- dwarf birch (*Betula nana*), shrub
- bog blueberry (*Vaccinium uliginosum*), shrub
- black crowberry (*Empetrum nigrum*), shrub
- water sedge (*Carex aquatilis*), grass
- tall cottongrass (*Eriophorum angustifolium*), grass
- Bigelow's sedge (*Carex bigelowii*), grass

- splendid feather moss (*Hylocomium splendens*), other herbaceous
- Schreber's big red stem moss (*Pleurozium schreberi*), other herbaceous

**Table 7. Ground cover**

Tree foliar cover	0%
Shrub/vine/liana foliar cover	25-75%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

## Additional community tables

### Animal community

Mammals common to MLRA 243X include brown bear, caribou, moose, musk ox, black bear, wolf, red fox, and rodents. Many species of migratory waterfowl and shore birds nest in the abundant ponds and wetlands. Raptors include gyrfalcon, peregrine falcon, golden eagle, hawks, and owls. Arctic char and Arctic grayling are in most of the rivers. Lake trout and northern pike are common in many lakes.

### Hydrological functions

Not available.

### Recreational uses

This remote area of Alaska is mostly undeveloped wildland and is sparsely populated. The area provides excellent opportunities for hunting and other kinds of wildland recreation. Most visitors are served by air taxi, guiding, and outfitting companies operating out of the major Alaska communities. Most of the communities are along the major rivers or lakes or on the coast. Kobuk Valley National Park is located within the Western Brooks Range MLRA. Visitors to Kobuk Valley NP typically fly in and camp / backpack in the park (USDA Agriculture Handbook 296).

Local residents use this area primarily for subsistence hunting, fishing, and gathering. In the Noatak Valley, many families leave their homes in villages in the spring and spend the summers at subsistence camps gathering salmon, caribou, and various wild greens and berries on the tundra.

## **Wood products**

Not available

## **Other products**

Not available

## **Other information**

Not available

## **Inventory data references**

The vegetation modeled for this site has limited data and is considered provisional. The associated model was largely developed from NRCS staff with working knowledge of the area and literature review.

## **References**

Smith, R.D., A.P. Ammann, C.C. Bartoldus, and M.M. Brinson. 1995. An approach for assessing wetland functions using hydrogeomorphic classification, reference wetlands, and functional indices.

United States Department of Agriculture Staff. 2008. Hydrogeomorphic Wetland Classification System: An Overview and Modification to Better Meet the Needs of the Natural Resources Conservation Service.

## **Other references**

Alaska Interagency Coordination Center (AICC). 2024. <http://fire.ak.blm.gov/>

Higuera, Philip E., Melissa L. Chipman, Jennifer L. Barnes, Michael A. Urban, and Feng Sheng Hu. 2011. Variability of Tundra Fire Regimes in Arctic Alaska: Millennial-Scale Patterns and Ecological Implications. *Ecological Applications* 21, no. 8: 3211–26.

Jones, B. M., C. A. Kolden, R. Jandt, J. T. Abatzoglou, F. Urban, and C. D. Arp. 2009. Fire

behavior, weather, and burn severity of the 2007 Anaktuvuk River Tundra Fire, North Slope, Alaska. *Arctic, Antarctic, and Alpine Research* 41:309-316.

Joly, K., F. S. Chapin, and D. R. Klein 2010. Winter habitat selection by caribou in relation to lichen abundance, wildfires, grazing, and landscape characteristics in northwest Alaska. *Ecoscience* 17:321-333.

LANDFIRE. 2009. Alaska Arctic Dwarf-Shrubland - Infrequent Fire. In: LANDFIRE National Vegetation Dynamics Models. USDA Forest Service and US Department of Interior. Washington, DC.

Racine, C. H., L. A. Johnson, and L. A. Viereck. 1987. Patterns of vegetation recovery after tundra fires in northwestern Alaska, U.S.A. *Arctic and Alpine Research* 19:461-469

Racine, C. H., J. L. Allen, and J. G. Dennis. 2006 Long-term monitoring of vegetation change following tundra fires in Noatak National Preserve, Alaska.

NPS/AKRARCN/NRTR-2006/02, Arctic Network of Parks Inventory and Monitoring Program, National Park Service, Alaska Region. U.S. Department of the Interior, Washington, D.C., USA.

Schoeneberger, P.J., and D.A. Wysocki. 2012. Geomorphic Description System, Version 4.2. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, Nebraska.

Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and W.D. Broderson, editors. 2012. Field book for describing and sampling soils. Version 3.0. U.S. Department of Agriculture, Natural Resources Conservation Service.

Soil Survey Division Staff. 2017. Soil survey manual. U.S. Department of Agriculture Handbook 18.

United States Department of Agriculture. 2024. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin.  
(<https://www.nrcs.usda.gov/resources/data-and-reports/major-land-resource-area-mlra>)

United States Department of Agriculture-Natural Resources Conservation Service. 2016. U. S. General Soil Map (STATSGO2). Web Soil Survey. Available online at <http://websoilsurvey.nrcs.usda.gov>. Accessed May 8 2024.

Viereck, L.A., C.T. Dyrness, A.R. Batten, and K.J. Wezlick. 1992. The Alaska vegetation classification. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station General Technical Report PNW-GTR-286.

## Contributors

Matthew Mayer  
Blaine Spellman

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	02/08/2026
Approved by	Blaine Spellman
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

### 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

---

### 5. Number of gullies and erosion associated with gullies:

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. Average percent litter cover (%) and depth ( in):

---

15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):

---

16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

---

17. Perennial plant reproductive capability:

---