

# Ecological site group R014XG908CA

## Dry Sandy Bottom

Last updated: 09/07/2023  
Accessed: 04/26/2024

---

### Key Characteristics

None specified

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### Physiography

This ESG is found typically on flood plains and alluvial fans and have slopes of 0 to 15 percent.

### Climate

The average annual precipitation in this area is 11 to 53 inches (272 to 1,353 millimeters). This ESG is found in the drier portions of the MLRA, averaging 11-18 inches. The higher amounts of precipitation occur at the higher elevations in the area north of San Francisco. Most of the rainfall occurs as low- or moderate-intensity, Pacific frontal storms during winter. This area is very dry from midspring to midautumn. Snowfall is rare. The average annual temperature is 54 to 61 degrees F (12 to 16 degrees C). The freeze-free period averages 315 days and ranges from 265 to 365 days. It is longest near the coast, and it becomes shorter with elevation.

### Soil features

This ESG is found on sandy soils that are very deep and somewhat excessively drained that formed in alluvium from granitic sources.

Metz, a sandy, mixed, thermic Typic Xerofluvent  
Tujunga, a mixed, thermic Typic Xeropsamment

### Vegetation dynamics

This ESG covers the areas of the valleys in MLRA 14 that were at one time part of a vast complex of marshes, tidal flats, estuaries, wetlands, riparian areas, and wet meadows. The urbanized landscape in the valleys within this MLRA that exists today makes it difficult to imagine the natural landscape prior to human development.

These dry sandy bottoms were likely the sandy-textured fluventic soils related to fresh sandy deposits of moving waters in the network of mainly freshwater, and some salt marshes, rivers and streams that ran through these valleys as their seasonal and tidally influenced flood waters stretched across the flood plains and terraces in spring and deposited sediment as they receded during summer. Once the area began to be settled by humans, many of these water dominated ecosystems were drained, leveed, cleared for crops and other agriculture, and urbanized.

As this landscape was de-watered and houses and agriculture took over, the water table for many of these habitats moved deeper and deeper, creating soils that would no longer offer the available soil moisture for many of the plants that had evolved with the hydrologic function of the natural system that no longer existed. These dry sandy fans and flood plains may have remained wetter than many of the surrounding soils, due to their relation to run on and river/stream waters, increasing seasonal available water capacity. These sandier textures will dry out more rapidly through both drainage and evapotranspiration in the summer months making them less hospitable for many of the native perennial grasses and more dominated by annual grasses and forbs. Annual production will still be higher

than the ESGs in uplands, due to the available waters to these soils each growing season.

Historically, this site may have looked similar to a combination of the CWHR wet meadow and riparian classifications, however with the introduction of non-native annual grasses and the impacts from cultivation, grazing, fragmentation, continued de-watering, and human alterations such as homes and roads, this site now reflects a lower producing, dry, annual grassland where it is not converted.

Currently, where this site is not under cultivation or urban developments, it is dominated by non-native annual grasses. These include wild oats, soft chess, ripgut brome, red brome, wild barley, and foxtail fescue. Common forbs include broadleaf filaree, redstem filaree, turkey mullein, true clovers, bur clover, popcorn flower, and many others. Perennial grasses, found in moist, lightly grazed, or relic grassland areas, may include creeping wildrye and blue wildrye and areas of intermittent ponding will have sedges and occasional willows and/or coyotebrush. Species composition is also related to water availability with greater amounts of relic perennial grasses and forbs in areas of greater precipitation or soil moisture.

Information from:

John G. Kie

California Wildlife Habitat Relationships System

California Department of Fish and Game

California Interagency Wildlife Task Group

## Major Land Resource Area

MLRA 014X


Central California Coastal Valleys

## Stage

Provisional

## State and transition model

### Dry Sandy Bottom



## Reference State

This ESG represents the alluvial fans, terraces and flood plains and somewhat excessively well drained grass and grass-like dominated riparian areas and wet meadows. The single most important characteristic of a wet meadow or riparian area is its hydrology. Seasonality and reliability of yearly water inflows and outflows largely determine the vegetational stability. Therefore, these habitats exist indefinitely unless the hydrologic regimes are altered. Wet meadows and riparian sites occur where water is available for much of the growing season, following spring runoff and flooding. Hydrologically, they occupy lotic and lentic stream and river lowlands and flood plains. This ESG gets its main input flow (other than precipitation) from upstream sources. Downstream runoff is the principal output flow however evapotranspiration in the warmer months also plays a role. They are in topographic toe slopes to flat lands but have a slight slope, which permits drainage of surface water. Percolation is faster due to the permeable nature of soils.

## Reference State Community Phase

Community 1.1 This reference community phase is dominated by a mixture of perennial grasses, forbs, grass-like and willows that vary depending on what the water source dominates and where the site is located across the landscape.

## State 2 Community Phase

Community 2.1 This community represents the mix of mainly non-native annual invaders, scattered willows and coyotebrush (*Salix* and *Baccharis* spp.) and some native perennial grasses like *Leymus triticoides*, *Elymus glauca*, that are able to withstand the drying out periods that occur during the warm summer months. There may be some *Juncus* spp. and *Distichlis spicata* intermixed, depending on the soil type and location on the landscape that allows for longer water inundation and greater saline conditions.

## State 3 Community Phase

Community 3.1 - This community phase represents all the varied land uses that significantly alter this ecological site, in MLRA 14 this will primarily be urban lands and cultivation. This is an extremely varied community phase that includes all types of alterations that so significantly alter the ecological site that it is permanently changed and no longer has typical or even representative ecological dynamics.

## Transitions

T1 This transition is caused by de-watering due to urbanization, levees, channelized rivers, and irrigation needs that allowed for the invasion of non-native annual plants.

R1 This restoration pathway occurs when significant time and money inputs are focused on addressing the hydrologic functions that existed historically in order to return the water table that once supported perennial grasses and grass-like.

T2 This transition is caused by significant human alterations that remove essential topsoil horizons, alter hydrologic functions, and/or add significant inputs (such as fertilizers) that change soil chemistry and soil properties for the purposes of housing developments, urban infrastructures or intensive cropping systems and force this ecological site over a threshold and change the function and structure of this site in extensive ways.

R2 This restoration pathway occurs only when significant time and money inputs are focused on returning ecological function and hydrology.

T3 This transition is caused by significant human alterations that remove essential topsoil horizons, alter hydrologic functions, and/or add significant inputs (such as fertilizers) that change soil chemistry and soil properties for the purposes of housing developments, urban infrastructures or intensive cropping systems and force this ecological site over a threshold and change the function and structure of this site in extensive ways.

## Citations