

# Ecological site group R015XY011CAESG

## Clayey Mountains <20"ppt

Last updated: 09/07/2023  
Accessed: 04/24/2024

---

### Key Characteristics

- located on mountains
- clayey texture
- <20" ppt

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### Physiography

This ESG is typically found on gently sloping to very steep uplands and have slopes of 10 to 75 percent. Elevations typically range from 50 to 3300 feet.

### Climate

The average annual precipitation in this area ranges from 6 to 20 inches (150 to 500 millimeters). Precipitation is evenly distributed throughout fall, winter, and spring but is very low in summer. Coastal areas receive some moisture from fog in summer. Most of the rainfall occurs as low- or moderate-intensity, Pacific frontal storms during the period October to May. The average annual temperature is 52 to 65 degrees F (11 to 18 degrees C), decreasing from south to north. The freeze-free period averages 275 days and ranges from 180 to 365 days, decreasing in length with elevation and from south to north.

### Soil features

Soils in this ESG consist mainly of moderately deep to deep, well drained soils that formed in material weathered from calcareous shale and sandstone or fine-grained sandstone and shale.

The soils that represent this ESG include:

Altamont, a fine, smectitic, thermic Aridic Haploxererts  
Nacimiento, a fine-loamy, mixed, superactive, thermic Calcic Haploxerolls  
Zakme, a very-fine, smectitic, mesic Vertic Haploxerolls

### Vegetation dynamics

This ESG covers the thermic (warm) areas of the low elevation mountains that receive less than 20 inches of precipitation each year in MLRA 15 that were at one time part of a vast complex of oak savannah, chaparral, and perennial native grasslands. The fragmented and urbanized landscape in these low elevation foothill terraces within this MLRA that exists today makes it difficult to imagine the natural landscape prior to human development.

Blue oak woodlands are generally associated with warmer, or rocky, or infertile, and well-drained soils from a variety of parent materials. Blue oaks are well adapted to dry, hilly terrain where the water table is usually unavailable. The climate is Mediterranean, with mild wet winters and hot dry summers. Climatic extremes are relatively great in these woodlands, because they have a considerable geographic and elevational range. Blue oaks have an unusual tolerance of severe drought, even shedding their leaves during periods of extreme moisture stress. This survival trait contributes to its pattern of distribution, as it competes most successfully with other tree species

on drier sites.

Generally, these woodlands have an overstory of scattered trees, although the canopy can be nearly closed on better quality sites. The density of blue oaks on slopes with shallow soils is directly related to water stress. The canopy is dominated by broad-leaved trees 5 to 15 m (16 to 50 ft) tall, commonly forming open savanna-like stands on dry ridges and gentle slopes. Blue oaks may reach 25 m (82 ft) in height; the tallest tree, found in Alameda County, measured 28.7 m (94 ft) high and had a crown spread of 14.6 m (48 ft). Shrubs are often present but rarely extensive, often occurring on rock outcrops. Typical understory is composed of an extension of Annual Grassland vegetation.

Blue oak is the dominant species, comprising 85 to 100 percent of the trees present. Common associates in the canopy are coast live oak in the Coast Range, interior live oak in the Sierra Nevada, valley oak where deep soil has formed, and western juniper in the Cascade Range. In the Tehachapi and Paiute Ranges in Kern County, this habitat mixes with species from east of the mountains California juniper and single-leaf pinyon. In interior sections of the southern Coast Range, as in San Luis Obispo County, it mixes with California juniper. Associated shrub species include poison-oak, California coffeeberry, buckbrush, redberry, California buckeye, and manzanita spp. The ground cover is comprised mainly of annuals, such as brome grass, wild oats, foxtail, needlegrass, filaree, fiddleneck, and others.

Details of successional trends in this habitat type are poorly known. Succession presumably proceeds directly from annual grasslands to tree stages. Most stands of BOW exist as medium or large tree stages with few or no young blue oaks present. Therefore, only structural classes 3-5:S-D are likely to be found. Few areas can be found in California where successful recruitment of blue oaks has occurred since the turn of the century. This may be due to changes in land use; increased consumption or damage of acorns and seedlings by insects, livestock, and native animals; competition between seedlings and introduced annuals for available soil nutrients and moisture; and the absence of appropriate climatic conditions. Where germination of acorns occurs, survival and growth of the seedlings typically fail. Probably in the drier savanna-like stands, the grassland openings will simply become larger as older trees die. Live oaks may replace deciduous oaks in some areas, because their seedlings are more browse resistant. Many authorities question whether conditions will ever again support the recruitment of blue oaks needed to maintain these important woodlands.

Valid generalizations about the duration of various successional stages leading to mature stands of BOW are not possible, because adequate quantitative studies have never been done. The successional sequence probably takes at least 50 years, even on good sites. Blue oaks are relatively slow-growing, long-lived trees. Large blue oaks range in age from 153 to 390 years. Estimation of tree age based on dbh measurements is risky, however, because the dbh relationship varies tremendously depending on site quality. Moreover, height growth is extremely slow or even ceases after trees reach 65 cm (26 in) in dbh.

Information from:

California Wildlife Habitat Relationships System  
California Department of Fish and Game  
California Interagency Wildlife Task Group  
Lyman V. Ritter

## **Major Land Resource Area**

MLRA 015X  
Central California Coast Range

## **Stage**


Provisional

## **Contributors**

Kendra Moseley  
Darren Pinnegar

## State and transition model

### Clayey Mountains <20 inches ppt.



Reference State Community Pathways (Natural disturbance regime only – no management scenarios)

Community Phase 1.1 – Tree cover (oaks) (<15% cover) with scattered evergreen shrub species (chamise, ceanothus, etc.). The understory is dominated by non-native annual grasses and forbs, with limited amounts of native grasses and forbs.

Community Pathway 1.1a: The historical fire regime is approximately 70 to 200 years, with lightning being the primary ignition source. Low severity, surface fires are most typical with severity of the fire increasing with the density of shrubs in the understory. Fires result in a decrease in mostly small or young trees, most shrubs and all herbaceous cover.

Community Pathway 1.1b: Fires become more frequent (less than 10-year intervals).

Community Phase 1.2 - Dominated primarily by non-native grasses and forbs with some remnant native herbs and less than 5% tree cover. This phase also includes post-fire endemics that only surface during first few years following fire.

Community Pathway 1.2a: The shift occurs as the oaks and evergreen shrubs regenerate and begin to shade out the herbaceous species.

Community Phase 1.3 – tree cover (1-5% cover) with scattered, re-sprouting evergreen shrub species (chamise, ceanothus, etc.). The understory is dominated by non-native annual grasses and forbs, with limited amounts of native grasses and forbs.

Community Pathway 1.3a: This occurs as the coast live oak and other species becomes taller and older.

Community Pathway 1.3b: This occurs when a fire takes place before the coast live oak and other species have time to create a denser, closed canopy.

State 2

Community Phase 2.1 - This community phase represents all the varied land uses that significantly alter this ecological site, typically removing the tree cover but not always. This is an extremely varied community phase that includes all types of alterations that so significantly alter the ecological site that it is permanently changed and no longer has typical or representative ecological dynamics.

Transitions

T1 This transition is caused by repeated fires that have removed most to all of the trees and evergreen shrub seed sources and/or significant human alterations that force this ecological site over a threshold and change the function and structure of this site in extensive ways that serve mostly intensive agriculture or urban/housing developments.

R1 This restoration pathway occurs only when significant time and money inputs are focused on returning ecological function and native seed source to the site.

## Citations