

# Ecological site group F022AW012CAESG

## Deep Frigid Mountains >40"ppt

Last updated: 08/07/2023  
Accessed: 04/17/2024

---

### Key Characteristics

- mountains
- frigid temp regime
- >40" ppt
- >40" depth

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### Physiography

This ESG is found on mountains generally above 5000 ft. on variable slopes.

### Climate

The average annual precipitation is 40 to 80 inches (1,015 to 2,030 millimeters) in much of this area, and as much as 100 inches (2,540 millimeters) on the mountain peaks. The amount of precipitation increases with elevation and from south to north. Summers are dry, but there are occasional thunderstorms. Much of the winter precipitation occurs as snow. The average annual temperature is 25 to 63 degrees F (-4 to 17 degrees C), decreasing with elevation. The freeze-free period averages 205 days and ranges from 65 to 345 days, decreasing in length with elevation. It is longest at the lower elevations along the western edge of the area.

### Soil features

The soils most dominant in this ESG are very deep, somewhat excessively drained soils that formed in colluvium, alluvium, and/or till from granitoid rock or andesitic mudflows. The Tallac soils are moderately well to well drained and formed in material weathered from glacial deposits.

Badgerpass, a sandy, isotic, frigid Humic Dystroxerepts

Craneflat, a sandy-skeletal, isotic, frigid Humic Dystroxerepts

Tallac, a loamy-skeletal, mixed, superactive, frigid Humic Dystroxerepts

Windy, a medial-skeletal, amorphic, frigid Humic Vitrixerands

### Vegetation dynamics

This ESG is characterized by deep to very deep, frigid soils on mountain landforms with greater than 40 inches of precipitation. It is part of the red fir habitat type, which is typically found on frigid soils over a wide range of topography exclusive of very wet sites. Annual precipitation ranges from 1000 to 3000 mm (40 to 50 in) per year, primarily as snow that forms packs up to 5 m (15 ft) in winter. Summers are dry, limiting tree growth to seasonally available soil moisture. Small pockets of lodgepole pine also occur in wet sites scattered throughout large tracts of red fir. Similar inclusions of aspen may be found along riparian zones. Mountain meadows of various sizes, sometimes associated with small lakes, are also typical of red fir habitats. At higher elevations, red fir habitats include increasing numbers of mountain hemlock, western white pine, whitebark pine, and to a lesser extent, foxtail pine and limber pine. Noble fir is associated with red fir in the northwestern region of California.

Stand structure is typified by even-aged (established within 20-year span) groups of trees that cover several to

thousands of square meters. The cause of this pattern is probably a history of recurrent lightning fires, windthrows, and insect outbreaks acting to kill groups of trees. Natural regeneration occurs on the disturbed site following the next good cone crop. Young seedling stands are thinned by competition for soil moisture during summer. Logging is becoming a more common source of disturbance, creating larger openings on average than historic disturbances.

Mature red fir stands normally are monotypic, with very few other plant species in any layer. Heavy shade and a thick layer of duff tends to inhibit understory vegetation, especially in dense stands. To the north, in the Klamath Mountains, red fir gives way to noble fir.

After disturbance (typically logging or fire) red fir vegetation on a site proceeds through 4 seral stages: grass/forb, shrub/sapling, pole/medium tree, and large tree. The grass/forb stage occurs when red fir seedlings become established on mineral soil or shallow litter and require about 5 years to reach a height of 15 cm (6 in). Herbs, are often sparse due to competition for soil moisture on light soils. In the shrub/sapling stage, large brush fields may develop after hot wildfires and are dominated by *Ceanothus* or other shrub species for many years. The pole/medium tree stage produces dense stands of young red fir that grow slowly with little mortality for many years. In the large tree stage, subdominant trees die and add to a growing layer of duff and downed woody material, and dominant trees continue to grow for several hundred years to heights of 40 m (130 ft). Old growth stands on poor sites in the Sagehen Creek drainage of Nevada County average about 400 years old. The understory of mature stands is limited to less than 5 percent cover of shade tolerant forbs (e.g., *Chimaphila menziesii*, *Pyrola picta*).

Seral patterns are defined here for both good and poor sites. The seral pattern on good sites includes 10 years in the grass/forb stage, 20 years in the shrub/seedling stage, 80 years in the pole/medium tree stage and 110 years in the large tree stage. The pattern on poor sites includes 20 years in the shrub/seedling stage, 100 years in the pole/medium tree stage and 250 years in the large tree stage. Hence the cumulative year totals are 200 from the good site and 400 from the poor site.

California Wildlife Habitat Relationships System  
California Department of Fish and Game  
California Interagency Wildlife Task Group  
Reginald H. Barrett

## Major Land Resource Area

MLRA 022A  
Sierra Nevada and Tehachapi Mountains

### Subclasses

- F022AC003CA–Frigid-Cryic Sandy Slopes
- F022AD102CA–Cool Moraines
- F022AD103CA–Very Deep Cryic Slopes
- F022AD104CA–Colluvial Frigid Mountain Slopes
- F022AE007CA–Frigid, Sandy, Moraines And Hill Slopes
- F022AE008CA–Frigid Loamy Moraine Slopes
- F022AE013CA–Frigid, Loamy, Volcanic Mountain Slopes
- F022AE024CA–South-Facing Cryic Loamy Mountain Slopes
- F022AW012CA–Deep Frigid Mountains >40"ppt
- F022AY118NV–ABMA-PICO/ARTRV/BRMA4
- R022AD010CA–Frigid Talus Slopes
- R022AE213CA–Steep Rubbly Slope

### Correlated Map Unit Components

22823477, 22823005, 22823398, 22823004, 22823189, 22823187, 22823182, 22823018, 22823410, 22938043, 22938082, 22938076, 22938067, 22938065, 22938358, 22937066, 22936608, 22936614, 22936819, 22936822, 22937898, 22937897, 22936825, 22936829, 22938492, 22938491, 22521661, 22521802, 23015818, 23015939, 23016192, 23016088, 23015761, 23016184, 23015928, 23015913, 23016618, 23016575, 23015898, 23016434, 23015892, 23025679, 23025456, 23025973, 23026059, 23025980, 23026023, 23026028, 23026032, 23025969,

23026047, 23026894, 23026899, 23027185, 23027442, 23026940, 23026943, 23027464, 23027467, 23026991, 23027710, 23027510, 23027561, 23027563, 23027764, 23027762, 23027768, 23027769, 23027771, 23027770, 23027779, 23027576, 23027581, 23027329, 23027089, 23027932, 23028013, 23027937, 23027993, 23027992, 23027100, 23028332, 23028517, 23028165, 23028166, 23028169, 23028168, 23028524, 23028525, 23028346, 23028348, 23028185, 23028194, 23028364, 23028367, 23028370, 23028373, 23028199, 23028551, 23028201, 23028205, 23028568, 23028575, 23028143, 23028801, 23028754, 23028237, 23028238, 23028761, 23028762, 23028765, 23028763, 23028764, 23028768, 23028767, 23028766, 23028770, 23028240, 23028783, 23028598, 23028253, 23028422, 23029048, 23029049, 23028786, 23028785, 23029053, 23028893, 23029063, 23028276, 23028665, 23028667, 23028476, 23028118, 23028283, 23028507, 23028285, 23028284, 23028287, 23028286, 23028289, 23028288, 23028121, 23028480, 23028292, 23028130, 23028491, 23028492, 23028682, 23028691, 23028132, 23028497, 23028305, 23028309, 23028499, 22589687, 22589686, 22589697, 22590020, 22590023, 22590025, 22589887, 22589889, 22589797, 22589698, 22589900, 22589901, 22589810, 22589811, 22590032, 22590031, 22589824, 22589913, 22589912, 22589825, 22589826, 22589915, 22589827, 22589830, 22589721, 22589722, 22589724, 22589723, 22589725, 22589921, 22589835, 22590052, 22590055, 22589926, 22589841, 22589728, 22589729, 22589731, 22590094, 22589750, 22589852, 22589952, 22589854, 22590013, 22589782, 22589874, 22591193, 22591966, 22591702, 22591970, 22591707, 22591431, 22591440, 22591437, 22591442, 22591441, 22591447, 22591714, 22591232, 22591741, 22592003, 22591786, 22592053, 22591791, 22592064, 22591793, 22591275, 22591281, 22591287, 22591283, 22591290, 22591292, 22591167, 22591589, 22591596, 22591594, 22591598, 22591600, 22591609, 22591611, 22591815, 22591813, 22591859, 22591863, 22591308, 22591309, 22592106, 22591323, 22591325, 22595563, 22595565, 22595815, 22595818, 22595582, 22595598, 22595849, 22595622, 22595400, 22609676, 22609679, 22609683, 22610153, 22610158, 22609836, 22609847, 22609715, 22609738, 22610049, 22610047, 22610055, 22610058, 22610056, 22609767, 22609766, 22610063, 22610062, 22609906, 22609910, 23029427, 23029290, 23029186, 23029287, 23029285, 23031939, 23031938, 23031929, 23031918, 23031921, 23031924, 23031911, 23031912, 23031908, 23031905, 23031903, 23031810, 23031808, 23031594, 23031736, 23031737, 23031539, 23031538, 23031896, 23031897, 23031395, 23031394, 23031732, 23031733, 23031280, 23031411, 23031412, 23031816, 23031817, 23031598, 23031599, 23031604, 23031610, 23031265, 23031998, 23031996, 23031438, 23031439, 23031433, 23031435, 23031661, 23031660, 23032257, 23032375, 23032129, 23032203, 23032125, 23032127, 23149564, 23149558

## **Stage**


Provisional

## **Contributors**

Kendra Moseley  
Curtis Talbot  
Darren Pinnegar

## **State and transition model**

## Deep Frigid Mountains >40"ppt



### Reference State Community Pathways (Natural dynamics only – no management scenarios)

After disturbance typically logging or fire red fir vegetation on a site proceeds through 4 seral stages: grass/forb, shrub/sapling, pole/medium tree, and large tree. The grass/forb stage occurs when red fir seedlings become established on mineral soil or shallow litter and require about 5 years to reach a height of 15 cm (6 in). Herbs, are often sparse due to competition for soil moisture on light soils. In the shrub/sapling stage, large brush fields may develop after hot wildfires and are dominated by *Ceanothus* or other shrub species for many years. The pole/medium tree stage produces dense stands of young red fir that grow slowly with little mortality for many years. In the large tree stage, subdominant trees die and add to a growing layer of duff and downed woody material, and dominant trees continue to grow for several hundred years to heights of 40 m (130 ft). Old growth stands on poor sites in the Sagehen Creek drainage of Nevada County average about 400 years old. The understory of mature stands is limited to less than 5 percent cover of shade tolerant forbs (e.g., *Chimaphila menziesii*, *Pyrola picta*). Seral patterns are defined here for both good and poor sites. The seral pattern on good sites includes 10 years in the grass/forb stage, 20 years in the shrub/seedling stage, 80 years in the pole/medium tree stage and 110 years in the large tree stage. The pattern on poor sites includes 20 years in the shrub/seedling stage, 100 years in the pole/medium tree stage and 250 years in the large tree stage. Hence the cumulative year totals are 200 from the good site and 400 from the poor site.

- 1.1a This community pathway occurs following a high severity fire. A prescription of mechanical clearing and burning of slash may also produce the same results.
- 1.1b This community pathway occurs over time without fire (60+ years), as the more shade-tolerant conifers over top and shade out some of the more shade-intolerant species.
- 1.1c This community pathway occurs following a low to moderate severity fire that removes younger subcanopy conifers, some less fire tolerant trees and understory species.
- 1.2a This community pathway occurs over time without vegetation management or major disturbances.
- 1.3a This community pathway occurs over time without vegetation management or major disturbances and normal progression.
- 1.3b This community pathway occurs following a low to moderate severity fire.
- 1.4a This community pathway occurs following a high severity fire that removes everything.
- 1.4b This community pathway occurs following a moderate severity fire.

Montane chaparral species are common post-disturbance on these deeper forest soils. After disturbance (logging, fire, erosion) chaparral proliferates and may exclude conifers and other vegetation for many years. However, chaparral may facilitate the germination of red fir seedlings and other shade tolerant conifers by providing a protective cover, moderating microclimate, and improving soil conditions. Chaparral shrubs may be an essential link in forest succession by building up soil nutrient levels, especially nitrogen, to the point where trees can survive. In mature timber stands, chaparral species may senesce due to insufficient light through the canopy and are only present as a sparse understory. Thus, silvicultural practices have a strong influence on the structure of montane chaparral.

## Citations