Ecological site group DX035X02CESG05 Coconino Transition - Aridic Ustic - Limestone or Loamy Hills

Last updated: 10/25/2022 Accessed: 04/19/2024

Key Characteristics

- Coconino Transition (C)
- Site parent material is limestone or dolomite, or soil is loamy.
- Site soils are aridic ustic or within a 13-17" precipitation zone.
- Site is and/or located on a hill with slopes >15%.

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Physiography

Site is and/or located on hills with slopes >15%. Aspects tend to be northeast except valleys near Truxton Wash and Aubrey Valley.

Climate

Site soils are aridic ustic or within a 13-17" precipitation zone. Precipitation comes predominantly from monsoonal patterns during months of July, August, and September. Winter precipitation is equally predominant in the northern half of the LRU.

Soil features

Limestone or loamy soils. Site consists of gently dipping shallow residuum weathered from sedimentary rocks eroded into steep cliff faces and canyons.

Major Land Resource Area

MLRA 035X Colorado Plateau

Subclasses

- F035XF613AZ-Limestone Hills 13-17" p.z. (PIED, JUOS)
- F035XG712AZ-Limestone Hills 14-18" p.z. (JUOS, PINUS)

Correlated Map Unit Components

22391267, 22391266

Stage


Provisional

Contributors


Curtis Talbot

State and transition model

Ecosystem states



State 1 submodel, plant communities



State 1 Reference state

Pinyon-juniper with understory of shrubs and grasses.

Community 1.1 Pinion - Juniper Woodland.

Pinion - Juniper with shrubs and grasses

Community 1.2 Pinion - Juniper Woodland

Pinion - Juniper Woodland with light understory of shrubs and grasses

Pathway P1.1 Community 1.1 to 1.2

A decrease in understory due to excessive grazing and browsing.

Pathway P1.2 Community 1.2 to 1.1

Management to improve understood. It may include thinning o pinyon and juniper.

State 2 Degraded State

Loss of plant and soil resources have increased erosion.

Transition T1A State 1 to 2

Severe ecosystem degredation and loss of plant and soil resources due to excessive herbivory, drought, or disease.

Restoration pathway R2A State 2 to 1

Very long term restoration of ecosystem health.

Citations