

# Ecological site group DX035X04DESG02

## Canyon Seboyeta LRU Subset - Salty Sites subgroup

Last updated: 11/02/2022  
Accessed: 04/19/2024

---

### Key Characteristics

- Canon Seboyeta. This LRU subset drains eastward toward the Acoma Valley, and is confined to Cretaceous sedimentary parent materials. It is bounded to the west by the Mt. Taylor Volcanic field, to the north by a watershed divide, and to the east and south by a break between Cretaceous and Jurassic strata.
- Sites that occur on "upland", water-shedding landforms. Elevated terraces are included in this group.
- Sites that have saline and/or sodic soils. In these cases soils regularly have an EC > 4.0 and/or SAR > 10 or ESP > 15.

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### Physiography

Sodic and saline soils, well-drained.  
Upland fans etc.

### Soil features

Saline and sodic soils, lots of carbonates, gypsum is common. These sites are derived from shale parent materials, typically have clay loam or finer textures that result in capillary redistribution of salts, seasonally, within the profile.

### Vegetation dynamics

Use STM from the Salt Flats 126NM site

### Major Land Resource Area

MLRA 035X  
Colorado Plateau

### Subclasses

- R035XA126NM–Salt Flats

### Correlated Map Unit Components

22977262, 22975952

### Stage

Provisional

### Contributors

Curtis Talbot

**State and transition model**

**Citations**