

Ecological site group ESG1

Sandy

Last updated: 08/16/2022
Accessed: 04/25/2024

Key Characteristics

- Not flooded (hills, convex portions of piedmont slopes, broad basin floors)
- Exposed bedrock absent and few if any cobbles and stones
- Soil surface texture loamy sand to medium sandy loam, subsoil non gravelly, gypsum absent
- Soil is shallow (< 50 cm) or has a calcic, cambic, or argillic horizon within 1 m, dunes may occur but are separated by areas of finer soil texture

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Physiography

Basin floors and fan piedmonts

Climate

Typic aridic, thermic

Soil features

Sandy surface and a subsurface with increased clay or carbonates, usually sandy loam to sandy clay loam.

Vegetation dynamics

Perennial grassland, mostly black grama and dropseeds, experience encroachment and progressive dominance by mesquite associated with grass loss, accelerated wind erosion, and soil redistribution and loss.

Major Land Resource Area

MLRA 042B
Southern Rio Grande Rift

Subclasses

- R042BB012NM–Sandy, Desert Shrub
- R042BB015NM–Shallow Sandy, Desert Shrub

Correlated Map Unit Components

21842962, 21843091, 21842891, 21840028, 22054094, 22054291, 21838361, 21838238, 21838112, 21838153, 21838100, 21838098, 21838044, 21838045, 21837941, 21837778, 21837779, 21837967, 21837733, 21837985, 21837904, 21837910

Stage

Provisional


Contributors

Brandon Bestelmeyer

Curtis Talbot

State and transition model

Ecosystem states



T1a - Mesquite establishment facilitated by seed transport by cattle, bare patches > 50 cm, and relatively wet springs

T1b - Black grama is reduced below ca. 5% foliar cover by heavy grazing in drought

R1 - Shrub removal via herbicide or fire followed by black grama recovery

T2a - Black grama is reduced below ca. 5% foliar cover by heavy grazing in drought

T2b - At perennial grass cover < 5%, wind and storm events trigger deep, spreading soil erosion

T3a - At perennial grass cover < 5%, wind and storm events trigger deep, spreading soil erosion

T3b - Invasion and spread of Lehman lovegrass

T4a - Sequences of wet years lead to colonization of bunchgrasses

T6a - Drought or overgrazing leads to widespread mortality of bunchgrasses

State and community key

- I. Black grama (*Bouteloua eriopoda*) foliar cover > 5%
 - A. Honey mesquite (*Prosopis glandulosa*) foliar cover ≤ 5% ... State 1 – Black grama grassland
 - B. Honey mesquite (*Prosopis glandulosa*) foliar cover > 5% ... State 2 – Mesquite invaded black grama grassland
- II. Black grama (*Bouteloua eriopoda*) foliar cover ≤ 5%
 - A. Honey mesquite (*Prosopis glandulosa*) foliar cover ≤ 15%
 - 1 Lehmann lovegrass (*Eragrostis lehmanniana*) foliar cover ≤ all other perennial grass foliar cover ... State 3 – Bunchgrass/shrubs
 - 2 Lehmann lovegrass (*Eragrostis lehmanniana*) foliar cover all other perennial grass foliar cover ... State 5 – Exotic perennial grassland
 - B. Honey mesquite (*Prosopis glandulosa*) foliar cover > 15%
 - 1 Bunchgrass cover ≤ 10% ... State 4 – Shrubland
 - 2 Bunchgrass cover > 10% ... State 6 – Grassy shrubland

State 1

Black grama grassland



The historic plant community is dominated by black grama (*Bouteloua eriopoda*) and dropseeds (*Sporobolus flexuosus*, *S. cryptandrus*, and *S. contractus*) may be secondary dominants. Bush muhly (*Muhlenbergia porteri*) and threeawns (*Aristida* spp.) are other common grasses. Soaptree yucca (*Yucca elata*), longleaf ephedra (*Ephedra trifurca*), sand sage (*Artemisia filifolia*), and snakeweed (*Gutierrezia sarothrae*) are common shrubs. Through its high basal cover, high litter cover, and consequent low rates of erosion and high infiltration rates, black grama can regenerate through tillering and possibly by seed in some years. Overutilization of black grama and increasing rates of soil erosion and deposition when bare ground cover is high may result in an increasing relative abundance of dropseeds, threeawns, or snakeweed. It is also possible that in coarser soils, such as loamy sands, dropseeds dominate naturally. Two seasons without summer rains will also lead to black grama decline, but grasses such as dropseeds and threeawns are thought to be more sensitive to drought than black grama (Herbel et al. 1972). Snakeweed or dropseeds may become dominant within this state but black grama can recover as long as some plants persist. Gibbens and Beck (1987) and Bestelmeyer et al. (2013) provide evidence for recovery of black grama within communities dominated by dropseeds or other plants. Campbell and Bombarger (1934) indicate that black grama can recover within snakeweed-dominated grassland.

Characteristics and indicators. Black grama is highly dominant and cover is largely continuous. There is evidence of black grama reproduction by seed and stolon. There are few large bare gaps (>1 m). Litter cover is abundant. Soil stability values range from 4-6 and poorly-developed biological crusts (Pedoderm class PDB) may be common in wet periods. Mesquite (*Prosopis glandulosa*) is sparse or absent (< 5% foliar cover). Black grama foliar cover levels approaching 5%, especially with loss of plants in large patches, should be cause for concern.

Resilience management. Black grama is the key plant of this site class due to its dominance in recent historical conditions, its high forage value, and its consequent sensitivity to grazing. Shifts away from black grama dominance are due to overgrazing and/or multi-year periods of summer or spring drought, or due to an accelerated rate of honey mesquite (*Prosopis glandulosa*) seed flow into black grama grassland and the establishment and growth of these shrubs. Continuous heavy grazing leads to declines in the proportional representation of black grama declines because it is preferred by cattle over other species, particularly outside the growing season (Paulsen and Ares 1962). Once black grama is locally extirpated, it tends not to recover because reproduction is primarily vegetative via stolons (Canfield 1939, Neilson 1986). Competition for water with neighboring shrubs, particularly in drought, can also limit grass growth, survival, and reestablishment (Snyder and Tartowski 2006). Consequently, resilience of the black grama grassland state is focused on 1) grazing management to avoid substantial reductions of black grama and 2) control of mesquite shrubs when mesquite recruitment is occurring. Stocking rates and grazing periods that leave stems (stolons) long enough for new plants (ramets) to take root and adequate cover to protect soils through the spring windy season is critical. This management apparently allowed black grama populations to survive extreme drought episodes in the 1930s and 1950s that led to *B. eriopoda* losses elsewhere (Bestelmeyer et al., 2013). Timely control of mesquite using aerial or spot treatments of a foliar herbicide blend (see http://aces.nmsu.edu/pubs/_b/B822/welcome.html). Herbicide treatment should be considered when there is evidence that mesquite recruitment is occurring. Individual, large mesquite plants, however, may be desirable wildlife habitat elements.

Management interpretations

Brush management Critical values	Interpretations
> 1% mesquite cover	Spot treatments with appropriate foliar herbicides can be considered, although the presence of sparse mesquite cover may have benefits for wildlife. A large number of small plants, on the other hand, may indicate risk of transition to a shrub-invaded state

Grazing Critical values	Interpretations
BOER 35-60% foliar cover	Perennial grass production averages ca. 500 lbs/acre; maximum possible cover of black grama, which is so dominant that it may exclude other species such as dropseeds and perennial forbs that may extend the length of growing season production. Advise to leave stems (stolons) long enough for new plants (ramets) to take root and adequate cover to protect soils through the spring windy season.
BOER 15-35% foliar cover	Black grama 15-35% cover Perennial grass production is ca. 200 lbs/acre; pastures including other highly palatable species can reduce grazing on black grama during growing season, but without these species present, growing season rest is advised to avoid overgrazing black grama and promote its spread.
BOER 5-15% foliar cover	Perennial grass production is ca. 100 lbs/acre; increasing risk of extinction of black grama and persistent loss of forage base as foliar cover approaches 5%; growing season grazing only when other palatable species are present or yearlong rest advised to recover black grama to a value > 15%. Rest during wet years can lead to rapid reestablishment of this grass.

Soil health management Critical values	Interpretations
Perennial grasses > 35% foliar cover	Large canopy gaps (> 1 m) are rare and usually associated with nests of kangaroo rats (<i>Dipodomys</i> spp). Undisturbed soils can have surface aggregate stability values of > 4 and poorly-developed biological crust (PDB) pedodermis. Organic carbon is maintained at 0.3% to a 18-50 cm depth. Maintain continuous cover and few large gaps to preserve these conditions.

Perennial grasses 15-35% foliar cover	Large canopy gaps are frequent and evidence of soil redistribution (Soil Redistribution Classes 2-3) is widespread. Changes to grazing management should be considered to attain higher grass cover and greater soil stability.
Perennial grasses < 15% foliar cover	Large canopy gaps dominate the site and bare soil areas are interconnected (Resource Retention Class 4-5). Measured surface aggregate stability values are between 2-3 but subsurface values may be < 2, and poorly developed biological crusts are not widely observed and are replaced by weak physical crusts (WP) or loose soil (S) pedodermis. These patterns indicate a substantial decline in resistance to wind erosion. Changes to grazing management should be considered to attain higher grass cover and greater soil stability and surface disturbance minimized prior to the spring windy season.

State 2

Mesquite invaded black grama grassland



Black grama continues to be the dominant perennial grass, but mesquite cover is sufficiently high to compete for space and resource with grasses. Mesquite plants may be very small and difficult to detect. Although fire may kill small (< 1.5 yr old; Wright et al. 1976) mesquite, it is unlikely that fire frequencies have been sufficient to remove mesquite from a grassland once a source of mesquite propagules has been connected to a grassland. Valentine (1936) indicates that beyond a height of 1-2 feet, mesquite begins to exclude grasses from around plant bases.

Characteristics and indicators. Black grama is the dominant perennial grass and mesquite are present and usually conspicuous, exceeding 5% foliar cover when leafed out. Black grama cover may be substantial but areas around shrubs devoid of grass. Typically, large patches devoid of black grama are common. Black grama foliar cover levels approaching 5% should be cause for concern.

Resilience management. Management of black grama is similar to State 1, but there should be additional concern for the effects of competition with shrubs. Bestelmeyer et al. (2013) showed that, under heavy grazing pressure, the presence of shrubs (estimated at 7% foliar cover) increased the loss of black grama. Thus, managers might consider more conservative stocking rates in the presence of shrubs than indicated by grass production alone. Brush control using herbicides (e.g. 2,4,5-T) resulting in at least a 30% mesquite kill can result in increases in grasses (Herbel et al. 1983). Newer herbicides, such as remedy/reclaim with electrostatic application may be most effective at killing shrubs. The use of fire may also be possible in more productive environments with continuous grass cover. Wright et al. (1976) observed that honey mesquite less than 2-3 years old were killed by fire, apparently because the bud zone meristem was still exposed. Older mesquite tolerate fire or other disturbances by resprouting from the bud zone if above ground parts are destroyed or damaged.

State 3

Bunchgrass/shrubs



Bunchgrasses, including dropseeds (*Sporobolus airoides*, *S. cryptandrus*, and *S. contractus*) and threeawns (*Aristida purpurea*, *A. ternipes*) are the dominant perennial grasses and black grama is rare (<5%).

Characteristics and indicators. Black grama foliar cover is < 5% and lower than that of bunchgrasses, but total plant canopy cover may range from 60-75% where perennial grasses remain healthy. Black grama exists as isolated patches, often featuring evidence of soil aggradation within these patches and soil erosion surrounding the patches where bunchgrasses occur (soil redistribution class is 3a). The % of continuous line intercept that is gap >1 m may range from 37-56. Poorly developed biological crusts (pedoderm PDB) are rare, but average surface soil stability values range from 3-5, subsurface (2.5 cm) values are 1-2. Mesquite cover is variable, but usually > 5% and must be < 15%.

State 4 Shrubland



Mesquite are dominant and intershrub areas are typically eroded with a sparse vegetation cover of annual plants, perennial forbs, and subshrubs. On soils that are moderately deep or deep to restrictive soil horizons, soil accumulates on mesquite to form coppice dunes (or nabkas; Langford 2000) over time, which attain heights over 2 m on deeper soils. Saltbush (*Atriplex canescens*) may be common in mesquite coppice dunes. Perennial grasses, restricted to bunchgrasses, may establish and attain high cover in shrub interspaces during sequences of wet years (Peters et al., 2014). In other periods, grasses cannot establish due to soil water limitations, the instability of the substrate, and rodent and lagomorph herbivory on grass seedlings. In some cases, soil truncation exposes clay- and carbonate-rich subsoil that is very hard and does not permit plant establishment.

Characteristics and indicators. Mesquite is dominant and may form coppice dunes from 0.5-3 m high. Bunchgrasses are usually absent or restricted to coppice dunes. In extremely wet years, bush muhly, threeawns, and dropseeds may colonize interdunes. There is usually abundant evidence of wind erosion and deposition including extreme pedestalling, plant burial, highly sorted sand, ripples, and an exposed B horizon in interdunes.

Resilience management. Grazing rest during wet seasons to promote a transition to a Grassy shrubland state.

State 5

Exotic perennial grassland



Lehmann lovegrass monocultures occur, usually as large patches occurring within areas otherwise dominated by bunchgrass/mesquite states.

Characteristics and indicators. Lehmann lovegrass is the dominant perennial grass.

Resilience management. Fire may promote Lehmann lovegrass dominance, but fire is rare in typical aridic, thermic soil climates. Little is known about management practices to promote or limit its expansion.

State 6

Grassy shrubland



Abundant cover of bunchgrasses, including bush muhly, dropseeds, and threeawns in the interspaces of mesquite shrubs or coppice dunes. Grass patches may accumulate soil and litter, creating environments that favor grass establishment and persistence. Black grama is not known to colonize shrubland states.

Characteristics and indicators. Mesquite is dominant, and bunchgrass cover may be patchy or co-dominant with mesquite (up to 30% foliare cover) nearly continuous in mesquite interspaces. Soil accumulation may be observed.

Resilience management. Care should be taken to manage grazing use to promote bunchgrass survival and recruitment during dry years. Severe drought may lead to a transition back to the shrubland state irrespective of grazing management.

Transition T1a

State 1 to 2



Black grama grassland



Mesquite invaded black grama grassland

Mesquite establishment rate is thought to be accelerated by 1) increased seed availability when cattle forage in shrubby areas and then transport and defecate mesquite seeds in grass-dominated patches and 2) reduced perennial grass cover that reduces competition for surface soil water between grasses and shrub seedlings. On the other hand, Herbel and Gibbens (1996) suggest that mesquite invasion can occur within apparently intact black grama stands, suggesting that competition may be less important than seed dispersal. In addition, a reduction of historical fire frequencies associated with fragmentation of perennial grass cover has been proposed, but there is little evidence that fire can be used to limit mesquite establishment in all but the most productive landscapes due to natural limitations in fuel continuity. The factors that have led to existing variations in mesquite density within black grama grasslands are poorly understood.

Constraints to recovery. Once established, mesquite shrubs are unlikely to suffer natural mortality due to drought, herbivory, or age (Goslee et al. 2003). Mesquite can live at least 60 years on the Jornada Experimental Range in south-central New Mexico and in desert grasslands of the Santa Rita Experimental Range in southeastern Arizona, mesquite live for over 200 years (McClaran 2003). Consequently, the spread of mesquite in black grama grasslands tends to accelerate over time without mesquite control (Goolsby 2012).

Context dependence. Increases in mesquite dominance within grasslands and associated grass declines have been shown to be distance-dependent (Yao et al. 2006, Goolsby 2012). Subsoil texture also controls the rate of shrub spread and grass loss over decadal timescales (Browning et al. 2012). Soils that are deep and with lower maximum clay content (9%) in the top 100 cm experienced a much faster rate of shrub encroachment than soils that had a root-limiting horizon (a petrocalcic or "caliche" layer) and higher maximum clay content (13.5 %) within the top 100 cm.

Transition T1b State 1 to 3



Black grama grassland



Bunchgrass/shrubs

Overgrazing of black grama plants in the context of multi-year drought is known to cause black grama mortality and persistent black grama reductions or extinction from large areas (Bestelmeyer et al. 2011), which may subsequently become dominated by bunchgrasses. The loss of perennial grass cover may create opportunities for mesquite shrub establishment and gradual increases in shrub dominance (see Transition 1a).

Constraints to recovery. Black grama does not recover due to limitations in seed establishment, and because too few plants are available to recolonize areas vegetatively.

Context dependence. It is suspected that this transition is most likely on soils that are > 1 m deep to a petrocalcic (caliche) horizon and have relatively low clay content, but this has not yet been quantified.

Restoration pathway R1 State 2 to 1



Mesquite invaded black grama grassland



Black grama grassland

Mesquite can be killed by application of foliar herbicides (see http://aces.nmsu.edu/pubs/_b/B822/welcome.html). Black grama recovery into large bare patches surrounding killed shrubs can occur over time via clonal growth, or areas may be colonized initially by other grasses such as *Sporobolus* species. In wetter areas (above 10 inches mean annual precipitation), fire might be used as a tool to prevent mesquite regrowth or reestablishment. The successful use of fire in black grama grasslands, however, depends strongly upon the size of mesquite and probably on post-fire precipitation patterns that favor black grama recovery (Drewa and Havstad 2001). At this point, it is unclear if fire can be effectively used as a management tool to promote black grama dominance in most areas within MLRA 42.

Context dependence. Causes of spatial or temporal variations in the efficacy of herbicides is unknown. Fire management may only be possible in the wettest parts of this MLRA.

Transition T2a State 2 to 3



Mesquite invaded black grama grassland



Bunchgrass/shrubs

Overgrazing of black grama plants in the context of multi-year drought is known to cause black grama mortality and persistent black grama reductions or extinction from large areas (Bestelmeyer et al. 2011), which may subsequently become dominated by bunchgrasses. An increased density of shrubs in state 2 exacerbates black grama loss by constraining recovery during wet years and hastening grass mortality due to competition for soil water (Pierce et al., 2018).

Constraints to recovery. Black grama does not recover due to limitations in seed establishment, and because too few plants are available to recolonize areas vegetatively.

Context dependence. It is suspected that this transition is most likely on soils that are > 1 m deep to a petrocalcic (caliche) horizon and have relatively low clay content, but this has not yet been quantified.

Transition T2b State 2 to 4



Mesquite invaded black grama grassland



Shrubland

Loss of interspace grass cover due to heavy grazing and/or drought leads to increasing erosion and redistribution of soil to shrubs (Schlesinger et al., 1990) and out of the site (Gillette and Monger 2006). Erosion leads to loss of remaining grasses due to soil destabilization, exacerbated by increasingly concentrated rodent and livestock herbivory on grasses. Sediment deposition around mesquite shrubs leads to coppice dune formation. The factors responsible for the apparently great variation in the occurrence of this transition are not well understood.

Constraints to recovery. Accelerated wind erosion, sediment transport, and abrasion of grass seedlings is likely an important constraint to overall grass recovery, alongside seedling and grass herbivory by rodents and

lagomorphs. Exposure of infertile soil horizons at the surface may also limit grass recovery. In addition, black grama recovery from seed is rarely observed due to unknown recruitment limitations in this climatic zone.

Context dependence. Grass loss and shrub encroachment has been observed to be more rapid on weakly developed soils without significant carbonate or clay accumulation (Browning et al. 2012).

Transition T3a

State 3 to 4



Bunchgrass/shrubs



Shrubland

Similar to transition 2b

Constraints to recovery. Similar to transition 2b

Context dependence. Similar to transition 2b

Transition T3b

State 3 to 5



Bunchgrass/shrubs



Exotic perennial grassland

Lehmann lovegrass invades (usually associated with deliberately introduced seed used in roadside plantings) and becomes dominant, particularly during wet years. The circumstances under which lovegrass dominance occurs are unknown, but high late spring/early summer precipitation and warmer temperatures might promote this species.

Constraints to recovery. Lehmann lovegrass patches persist and spread, especially during wet years.

Context dependence. None yet known.

Transition T4a

State 4 to 6



Shrubland



Grassy shrubland

Sequences of wet summers in the context of limited grazing use can catalyze bunchgrass establishment, sometimes featuring multiple establishment events in growing season (Peters et al. 2014).

Constraints to recovery. The formation of bunchgrass patches leads to soil stabilization and soil and litter accumulation, which might increase seasonal soil water availability. The abundance of plants might also surpass a threshold beyond which rodent and lagomorph herbivory has reduced impact on overall grass survival and recruitment.

Context dependence. This transition is not possible on soil or soil patches that have eroded to clay-rich hardpans.

Transition T6a State 6 to 4



Grassy shrubland



Shrubland

Drought alone or overgrazing leads to widespread mortality of bunchgrasses. Soil erosion and other feedbacks constraining grass recover resume.

Context dependence. None currently known

Citations